

1 **Emergent synergistic lysosomal toxicity of chemical mixtures in**  
2 **molluscan blood cells (hemocytes)**

3

4 **Moore, MN<sup>a, 1,2, 3</sup>, Wedderburn, RJ<sup>1</sup>, Clarke KR<sup>1, 4</sup>, McFadzen, IRB<sup>1, 3</sup>, Lowe, DM<sup>1</sup>, Readman, JW<sup>1, 5</sup>**

5

6 <sup>1</sup>Plymouth Marine Laboratory, Prospect Place, The Hoe, Plymouth PL1 3DH, UK

7 <sup>2</sup>University of Exeter Medical School, European Centre for Environment & Human Health (ECEHH), Truro,  
8 TR1 3HD, UK

9 <sup>3</sup>University of Plymouth, School of Biological and Marine Sciences, Drake Circus, Plymouth PL4 8AA, UK

10 <sup>4</sup>PRIMER-C, Plymouth Marine Laboratory, Prospect Place, The Hoe, Plymouth PL1 3DH, UK

11 <sup>5</sup>University of Plymouth, School of Geography, Earth and Environmental Sciences, Drake Circus,  
12 Plymouth PL4 8AA, UK

13

14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48

## **Abstract**

The problem of effective assessment of risk posed by complex mixtures of toxic chemicals in the environment is a major challenge for government regulators and industry. The biological effect of the individual contaminants, where these are known, can be measured; but the problem lies in relating toxicity to the multiple constituents of contaminant cocktails. The objective of this study was to test the hypothesis that diverse contaminant mixtures may cause a greater toxicity than the sum of their individual parts, due to synergistic interactions between contaminants with different intracellular targets. Lysosomal membrane stability in hemocytes from marine mussels was used for *in vitro* toxicity tests; and was coupled with analysis using the isobole method and a linear additive statistical model. The findings from both methods have shown significant emergent synergistic interactions between environmentally relevant chemicals (i.e., polycyclic aromatic hydrocarbons, pesticides, biocides and a surfactant) when exposed to isolated hemocytes as a mixture of 3 & 7 constituents. The results support the complexity-based hypothesis that emergent toxicity occurs with increasing contaminant diversity, and raises questions about the validity of estimating toxicity of contaminant mixtures based on the additive toxicity of single components. Further experimentation is required to investigate the potential for interactive effects in mixtures with more constituents (e.g., 50 –100) at more environmentally realistic concentrations in order to test other regions of the model, namely, very low concentrations and high diversity. Estimated toxicant diversity coupled with tests for lysosomal damage may provide a potential tool for determining the toxicity of estuarine sediments, dredge spoil or contaminated soil.

### ***Two line capsule of paper:***

**Synergistic interactions have been observed in mixtures of toxic chemicals and relatively non-toxic chemicals increase the toxicity of the mixture. Toxicity increases with chemical diversity.**

**Key words:** *complex pollutant mixtures; effect isobole; molluscan hemocytes; lysosomal membrane stability; neutral red retention test; synergistic interactions*

***a – Corresponding author - Prof. Michael N Moore; e-mail: [mnmm@pml.ac.uk](mailto:mnmm@pml.ac.uk)***

## 49 Introduction

50 In the past, the problem of chemical contamination in the environment has generally been  
51 addressed in terms of chemical characterisation of micropollutants such as polycyclic  
52 aromatic hydrocarbons, organochlorines, pesticides (Cassee et al., 1998; European  
53 Commission, 2011; LeBlanc & Olmstead, 2004; Readman, 1996; Readman et al., 1986,  
54 1992a & b, 1993a & b; Smith et al., 2013; Tolosa & Readman, 1996; Tolosa et al., 1996,  
55 1997). However, chemical analysis only provides limited windows into a very complex  
56 mixture often believed to contain anything from 1000 to >100,000 compounds. Such  
57 analyses are very expensive and provide no direct information on harmful effects. However,  
58 effective measurement of direct toxicity *in situ* is now possible and has been applied  
59 increasingly to earthworms, fish and shellfish (Sforzini et al., 2015; Koehler et al., 1992;  
60 Lowe et al., 1992; Moore, 1988). The major difficulty has been to relate toxicity in the real  
61 environment to the chemicals believed to be present in soils, sediments, effluents and  
62 dredge spoils. At present Toxicity Identification and Evaluation (TIE) method probably offers  
63 the best option (Mount & Anderson-Carnahan, 1988). First introduced by the US  
64 Environmental Protection Agency, TIE uses various procedures to fractionate the toxins  
65 within a sample. Bioassays and high level fractionation are used to determine causative  
66 agents and quantitative high resolution GC–MS or LC–MS analysis is then used to  
67 investigate the fractions producing the greatest toxicity. However, in many environmental  
68 situations there is limited knowledge of which toxic chemicals are actually present, as well as  
69 their physical chemical speciation and bioavailability, and this can impose a level of  
70 uncertainty on attempts to predict the toxic effects on the biota and potential human health  
71 impact (e.g., through consumption of seafoods).

72

73 Assessing the harmful impact of mixtures in the environment and food is a major concern to  
74 regulators (McCarty & Borgert, 2006; Bringhoff et al., 2007; Cedergreen, 2014; Kienzler et  
75 al., 2016; Kortenkamp et al., 2009; Sarigiannis & Hansen, 2012; Tallarida, 2012, 2016; Tang  
76 et al., 2014). Attempts have been made to relate measured harmful endpoints (e.g.,  
77 pathology, mortality) to data on the toxicity of individual constituents of the mixture (Doi,  
78 1994). However, success has been limited in this respect and the hypothesis that the  
79 “toxicity of a complex mixture is simply the summation of the toxicity of its individual  
80 constituents” is now treated with some scepticism, since in this model there is no accounting  
81 for emergent interactive effects (Sahai, 1997; Fig. 1). For instance, there are numerous  
82 instances of synergistic interactions, even in simple mixtures, of drugs used in medical  
83 therapeutics (Di Dodato & Sharom, 1997; Kanazawa et al., 1997; Piras, et al., 1997;  
84 Tallarida, 2012; Valenti et al., 1997). There is also evidence for this type of emergent effect

85 in estrogenic effect of mixed pollutants (Ashby et al., 1996; Kortenkamp & Altenburger,  
86 1998), and in the synergistic interactive effects of non-toxic sucrose polyester, a zero-calorie  
87 cooking food additive and the polycyclic aromatic hydrocarbon anthracene (Moore et al.,  
88 1997).

89

90 Nevertheless, confounding factors include the lack of information about which chemicals are  
91 actually present as mentioned above, as well as their concentrations and toxicities (Smith et  
92 al., 2013). Furthermore, in a complex mixture situation where the chemicals are often  
93 associated with particle surfaces or lipid-rich coatings of particles, the probability of catalytic  
94 reactions occurring to generate new compounds will be increased, since reactions such as  
95 oxidative changes will readily take place in a two dimensional environment (i.e., surfaces)  
96 where the chemicals are highly concentrated (Fig. 1; Li et al., 2017).

97

98 In essence, this situation presents a major problem and challenge for ecotoxicology and  
99 environmental toxicology, and one that is also widely recognised in mammalian and human  
100 pharmacology and toxicology; and due of the enormous diversity of chemicals, it is very  
101 difficult to develop generalised rules that will determine the toxicity. Consequently,  
102 understanding the toxicity of chemically diverse mixtures is one of the major challenges for  
103 the future in toxicology (Cassee et al., 1998; Cedergreen, 2014; European Commission,  
104 2011; Kienzler et al., 2016; Kortenkamp et al., 2009; LeBlanc & Olmstead, 2004;  
105 Kortenkamp et al., 2009; McCarty & Borgert, 2006; Sarigiannis & Hansen, 2012; Smith et  
106 al., 2013; Tallarida, 2012, 2016).

107

108 An alternative approach to the problem of predicting the toxicity of pollutant cocktails  
109 (complex mixtures) involves treating the probable harmful impact as supracritical cascades  
110 of self-propagating chemical interactions, leading to a burst of toxic molecular diversity  
111 which results in cell and tissue damage. By trying to ascertain the laws that govern the  
112 emergence of toxic interactions in the complexity of contaminated environments (Kauffman,  
113 1993), we have to consider the types of chemical and biochemical interactions that can  
114 occur within a highly diverse molecular environment. For instance, such a situation must  
115 have prevailed in the early prebiotic history of our planet and yet it was from this diverse  
116 molecular mixture, also containing many toxic chemicals that life originated. Living  
117 organisms are highly organised molecular and supramolecular aggregations where order  
118 and structure have emerged as a direct result of this very molecular diversity (i.e., self-  
119 organised criticality), but in which, destructive toxic cascades (**supracriticality**, see Fig. 1)  
120 are prevented by protective homeostatic regulation (Bak & Chen, 1991; Kauffman, 1993).  
121 However, vestiges of the prebiotic condition are probably still represented in the universal

122 use by cells of limited toxic cascades in intracellular signalling processes (e.g., free Ca<sup>2+</sup>,  
123 oxyradicals and nitric oxide; Yermolaieva et al., 2000).

124

125 Consider then, an environment containing a diversity of toxic chemicals (e.g., a  
126 contaminated sediment or soil) and the consequences of this on the indigenous animals and  
127 plants. At some critical diversity and contaminant concentration, the protective homeostatic  
128 processes within the cells will be overwhelmed and the cells will become supracritical (e.g.,  
129 cascades of reactive free radicals) leading to cell injury and death. Hence, toxic cascades  
130 will occur at a high concentration of total contaminants where the molecular diversity is very  
131 low. What is not known is whether in a highly diverse toxic mixture, the total concentration  
132 can be very much lower (Fig. 1). Essentially, the question that is posed here is as follows:  
133 does the diversity of pollutant molecules and multiplicity of modes of action increase toxicity  
134 or are the effects generally additive?

135

136 Consequently, the hypothesis being tested is that complex mixtures of contaminant  
137 chemicals will result in a cascade of toxicity (supracriticality) if the molecular diversity rises  
138 above a critical threshold (i.e., a phase transition). Examples of analogous behaviour have  
139 been demonstrated in autocatalytic systems and proposed for the behaviour of bacterial  
140 ecosystems (Kauffman, 1993). This hypothetical model is readily testable in relation to  
141 environmental toxicity (Fig. 1).

142

143 If this hypothesis provides a satisfactory explanation for mixture toxicity, then the total  
144 concentration of pollutant chemicals in a mixture is such that it would be relatively non-toxic,  
145 or have low toxicity, for any single compound (i.e., subcritical behaviour). However, when  
146 the chemicals are combined in a mixture, they will interact in a complex manner with cellular  
147 processes to produce toxicity (i.e., supracritical behaviour), if their molecular diversity  
148 exceeds the threshold of the critical phase-transition boundary. This would be an example  
149 of emergent behaviour.

150

151 Unfortunately, additive effects at single test concentrations cannot be used reliably to test for  
152 interactive effects, and dose responses are a necessary requirement, as demonstrated by  
153 Berenbaum (1989) and Kortenkamp and Altenburger (1998). Additionally, non-additive  
154 emergent interactions can readily be identified by using the method of effect isoboles, which  
155 is reliant on the concept of concentration additivity proposed by Loewe and Muischnek  
156 (1926), and used by Kortenkamp and Altenburger (1998) to demonstrate emergent  
157 estrogenic effects. Tallarida (2012 & 2016) has recently reviewed the use of the isobole  
158 method in relation to the interactions of pharmaceuticals.

159

160 The phagocytic blood cells (hemocytes) of marine mussels (*Mytilus galloprovincialis*), a  
161 common commercially and ecologically important animal, were used as the main  
162 experimental tool in this study. Mussels are used globally as sentinels for environmental  
163 monitoring and impact assessment, hence, they provide an appropriate model for this  
164 investigation (Cheung et al., 1998; Krishnakumar et al., 1994; Moore, 1988; Widdows et al.,  
165 1992). Their blood cells are immunocytes, and hence, are a key part of the cellular or innate  
166 immune system of the mussel, and there are many functional parallels with phagocytic  
167 mammalian white blood cells. They are also known to be the target for chemical pollutant  
168 impact, which relates directly to important pathological consequences such as suppression  
169 of immune function (Galloway & Depledge, 2001; Moore et al., 2009). *In vitro* tests with live  
170 cells can be performed rapidly and in large numbers (Loizou, 2016); and experimental  
171 exposures of molluscan blood cells (hemocytes) to mixtures will be tested for evidence of  
172 additive, synergistic and antagonistic interactions. Toxic cellular reactions induced by the  
173 various experimental treatments will be measured using a lysosomal membrane stability  
174 method as an indicator of cell injury involving damage to intracellular membranes (Lowe et  
175 al., 1992; Moore et al., 1996, 2009). Lysosomal membrane stability was chosen as it is an  
176 integrated biomarker of cellular health/dysfunction, which is functionally related to protein  
177 turnover (degradation component), endocytosis, autophagy, oxidative stress and correlated  
178 with DNA damage caused by benzo[a]pyrene (Moore et al., 2006; Sforzini et al., 2018).

179

180 Finally, the primary objective of this investigation will be to test the hypothetical model  
181 described in Figure 1 relating sublethal pollutant toxicity in complex mixtures to the  
182 molecular diversity of pollutant species; and the experimental results will be used to  
183 establish the subcritical-supracritical boundary for various concentrations and combinations  
184 of chemicals. Environmental contaminants that will be tested include polycyclic aromatic  
185 hydrocarbons, pesticides, a biocide and a surfactant (Readman, 1996; Tolosa et al., 1996a  
186 & b). These chemicals have been chosen on the basis of their well documented toxicity and  
187 their continued presence in the environment in various parts of the world (Cassee et al.,  
188 1998; European Commission, 2011; LeBlanc & Olmstead, 2004; Patel et al., 2016;  
189 Readman, 1996; Readman et al., 1986, 1992a & b, 1993a & b; Sapozhnikova et al., 2013;  
190 Smith et al., 2013; Tolosa & Readman, 1996; Tolosa et al., 1996, 1997). The concentrations  
191 of test chemicals used in this study were higher than would probably be encountered in the  
192 natural environment; however, the aim of the investigation was to demonstrate “proof of  
193 principle”.

194

195 **Materials and methods**

196 **Animal husbandry**

197 Blue mussels (*Mytilus galloprovincialis*, 40-50 mm shell length) were collected from Freathy  
198 Beach (Whitsand Bay, Cornwall; Grid ref: SX 39390 52066); and held for 24 hours without  
199 food in a seawater aquarium system at  $15 \pm 1^\circ\text{C}$  and 34 psu salinity with natural daylight  
200 prior to harvesting the blood cells (hemocytes).

201

202 **Lysosomal membrane stability (neutral red retention – NRR test)**

203 Lysosomal stability was assessed in the hemocytes or blood cells of mussels using neutral  
204 red as described by Lowe (1995) and Moore et al. (2008). Briefly, approximately 50  $\mu\text{l}$  of  
205 haemolymph was removed from the posterior adductor muscle of mussels ( $n = 20$  for each  
206 set of test concentrations) and added to 50  $\mu\text{l}$  ml of physiological saline (0.02 M HEPES, 0.4  
207 M NaCl, 0.1 M  $\text{MgSO}_4$ , 0.01 M KCl, 0.01 M  $\text{CaCl}_2$ , pH 7.3). 40  $\mu\text{l}$  of cell suspension was  
208 aliquoted onto a microscope slide and left in a dark moisture chamber at  $15^\circ\text{C}$  for 15 minutes  
209 to allow the cells to adhere, following which the cells were incubated in the test treatment  
210 solutions (see below). A stock solution of 100 mM neutral red in DMSO (28.9 mg of neutral  
211 red in 1 ml of DMSO) was prepared and stored in a refrigerator prior to use. However, the  
212 solution will solidify in the refrigerator and must be raised to room temperature for dilution in  
213 physiological saline to the working strength solution. The saline containing neutral red  
214 comprised 10  $\mu\text{l}$  of stock neutral red in DMSO in 5ml of mussel physiological saline. 40  $\mu\text{l}$  of  
215 neutral red saline solution was added to the slides and left for 15 min in a dark moist  
216 environmental chamber at  $15^\circ\text{C}$  to allow the neutral red to enter the cells and accumulate in  
217 the lysosomes. The slides were maintained under these conditions for the duration of the  
218 test, with slides only being removed briefly for microscopical examination before being  
219 returned. The cells were examined microscopically after 15, 30, 60, 90, 120, 150 and 180  
220 minutes. The test was terminated after 180 minutes, since the neutral red itself becomes a  
221 toxic xenobiotic stressor. The end point of the test was when  $> 50\%$  of the cells, based on a  
222 visual determination, exhibited lysosomal leakage of neutral red dye into the cytoplasm or  
223 showed significant abnormalities such as lysosomal enlargement (Lowe et al 1992, Moore et  
224 al., 2009).

225

226 **Experimental treatments**

227 The chemicals tested were phenanthrene, anthracene, lindane, malathion, irgarol-1051, *cis*-  
228 permethrin and sodium dodecylbenzene sulphonate (LAS). Dimethyl sulphoxide (DMSO)  
229 was used as an initial solvent to prepare the 100 mM stock solutions for the test compounds,  
230 with the exception of LAS which was water miscible. Exposures were conducted at 10, 50,  
231 and 100  $\mu\text{M}$  in physiological saline for 30 minutes at  $15^\circ\text{C}$ . Various test mixtures (Mix 2 –  
232 phenanthrene + anthracene; Mix 3 - phenanthrene + anthracene + LAS; and Mix 7 – all

233 seven test compounds) were made up to each of these concentrations with all of the  
234 constituents being an equal proportion of the final dose. The controls were actually vehicle  
235 controls with the equivalent concentration of DMSO (0.1 % v/v) in physiological saline in all  
236 cases. DMSO at the concentration used is non-toxic to mussels; and the NRR values for the  
237 vehicle controls were within the normal range (Banni et al., 2017; Bellas et al., 2005, 2006).

238

239 Microscope slides with attached hemocytes (see section above) were then shaken to  
240 remove the excess haemolymph and the slides placed into 50 ml Coplin jars containing the  
241 test chemicals dissolved in physiological saline. Slides were incubated in the Coplin jars, in  
242 the absence of light for a further 30 minutes, at a constant temperature of 15°C. Slides were  
243 then removed from the coplin jar, drained and 40µl of the neutral red working solution was  
244 added (see section above). All slides were coded to prevent operator bias and only decoded  
245 after the analyses of all samples were complete. The results were expressed as lysosomal  
246 toxicity (i.e., 100% - NRR [*lysosomal membrane stability*] as a % of control ± 95% CL).

247

#### 248 **Isobole method**

249 50% toxicity values were determined for the individual compounds and the mixtures and the  
250 sum of concentration additivity for the 50% isobole calculated from the generic formula  $d_a/D_a$   
251  $+ d_b/D_b$  ..... where  $d_a$  and  $d_b$  are the doses/concentrations of A and B in a mixture that  
252 produces a specified effect (50% toxicity) and  $D_a$  and  $D_b$  are the doses/concentrations of the  
253 single compounds, which on their own elicit the same effect as the mixture (Kortenkamp &  
254 Altenburger, 1998). 95% confidence bands were generated for the determination of the  
255 confidence limits (± 95%) for the 50% isoeffective concentrations and; and a pooled variance  
256 estimate was used to determine the estimated 95% confidence limits for the additivity  
257 concentrations (Cohen, 1988; Kortenkamp & Altenburger, 1998). 84% confidence limits were  
258 employed on the graphical plots rather than 95% confidence intervals, it then being true (for  
259 large n, as here) that non-overlapping intervals correspond to a significant difference in a 5%  
260 level test (Buzatto et al., 2015).

261

#### 262 **Linear-additive statistical method**

263 A conventional statistical modelling approach was also employed, to examine robustness of  
264 the conclusions to choice of predictive model. Rather than the threshold-based approach of  
265 supracritical cascades this postulates a simple linear additive structure for the dose-  
266 response models. Computations are performed on the raw NRR data, rather than as  
267 expressed by a percentage of controls, to preserve the statistical independence in formal  
268 inference from standard linear models and to allow visualisation of NRR levels under control  
269 conditions (higher NRR denotes lower toxicity). Observed NRR means and confidence



270 intervals for the 9 mixture experiments (3 mixtures at 3 concentrations, n = 20 replicates  
271 each) are contrasted with predictions from an appropriate linear combination of NRR  
272 estimates from the separate regressions for each of the 7 compounds. Confidence intervals  
273 for these predictors were based on standard errors computed under the usual rules for  
274 variance of a linear combination of independent random variables. Formal testing of a  
275 difference between a predicted and observed mean relied on standard normality  
276 assumptions, justifiable here by the central limit theorem since the two statistics are each  
277 means over a large number of observations. For this model, under the null hypothesis of no  
278 effect of toxicant diversity, only of toxicant concentration, predicted and observed means will  
279 not differ. These test results are conveniently visualised on means plots by employing 84%  
280 rather than 95% confidence intervals, it then being true (for large n, as here) that non-  
281 overlapping intervals correspond to a significant difference in a 5% level test (Buzatto et al.,  
282 2015).

283

284 Non-parametric Kruskal-Wallis tests were also used on the lysosomal toxicity data, to  
285 provide additional robustness to test conclusions.

286

## 287 **Chemicals**

288 All chemicals were obtained from Sigma-Aldrich. Anthracene and phenanthrene were  $\geq 99\%$   
289 pure; LAS (dodecylbenzene sulphonate) was Pharmaceutical Secondary Standard -  
290 Certified Reference Material; pesticides and herbicides were analytical standard grade;  
291 DMSO ( $\geq 99.9\%$ ); and neutral red powder (N4638) was graded as suitable for cell culture.  
292 All other reagents used were of ANALAR grade.

293

## 294 **Results**

### 295 **Lysosomal toxicity**

296 Effects of the test chemicals and the 3 mixtures on % lysosomal toxicity (100% - NRR  
297 [*lysosomal membrane stability*] as a % of control  $\pm$  95% CL) were measured at three  
298 concentrations (10, 50 & 100  $\mu\text{M}$ ; Fig. 2). Lysosomal toxicity was also plotted as a series of  
299 dose responses for all of the test conditions (Fig. 2A - H), as these were required to  
300 determine the 50% effect isoboles.

301

302 Dose/concentration responses for the individual test chemicals showed that LAS had the  
303 greatest effect on lysosomal toxicity (Fig. 2C). Phenanthrene and anthracene also had a  
304 significant effect on the lysosomal toxicity ( $p < 0.05$ ,  $n = 20$ ; Kruskal-Wallis test; Fig. 2A, B).  
305 Malathion showed slight toxicity, but only at 10  $\mu\text{M}$ ; while lindane, irgarol-1051, and *cis*-

306 permethrin caused no significant effect on lysosomal retention ( $p > 0.05$ ,  $n = 20$ ; Kruskal-  
307 Wallis test; Fig. 2D - G). Controls maintained high retention times throughout all the  
308 exposures and no significant differences could be found ( $p > 0.05$ ; Kruskal-Wallis test). The  
309 mixtures of test compounds caused a significant change in lysosomal toxicity for the mixture  
310 of phenanthrene and anthracene (Mix 2) at 100  $\mu\text{M}$  ( $p < 0.05$ ,  $n = 20$ ; Kruskal-Wallis test;  
311 Fig. 2 H). However, significant increases in lysosomal toxicity were observed at 10, 50 and  
312 100  $\mu\text{M}$  for the mixture of phenanthrene, anthracene and LAS (Mix 3), as well as in the  
313 mixture of all 7 test chemicals (Mix 7;  $p < 0.05$ ,  $n = 20$ ; Kruskal-Wallis test; Fig. 2H). When  
314 the dose responses for the individual compounds were compared with the mixtures, the  
315 mixture with 7 chemicals (Mix 7) was the most toxic (Fig. 2H), and the mixture with 3  
316 components (Mix 3) was comparable to the dose response for LAS (Fig. 2C, H).

317

### 318 **Application of isobole method**

319 Isoeffective concentration values were determined for the individual compounds and the  
320 mixtures; and the sum of concentration additivity for the 50% isobole was calculated for each  
321 mixture using the generic formula described by Kortenkamp & Altenburger (1998) (Fig. 3).  
322 Anthracene did not quite achieve 50% lysosomal toxicity anthracene, and was extrapolated  
323 slightly beyond the 100  $\mu\text{M}$  concentration (Fig. 2B). Some of the test compounds (i.e.,  
324 malathion, *cis*-permethrin, irgarol 1051 and lindane) did not achieve 50% lysosomal toxicity,  
325 and also, were not significantly different from the control (Fig. 2D - G). Consequently, these  
326 compounds could not be included in the 50% concentration additivity calculations. 95%  
327 confidence bands were generated for the determination of the confidence limits ( $\pm 95\%$ ) for  
328 the 50% isoeffective concentrations; and a pooled variance estimate was used to determine  
329 the estimated 95% confidence limits for the additivity concentrations (Cohen, 1988;  
330 Kortenkamp & Altenburger, 1998).

331

332 The isobole method demonstrated that there was an additive toxic effect with a mixture of  
333 phenanthrene and anthracene (Mix 2) at the 50% effect isobole ( $P < 0.05$ ,  $n = 20$ ; Fig. 3A,  
334 B). When LAS was added to phenanthrene and anthracene (Mix 3), there was a significant  
335 synergistic interactive effect for the observed isoeffective concentration at the 50% effect  
336 isobole (Fig. 3A, B; Kortenkamp and Altenburger, 1998). With a mixture of all seven  
337 compounds (Mix 7), there was a significant synergistic interaction for the observed  
338 isoeffective concentration at the 50 % effect isobole ( $P < 0.05$ ,  $n = 20$ ; Fig. 3A, B).

339

### 340 **Application of linear additive model**

341 The linear additive statistical model allows comparison of observed with predicted effects on  
342 lysosomal membrane stability in the absence of any effect of toxicant diversity (Fig. 5). Major

343 declines in NRR (i.e., greater toxicity) were seen for Mixtures 3 & 7 that cannot be explained  
344 solely by toxicant concentration under this linear additive model (Fig. 4); and these findings  
345 were indicative of synergistic interactions (Fig. 3A, B). These results have to be treated with  
346 caution as the dose responses for phenanthrene, anthracene and LAS were not strictly  
347 linear. However, the results from the linear additive model are in very good agreement with  
348 the findings from the use of the isobole method.

349

### 350 **Modelling**

351 The enhanced toxic effects were evident with the most diverse toxic mixture (Mix 3 & 7),  
352 when the data for the mixtures were plotted as % lysosomal toxicity (100% – NRR as % of  
353 control) against the diversity (i.e., number of test compounds) of the toxic mixture (Fig. 5).  
354 Results shown in Figure 6 tentatively indicated that there may be a phase shift in relative  
355 toxicity with the test mixtures with 3 & 7 components.

356

357 Finally, the lysosomal toxicity data was used to test the concentration & diversity model  
358 proposed in the hypothesis. The results are presented in the log concentration versus log  
359 chemical diversity matrix shown in Figure 6 that indicates that the hypothesis being tested is  
360 probably supported by the data. These findings are in agreement with the results of the 50%  
361 effect isobole method, and the linear additive statistical model, that synergistic interactions  
362 are in fact occurring in the more complex mixtures.

363

### 364 **Discussion**

365 Living cells as the basic units of life operate below or near to the subcritical-supracritical  
366 boundary, sometimes referred to as “the edge of chaos” (Kaufmann, 1993; Fig. 1). If cells  
367 were supracritical then the introduction of any foreign molecule (i.e., xenobiotic chemical)  
368 will probably unleash a potentially harmful and reactive molecular cascade (Kauffman,  
369 1993). Such a reactive cascade will propagate, since each new molecule can potentially  
370 interact with another biological molecule, such as membrane lipids, proteins and DNA and  
371 potentially initiate a further cascade (i.e., a chain reaction). Therefore, there is a high  
372 probability that many of these cascading toxic molecules will perturb the homeostatic  
373 regulation of cellular processes (i.e., sublethal toxicity) and lead to cell injury, pathology and  
374 cell death. Essentially, supracritical conditions within cells will be lethally destructive.  
375 However, cells have evolved a number of protective processes in order to protect  
376 themselves. These protective systems include sequestration within the interior  
377 microenvironment of a membrane-bound vesicle (e.g., the lysosome), membrane pumps to  
378 remove novel molecules from the cell (e.g., multidrug-resistance system - MDR), detoxifying

379 enzymes to metabolise toxic molecules (e.g., cytochromes P-450 [CYP superfamily] and  
380 esterases) and antioxidant enzymes to protect against free radicals (e.g., superoxide  
381 dismutase, catalase and glutathione peroxidase), as well as scavenging molecules that bind  
382 to toxic reactive molecules (Minier & Moore, 1996a, b; Sies, 1997).

383

384 Lysosomes are central to innate or cellular immune function, and normal turnover of cellular  
385 proteins and other biomolecules (Moore et al., 2004, 2006, 2007). Lysosomes are well  
386 established targets for many environmental xenobiotic chemicals that are also known to  
387 accumulate in lysosomes (i.e., metal ions and organic chemicals with many modes of toxic  
388 action; Moore et al., 2004; Rashid et al., 1991). Furthermore, the lysosomal vacuolar system  
389 has an important cellular protective function and, when lysosomal storage capacity is  
390 overloaded, the lysosomes display characteristic low membrane stability reactions to toxic  
391 injury (Minier & Moore, 1996a, b; Moore, 1985, 1986, 1990; Moore et al., 1996, 2004, 2006,  
392 2007).

393

394 Lysosomal integrity is an effective measure of integrated physiological function and,  
395 therefore, is more functionally relevant than many other biomarker tests that only measure a  
396 change in the level or function of a particular protein (Lowe et al., 1992; Moore et al., 2004,  
397 2006a; Regoli, 1998; Ringwood et al., 1992; Sforzini et al., 2015, 2017). The lysosomal  
398 system also has the propensity for accumulating many xenobiotic chemicals including  
399 polycyclic aromatic hydrocarbons (de Duve, 1974; Minier & Moore, 1996a, b; Moore et al.,  
400 1996, 2004, 2006a; Rashid et al., 1991). Only some of the compounds tested were  
401 lysosomotropic: these were anthracene and phenanthrene (Moore et al., 2006). Of the other  
402 chemicals, LAS is a detergent (surfactant) and disrupts cell membranes, while lindane, *cis*-  
403 permethrin and malathion are neurotoxic pesticides and irgarol 1051 is a herbicide (Patel et  
404 al., 2016; Sapozhnikova et al., 2013). The effects of the pesticides and irgarol 1051 have  
405 been tested in several species of mussel and found to have relatively low toxicity (Bellas,  
406 2006; Bringhoff et al., 2007; Khessiba et al., 2005; Lehtonen & Leiniö, 2003).

407

408 Four of the compounds tested in several species of mussel (i.e., malathion, lindane, *cis*-  
409 permethrin and irgarol 1051) had either very low or else no lysosomal toxicity and this is  
410 supported by other studies (Bellas, 2006; Bringhoff et al., 2007; Khessiba et al., 2005;  
411 Lehtonen & Leiniö, 2003; Fig. 2). It was, therefore, surprising that these relatively low toxicity  
412 chemicals, when combined with the two PAHs and LAS, apparently contributed to the  
413 enhanced lysosomal toxicity in the hemocytes (Smith et al., 2013; Figs. 2H, 3A). However,  
414 this type of effect has been observed previously in mice exposed to a mixture of estrogenic

415 methoxychlor and non-estrogenic dieldrin (Ashby et al., 1997; Kortenkamp & Altenburger,  
416 1998; Smith et al., 2013).

417

418 When the results from the isobole and linear additive method for interactions in the mixtures  
419 (Mix 2, Mix 3 & Mix 7) were applied to the concentration/diversity model, there was a clear  
420 demarcation between those test conditions showing evidence for a synergistic interaction  
421 and those with no interaction (Fig. 6). The 50% isoeffective concentrations for the three  
422 mixtures, with transposed axes from Figure 3B, were used to generate the subcritical-  
423 supraceutical boundary curve (Fig. 6). Consequently, the hypothesis is apparently supported  
424 by the data that increasing diversity in a mixture of contaminants contributes to emergent  
425 toxicity (Kauffman, 1993).

426

427 There is some indication that a phase shift is occurring at the level of the mixture with 3  
428 compounds leading to emergent synergistic toxicity as indicated in Figure 5; and this is  
429 supported by the evidence for increased variance for the isoeffective concentration of  
430 Mixture 3 (Fig. 3A, B). Although it is probable that in other combinations of chemical  
431 compounds, this type of phase transition will occur at a different level of chemical diversity; it  
432 is reasonable to argue that a phase transition will generally indicate that emergent behaviour  
433 is happening (Kauffman, 1993; Kortenkamp et al., 2009). Nevertheless, the consequences of  
434 synergy between contaminants in the natural environment may be greater than those  
435 observed in this investigation, under relatively simple conditions over a short time period.  
436 Hence, in more chronic conditions it is possible that the interaction between contaminants  
437 may be of much greater concern for animal health status (Doi, 1994).

438

439 Ideally we would have wished to explore the effects of a larger number of combinations of  
440 these chemicals, however, this was not logistically possible within the scope of the  
441 investigation. Further experimentation is required for investigating the potential for interactive  
442 effects in mixtures with more constituents (e.g., 50–100). Such investigation will provide  
443 data sets for testing regions of the current model with very low environmentally realistic  
444 concentrations and very high diversity (Fig. 6; Cedergreen, 2014).

445

## 446 **Conclusions**

447 The fact that the results support the complexity-based hypothesis that there is emergent  
448 toxicity with increasing contaminant diversity should perhaps urge a cautionary attitude to  
449 disposal of toxic mixtures until the hypothesis is disproved. However, if further support for  
450 the hypothesis is forthcoming, then serious questions arise as to the validity of estimating  
451 potential emergent toxicity of complex mixtures based on the additive toxicity of single

452 components. Regulation of discharges based on toxicity rather than chemical composition  
453 will probably provide a more practical solution to this problem, where the toxicity of the main  
454 components has been determined by bioassays or ecotoxicity tests (Cassee et al., 1998;  
455 Cedergreen, 2014; European Commission, 2011; Kienzler et al., 2016; Kortenkamp et al.,  
456 2009; LeBlanc & Olmstead, 2004; McCarty & Borgert, 2006; Sarigiannis & Hansen, 2012;  
457 Smith et al., 2013; Tallarida, 2012, 2016; Tang et al., 2014).

458

459 **Acknowledgements:** This research is part of the PRETOCS Project supported by the Defence  
460 Evaluation and Research Agency (DERA, UK) Contract No. HO/TD/113/1), the PREDICT 2 Project  
461 supported by the Department for Environment, Food and Rural Affairs (Defra, UK), Contract No.  
462 AE1136 & the Natural Environment Research Council (NERC, UK). There are no conflicts of interest  
463 in the research reported here.

464

## 465 References

- 466 Ashby, J., Lefevre, P.A., Odum, J., Harris, C.A., Routledge, E.J., Sumpter, J.P. (1997). Synergy  
467 between synthetic oestrogens? *Nature*, 385: 494.
- 468 Bak, P. & Chen, K. (1991). Self-organized criticality. *Scientific American*, 264: 26-33.
- 469 Banni, M., Sforzini, S., Arlt, V.M., Barranger, A., Dallas, L.J., Oliveri, C., Aminot, Y., Readman, J.W.,  
470 Moore, M.N., Viarengo, A., Jha, A.N. (2017). Assessing the impact of benzo[a]pyrene on marine  
471 mussels: use of a novel targeted low density microarray complementing biomarkers data. *PLOS*  
472 *ONE*. 2017 Jun 26;12(6):e0178460. doi: 10.1371/ journal.pone.0178460.
- 473 Bellas, J. (2006). Comparative toxicity of alternative antifouling biocides on embryos and larvae of  
474 marine invertebrates. *Sci. Total Environ.*, 367: 573–585.
- 475 Bellas, J., Granmo, Å., Beiras, R. (2005). Embryotoxicity of the antifouling biocide zinc pyrithione to  
476 sea urchin (*Paracentrotus lividus*) and mussel (*Mytilus edulis*). *Mar. Poll. Bull.*, 5: 1382–1385.
- 477 Berenbaum, M.C. (1989). What is synergy? *Pharmacol. Rev.*, 41: 93-141.
- 478 Bringolf, R.B., Cope, W.G., Barnhart, M.C., Mosher, S., Lazaro, P.R. and Shea, D. (2007), Acute and  
479 chronic toxicity of pesticide formulations (atrazine, chlorpyrifos, and permethrin) to glochidia and  
480 juveniles of *Lampsilis siliquoidea*. *Environ. Toxicol. Chem.*, 26: 2101–2107.
- 481 Buzatto, B.A., Kotiaho, J.S., Tomkins, J.L., Simmons, L.W. (2015). Intralocus tactical conflict: genetic  
482 correlations between fighters and sneakers of the dung beetle *Onthophagus taurus*. *J. Evolut.*  
483 *Biol.*, 28: 730-738.
- 484 Cassee, F.R., Groten, J.P., van Bladeren, P.J., Feron, V.J. (1998). Toxicological evaluation and risk  
485 assessment of chemical mixtures. *Crit Rev Toxicol.*, 28: 73-101.
- 486 Cedergreen, N. (2014). Quantifying Synergy: A Systematic Review of Mixture Toxicity Studies within  
487 Environmental Toxicology. *PLoS ONE*, 9(5), e96580. [http://doi.org/10.1371/ journal.pone.0096580](http://doi.org/10.1371/journal.pone.0096580).
- 488
- 489 Cheung, V. V., Wedderburn, R. J., Depledge, M. H. (1998). Molluscan lysosomal responses as a  
490 diagnostic tool for the detection of a pollutant gradient in Tolo Harbour, Hong Kong. *Mar. Env.*  
491 *Res.* 46: 237-241.
- 492 Cohen, J. (1988). Statistical power analysis for the behavioural sciences New York. NY: *Academic*.
- 493 DeDuve, C.T., DeBary, B., Poole, A., Trouet, P., Tulkens, A., Van Hoof, F. (1974). Lysosomotropic  
494 agents. *Biochem. Pharmacol.* 23: 2495-2531.
- 495 Di Diodato, G. & Sharom, F.J. (1997). Interaction of combinations of drugs, chemosensitisers, and  
496 peptides with the P-glycoprotein multidrug transporter. *Biochem. Pharmacol.*, 53: 1789-1797.
- 497 Doi, J. (1994). Complex mixtures. In: Handbook of Ecotoxicology, vol. 2 (Ed. P. Calow), pp. 289-310.  
498 Blackwell Scientific, Oxford, London, Boston.
- 499 European Commission (2011). Toxicity and Assessment of Chemical Mixtures - [ec.europa.eu/health/](http://ec.europa.eu/health/scientific_committees/environmental_risks/.../scher_o_155.pdf)  
500 [scientific\\_committees/environmental\\_risks/.../scher\\_o\\_155.pdf](http://ec.europa.eu/health/scientific_committees/environmental_risks/.../scher_o_155.pdf)
- 501 Galloway, T.S. & Depledge, M.H. (2001). Immunotoxicity in invertebrates: measurement and  
502 ecotoxicological relevance. *Ecotoxicol.*, 5: 10-23.
- 503 Kanzawa, F., Nishio, K., Fukuoka, K., Fukuda, M., Kunimoto, T. Saijo, N. (1997). Evaluation of  
504 synergism by a novel three-dimensional model for the combined action of cisplatin and etoposide  
505 on the growth of a human small-cell lung-cancer cell line, SBC-3. *Int. J. Cancer*, 1997, 71: 311-  
506 319.

507 Kauffman, S.A. (1993). *The Origins of Order*. Oxford University Press, Oxford, New York, 709p..

508 Kienzler, A., Bopp, S.K., van der Linden, S., Berggren, E., Worth, A. (2016). Regulatory assessment

509 of chemical mixtures: Requirements, current approaches and future perspectives. *Regulat.*

510 *Toxicol. Pharmacol.*, 80: 321-334.

511 Kortenkamp, A., Altenburger, R. (1998). Synergisms with mixtures of xenoestrogens: a reevaluation

512 using the method of isoboles. *Sci. Total Environ.*, 221: 59-73.

513 Kortenkamp, A., Backhaus, T., Faust, A. (2009). State of the art report on mixture toxicity. Final

514 Report to the European Commission under Contract 070307/2007/485103/ETU/D.1. Brussels,

515 Belgium.

516 Krishnakumar, P.K., Casillas, E., Varanasi, U. (1994). Effect of environmental contaminants on the

517 health of *Mytilus edulis* from Puget Sound, Washington, USA. I. Cytochemical measures of

518 lysosomal responses in the digestive cells using automatic image analysis. *Mar. Ecol. Prog. Ser.*

519 106: 249-261.

520 LeBlanc, G.A., Olmstead, A.W. (2004). Evaluating the toxicity of chemical mixtures. *Environ Health*

521 *Perspect.*, 112: A729–A730.

522 Lehtonen, K.K., S. Leiniö , S. (2003). Effects of exposure to copper and malathion on metallothionein

523 levels and acetylcholinesterase activity of the mussel *Mytilus edulis* and the clam *Macoma*

524 *balthica* from the northern Baltic Sea. *Bull. Environ. Contam. Toxicol.*, 71: 489-496.

525 Li, H., Xiao, J., Fu, Q., Bao, X. (2017). Confined catalysis under two-dimensional materials. PNAS,

526 114: 5930–5934.

527 Loewe, S., Muischnek, H. (1926). Über kombinationswerken. 1. Mitteilung: hilfsmittel der

528 fragestellung. *Naunyn-Schmiedebergs Arch. Exp. Pathol. Pharmacol.*, 114: 313-326.

529 Loizou, G. D. (2016). Animal-Free Chemical Safety Assessment. *Front. Pharmacol.*, 7: 218. doi:

530 10.3389/fphar.2016.00218

531 Lowe, D.M., Moore, M.N., Evans, B.M. (1992). Contaminant impact on interactions of molecular

532 probes with lysosomes in living hepatocytes from dab *Limanda limanda*. *Mar. Ecol. Prog. Ser.*

533 91: 135-140.

534 Lowe, D.M., Pipe, R.K. (1994). Contaminant induced lysosomal membrane damage in marine mussel

535 digestive cells: an *in vitro* study. *Aquat. Toxicol.* 30: 357-365.

536 Lowe, D.M., Fossato, V.U., Depledge, M.H. (1995a). Contaminant induced lysosomal membrane

537 damage in blood cells of mussels *M. galloprovincialis* from the Venice Lagoon: an *in vitro* study.

538 *Mar. Ecol. Prog. Ser.* 129: 189-196.

539 Lowe, D.M., Soverchia, C., Moore, M.N. (1995b). Lysosomal membrane responses in the blood and

540 digestive cells of mussels experimentally exposed to fluoranthene. *Aquat. Toxicol.* 33: 105-112.

541 McCarty, L.S., Borgert, C.J. (2006). Review of the toxicity of chemical mixtures: theory, policy, and

542 regulatory practice. *Regulat. Toxicol. Pharmacol.*, 45: 119–143.

543 Minier, C. Moore, M.N. (1996a). Induction of multixenobiotic resistance in mussel blood cells. *Mar.*

544 *Environ. Res.*, 42: 389-392.

545 Minier, C., Moore, M.N. (1996b). Rhodamine B accumulation and MXR protein expression in mussel

546 blood cells: effects of exposure to vincristine. *Mar. Ecol. Prog. Ser.* 142: 165-173.

547 Moore, M.N. (1985). Cellular responses to pollutants. *Mar. Pollut. Bull.*, 16: 134-139.

548 Moore, M.N. (1988). Cytochemical responses of the lysosomal system and NADPH-

549 ferrihemoprotein reductase in molluscs to environmental and experimental exposure to

550 xenobiotics. *Mar. Ecol. Prog. Ser.*, 46: 81-89.

551 Moore, M.N. (1990). Lysosomal cytochemistry in marine environmental monitoring. *Histochem. J.* 22:

552 189-191.

553 Moore, M.N., Allen, J.I. & McVeigh, A. (2006). Environmental prognostics: an integrated model

554 supporting lysosomal stress responses as predictive biomarkers of animal health status. *Mar.*

555 *Environ. Res.*, 61: 278–304.

556 Moore, M.N., Depledge, M.H., Readman, J.W. & Leonard, P. (2004). An integrated biomarker-based

557 strategy for ecotoxicological evaluation of risk in environmental management. *Mutat. Res.*, 552:

558 247-268.

559 Moore, M.N., Koehler, A., Lowe, D. & Viarengo, A. (2008). Lysosomes and autophagy in aquatic

560 animals. In: *Methods in Enzymology* (D.Klionsky, Ed), vol. 451, pp. 582-620. Academic

561 Press/Elsevier, Burlington.

562 Moore, M.N., Lowe, D.M., Soverchia., Haigh, S.D., Hales, S.G. (1997). Uptake of a non-calorific,

563 edible sucrose polyester oil and olive oil by marine mussels and their influence on uptake and

564 effects of anthracene. *Aquatic Toxicol.*, 39: 307-320.

565 Moore, M.N., Readman, J.A., Readman, J.W., Lowe, D.M., Frickers, P.E. & Beesley, A. (2009).  
566 Lysosomal cytotoxicity of carbon nanoparticles in cells of the molluscan immune system: an *in*  
567 *vitro* study. *Nanotoxicology*, 3: 40-45.

568 Moore, M.N., Viarengo, A., Donkin, P. & Hawkins, A.J.S. (2007). Autophagic and lysosomal reactions  
569 to stress in the hepatopancreas of blue mussels. *Aquatic Toxicol.*, 84: 80-91.

570 Moore, M.N., Wedderburn, R.J., Lowe, D.M., Depledge, M.H. (1996). Lysosomal reaction to  
571 xenobiotics in mussel haemocytes using BODIPY-FL-verapamil. *Mar. Environ. Res.*, 42: 99-105.

572 Mount, D.I., Anderson-Carnahan, D.M. (1988). Methods for aquatic toxicity identification evaluations.  
573 Phase 1. Toxicity characterisation procedures (EPA/600/3-88/034). US EPA, Duluth, MN, USA,  
574 1988.

575 Murphy, S. D. (1980) Assessment of the potential for toxic interactions among environment pollutants.  
576 In *The Principles and Methods in Modern Toxicology* (C. L. Galli, S. D. Murphy, and R. Paoletti,  
577 Eds.), pp 571-578. Wiley, New York.

578 Patel, V.M., Lambert, W.C., & Schwartz, R.A. (2016). Safety of topical medications for scabies and lice  
579 in pregnancy. *Indian J. Dermatol.*, 61: 583–587.

580 Piras, G., Nakade, K., Yuasa, S., Baba, M. (1997). Three-drug combination of MKC-442, lamivudine  
581 and zidovudine *in vitro*: potential approach towards effective chemotherapy against HIV-1. *Aids*,  
582 11: 469-475.

583 Rashid, F., Horobin, R.W., Williams, M.A. (1991). Predicting the behaviour and selectivity of  
584 fluorescent probes for lysosomes and related structures by means of structure-activity models.  
585 *Histochem. J.*, 23: 450-459.

586 Readman, J.W. (1996). Antifouling herbicides - a threat to the marine environment? *Mar. Poll. Bull.*,  
587 32: 320-321.

588 Readman, J.W., Albanis, T.A., Barcelo, D., Galassi, S., Tronczynski, J., Gabrielides, G.P. (1993a).  
589 Herbicide contamination of Mediterranean estuarine waters: results from a MEDPOL Pilot  
590 Survey. *Mar. Poll. Bull.*, 26: 613-619.

591 Readman, J.W., Fileman, T.W. (2000). Inputs and Fluxes of Estuarine Micropollutants to the North  
592 Sea. Report to the Department of the Environment, Transport and the Regions, Contract - CWO  
593 637, 56p.

594 Readman, J.W., Fowler, S.W., Villeneuve, J.-P., Cattini, C., Oregioni, B., Mee, L.D. (1992a). Oil and  
595 combustion-product contamination in the gulf marine environment following the war. *Nature*,  
596 358: 662-665.

597 Readman J.W., Liang Wee Kwong, L., Mee, L.D., Bartocci, J., Nilve, G., Rodriguez-Solano, J.A.,  
598 Gonzalez-Farias, F. (1992b). Persistent organophosphorus pesticides in tropical marine  
599 environments. *Mar. Poll. Bull.*, 24: 398-402.

600 Readman, J.W., Liang Wee Kwong, L., Grondin, D., Bartocci, J., Villeneuve J.-P., Mee, L.D. (1993b).  
601 Coastal water contamination from a triazine herbicide used in antifouling paints. *Environ.*  
602 *Science Technol.*, 27: 1940-1942.

603 Readman, J.W., Mantoura, R.F.C., Preston, M.R., Reeves, A.D., Llewellyn, C.A. (1986). The use of  
604 pollutant and biogenic markers as source discriminants of organic inputs to estuarine sediments.  
605 *Internat. J. Environ. Anal. Chem.*, 27: 29-54.

606 Readman, J.W., Bartocci, J., Tolosa, I., Oregioni, B., Fowler, S.W. (1996). Recovery of the coastal  
607 marine environment in the Gulf following the 1991 war-related oil spills. *Mar. Poll. Bull.*, 32: 493-  
608 498.

609 Regoli, F. (1992). Lysosomal responses as a sensitive stress index in biomonitoring heavy metal  
610 pollution. *Mar. Ecol. Prog. Ser.* 84: 63-69.

611 Ringwood, A.H., Conners, D.E., Hogue, J. (1998). Effects of natural and anthropogenic stressors on  
612 lysosomal destabilization in oysters *Crassostrea virginica*. *Mar. Ecol. Prog. Ser.* 166: 163-171.

613 Sahai, J. (1996). Risks and synergies from drug interactions. *Aids*, 10, (S1): S21-S25.

614 Sapozhnikova, Y., Wirth, E., Schiff, K., Fulton, M. (2013). Antifouling biocides in water and sediments  
615 from California marinas. *Mar. Pollut. Bull.*, 69:189-94.

616 Sarigiannis, D.A., Hansen, U. (2012). Considering the cumulative risk of mixtures of chemicals – a  
617 challenge for policy makers. *Environ. Health*, 11(Suppl 1): S18.

618 Sforzini, S., Moore, M.N., Boeri, M., Bencivenga, M., Viarengo, A. (2015). Effects of PAHs and dioxins  
619 on the earthworm *Eisenia andrei*: a multivariate approach for biomarker interpretation. *Environ.*  
620 *Pollut.*, 196: 60-71.

621 Sforzini, S., Moore, M.N., Mou, Z., Boeri, M., Banni, M., Viarengo, A. (2017). Mode of action of Cr(VI)  
622 in immunocytes of earthworms: Implications for animal health. *Ecotoxicol. Environ. Safety*, 138:  
623 298–308.



624 Sforzini, S., Moore, M.N., Oliveri, C., Volta, A., Jha, A., Banni, M., Viarengo, A. (2017). Probable role  
625 of mTOR in autophagic and lysosomal reactions to environmental stressors in molluscs. *Aquatic*  
626 *Toxicol.*, 195: 114-128.

627 Sies H. (1997). Oxidative stress: Oxidants and antioxidants. *Exp Physiol.*, 82: 291–5.

628 Smith, K.E.C., Schmidt, S.N., Dom, N., Blust, R., Holmstrup, M., Mayer, P. (2013). Baseline toxic  
629 mixtures of non-toxic chemicals: “solubility addition” increases exposure for solid hydrophobic  
630 chemicals. *Environ. Sci. Technol.*, 47: 2026–2033.

631 Tallarida, R.J. (2012). Revisiting the isobole and related quantitative methods for assessing drug  
632 synergism. *J. Pharm. Exper. Therapeut.*, 342: 2–8.

633 Tallarida, R.J. (2016). Drug combinations: tests and analysis with isoboles. *Curr. Protoc. Pharmacol.*,  
634 72: 9.19.1–9.19.19.

635 Tang, J.Y.M., Busetti, F., Charrois, J.W.A., Escher, B.I. (2014). Which chemicals drive biological  
636 effects in wastewater and recycled water? *Water Res.*, 60: 289-299.

637 Tolosa, I., Readman, J.W. (1996). Simultaneous analysis of the antifouling agents: tributyltin,  
638 triphenyltin and IRGAROL 1051 in marina water samples. *Anal. Chimica Acta*, 335: 267 -274.

639 Tolosa, I., Readman, J.W., Blaevoet, A., Ghilini, S., Bartocci, J., Horvat, M. (1996). Contamination of  
640 Mediterranean (Cote d’Azur) coastal waters by organotins and IRGAROL 1051 used in  
641 antifouling paints. *Marine Pollution Bulletin*, 32: 335-341.


642 Tolosa, I., Readman, J.W., Fowler, S.W., Villeneuve, J.-P., Bayona, J.M., Albaiges, J. (1997). PCBs  
643 in the Western Mediterranean. Temporal trends and mass balance assessment. *Deep Sea*  
644 *Research*, 44: 907-928.

645 Valenti, A.M., Niero, A., Monti, G., Marangolo, F., Marangolo, M. (1997). Paclitaxel and N-  
646 methylformamide: In vitro interactions in human colon cancer cell line. *Anticancer Res.*, 1997,  
647 17: 2491-2497.


648 Widdows, J., Bakke, T., Bayne, B.L., Donkin, P., Livingstone, D.R., Lowe, D.M., Moore, M.N., Evans,  
649 S.V., Moore, S.L. (1982). Responses of *Mytilus edulis* on exposure to the water-accomodated  
650 fraction on North Sea oil. *Mar. Biol.* 67: 15-31.

651 Yermolaieva, O., Brot, N., Weissbach, H., Heinemann, S.H., Hoshi, T. (2000). Reactive oxygen  
652 species and nitric oxide mediate plasticity of neuronal calcium signalling. *PNAS*, 97: 448-453.

653  
654



658 **Fig. 1.** Toxicity as a cellular supracritical reaction based on complexity theory (Kauffman,  
659 1993). Using logarithmic scales, the molecular diversity of pollutant chemical species  
660 is plotted against the total concentration of pollutants in a complex mixture. A variety  
661 of modes of action is implicit in the diversity (see Kauffman, 1993). Idealised  
662 subcritical-supracritical boundaries are shown for alternative models where the effects  
663 are either synergistic, additive or antagonistic.  
664



668

669

670

671


672

673

674


675

**Fig. 2.** The % lysosomal toxicity derived from the NRR time (as % of control) plotted against toxicant diversity (Mean  $\pm$  95% CL). Individual dose/concentration responses are shown in A-G with the dose responses of the 3 mixture groups shown in H. The 50% effect isoboles are shown as grey dashed lines. The dose response data for lysosomal toxicity was tested using the Kruskal-Wallis non-parametric test.




676  
677  
678  
679  
680  
681  
682  
683  
684  
685  
686  
687  
688  
689  
690  
691  
692  
693  
694  
695

**Fig. 3. A** - Results of the 50% effect isobole method for identification of toxic lysosomal interactions. 50% isoeffective and additivity concentrations were determined for the three mixtures as explained in the Results section. The sums of concentration additivity (shaded boxes) were calculated from the generic formula  $d_a/D_a + d_b/D_b + \text{etc.....}$  where  $d_a$  and  $d_b$  are the doses/concentrations of A and B in a mixture that produces a specified effect (50% toxicity) and  $D_a$  and  $D_b$  are the doses/concentrations of the single compounds, which on their own elicit the same effect as the mixture (Kortenkamp & Altenburger, 1998). Additivity is indicated by a sum concentration additivity of 1.0, synergy by a sum concentration additivity of  $<1.0$ , and antagonism by a sum concentration additivity of  $>1.0$ . Isoeffective concentrations determined from dose response graphs for individual compounds and mixtures based on the 50% lysosomal toxicity (100% – NRR value as % of control). **B** - The same data is also shown as a  $\text{Log}_{10}$  scale graphical plot for both axes, as these are used in Figure 6 with transposed axes. Significant differences were indicated by non-overlapping 84% CIs (i.e., employing 84% rather than 95% confidence intervals [Buzatto, et al., 2015], it then being true for large  $n$  [ $n = 20$ ], that non-overlapping intervals correspond to a significant difference in a 5% level test). \* - indicates significant difference ( $P \leq 0.05$ ).




697  
698  
699  
700  
701  
702  
703  
704  
705  
706  
707  
708  
709  
710

**Fig. 4.** Observed (Obs) NRR values for the three mixtures and controls at three concentrations, and predicted (Pred) estimates from separate components (linear additive model), plus 84% confidence intervals for both (non-overlapping intervals imply significant differences). An – anthracene, Ph – phenanthrene, LAS – dodecylbenzene sulphonate, I – irgarol 1051, Pe – *cis*-permethrin, Li – lindane, M – malathion. Significant differences were indicated by non-overlapping 84% CIs (i.e., employing 84% rather than 95% confidence intervals [Buzatto, et al., 2015], it then being true for large  $n$  [ $n = 20$ ], that non-overlapping intervals correspond to a significant difference in a 5% level test).



711  
712  
713  
714  
715  
716

**Fig. 5.** Lysosomal toxicity (%) calculated from the NRR time (as % of control) plotted against mixture diversity (Mean  $\pm$  95% CL). The possible phase transition observed with the 3-component mixture (Mix 3) is outlined by the shaded box.



717  
718

719 **Fig. 6.** Evidence for synergistic toxicity as a consequence of increasing diversity of harmful  
 720 chemical. Test matrix (circular symbols) for chemical molar concentrations versus  
 721 chemical diversity ( $\log_{10}$  scales) with the % lysosomal toxicity (shaded boxes) shown  
 722 for the corresponding treatment. The conjectured curve for the subcriticality  
 723 /supracriticality boundary (broad grey line; see Figure 1) employs the 50%  
 724 isoeffective concentrations (open grey diamond shapes) for the 3 mixtures  
 725 (transposed from Fig. 3B; Kortenkamp & Altenburger, 1998). The 50% additivity  
 726 concentrations are shown as a dashed line (transposed axes from Fig. 3B).

727  
728  
729