The Seasonal Distribution of Macroplankton as shown by Catches in the 2-metre Stramin Ring-trawl in Off-shore Waters off Plymouth.

By F. S. Russell,

Naturalist at the Plymouth Laboratory.

In past years during the study of the vertical distribution of macroplankton much information has been obtained on the abundance of the different animals in the catches at different times of the year. Results have already been published to show the seasonal distribution of pelagic young fish (4), and also on the abundance and composition of the Sagitta population (5, p. 565). In this latter study the change over from the predominance of Sagitta elegans to that of S. setosa during the years 1930 to 1932 is striking. During the examination of catches it has been noticed that each year has been characterised at some time by the abundance of a species which may occur only in small numbers, if at all, in other years. This has been especially noticeable in the composition of the medusa population. It is necessary to follow these changes through a number of years to see if the presence of certain plankton organisms cannot be used as an index of conditions that may lead to other fluctuations such as those of the fisheries.

During 1930 and part of 1931 weekly catches of the 2-metre stramin ring-trawl were counted in order to obtain precise information on the seasonal abundance of the different animals throughout a whole year. The collections were made by oblique hauls of half an hour duration at the station 2 miles east of the Eddystone (4). Counts were made as on previous occasions by picking out directly distinctive and unusual animals and counting every animal in a $_{10}^{1}$ th sample of the remainder (2, p. 776).

The full results are given in the attached table which gives instructive information on the seasonal variation in abundance of the different animals, and their relative importance in numbers one to another. It seems unnecessary to analyse the tables further since any worker can abstract from it the data he may desire. One or two points must, however, be borne in mind.

1. The animals are all of a size that is normally retained in good condition by the stramin net. For instance, the medusæ are generally well-grown individuals; the earliest stages, while perhaps not small enough to pass through the meshes, are generally too badly damaged to allow of

definite identification. A useful indication of the size of disregarded medusæ is given by the absence of *Rathkea octopunctata* in Table I, which is common in tow-nettings in the earlier months of the year. This applies also to all other animal groups; the earliest stages of decapod larvæ do not appear in numbers in the catches as many will pass through the meshes.

In considering the seasonal distribution of any animal it must therefore be remembered that its first appearance in the catches does not imply the actual beginning of its abundance in the plankton. A period of two or three weeks should be allowed for the growth of the earliest stages to a size sufficient to appear in the catches.

2. The table does not contain any data on the abundance of young fishes which has been dealt with in another report (4).

3. In examining the data the vertical distribution of the various animals must be considered. The figures for *Tomopteris helgolandica* show, for instance, a great increase in July and August. This must only be regarded as an increase in abundance in the plankton; at other times the Tomopteris may actually be present in the area, but in the unsampled layers very near the bottom (see 3, from which data can be obtained for most of the animals).

4. The names given in the table are those used in the Plymouth Marine Fauna (1931). Phialidium sp. will be mostly Phialidium hemisphæricum: Phialella cymbaloides will also occur though not so commonly, as also a few Mitrocomella brownei. It was not possible in the time to analyse all the Decapod larvæ into species: the species that comprise each group can be found in the Plymouth Marine Fauna, where also in many cases is given the time of year at which the larval stages are most prevalent. Pandalid larvæ must be taken to include all species contained under Pandalidæ, Hippolytidæ, and Processidæ. Mysid sp. in summer consist chiefly of young specimens.

5. Examination of catches over a number of years has shown that there are species which can normally be expected to appear regularly each year. It is felt that these results are a fairly reliable picture of the average ringtrawl plankton catches to be found throughout any year. The following species, however, appeared in exceptional numbers and cannot be regarded as characteristic, but only for the year in question at the times they appeared:—

Aglantha rosea, Liriope exigua, and Stephanomia bijuga: Clione limacina and adult Meganyctiphanes norvegica also appeared in unusual numbers at the beginning of the year, although the larvæ of the latter are usually present then.

The results are to form a basis with which future years can be compared. Marked differences have been noticed in other years and it is hoped to publish these results in a separate report when sufficient data have been gathered. It should then be possible to decide on certain species whose appearance in the plankton should be watched for in future years. It is especially desirable to see whether any correlation can be found between the appearance of one characteristic species and that of another. The desirability of such observations was already stressed many years ago when Allen (1) prepared a list of species for special observation. Now that regular collections of these larger plankton animals are obtainable throughout the year an opportunity is afforded to attempt this study over a period of a number of years.

REFERENCES.

- Allen, E. J. The Distribution of Marine Plankton. Journ. Mar. Biol. Assoc., N.S., Vol. IV, No. 4, pp. 408–409, 1897.
- Russell, F. S. The Vertical Distribution of Marine Macroplankton.
 An Observation on Diurnal Changes. *Ibid.*, Vol. XIII, No. 4, p. 769, 1925.
- The Vertical Distribution of Marine Macroplankton. V. The Distribution of Animals caught in the Ring-trawl in the Daytime in the Plymouth Area. *Ibid.*, Vol. XIV, No. 3, pp. 557–608, 1927.
- 4. The Seasonal Abundance and Distribution of the Pelagic Young of Teleostean Fishes caught in the Ring-trawl in Offshore Waters in the Plymouth Area. *Ibid.*, Vol. XVI, No. 3, pp. 707–722, 1930.
- 5. On the Biology of Sagitta. IV. Observations on the Natural History of Sagitta elegans Verrill and Sagitta setosa J. Müller in the Plymouth Area. *Ibid.*, Vol. XVIII, No. 2, pp. 559–574.

TABLE I.

Numbers of Animals in Half-hour Oblique Hauls with the 2-metre Stramin Ring-trawl taken 2 miles East of the Eddystone.

				Dipurena (Slabberia) halterata.	Steenstrupia nutans.	Bougainvillia sp.	Amphinema dinema.°	Leuckartiara octona.	Laodicea undulata.	Cosmetira pilosella.	Obelia sp.	Phialidium sp.	Saphenia gracilis.	Aequorea forskalia	Aglantha rosea.	Liriope exigua.	Muggiaea atlantica.	Stephanomia bijuga.	Chrysaora isoscenes.	Cyanea lamarcki.	Aurelia aurita.	Pleurobrachia pileus.	Beroe cucumis.	Peachia arvæ.	Arachnactis larvæ.
193																									
Feb.	4th			-	_		-	_	-	-	-	-	-	***	52	_	_		-	-	-	40	-		-
,,	12th			-	-	-	_	-	-	_	-	-	_	-	2		-	880* -	-	-	-	80	-		-
,,	19th			-	-		-	-	-	_	-	-	_	-	7	-	_		-	-	-	3	-	-	_
. ,,	26th			-	-	-	_	-	-	-	-	_	_	-	8	-	-	1 -	_	-	-	3	_	_	-
Mar.	5th			_	_		_	_	_	_		_	-	-	5	-	-		-	_	-	6	2		
,,	12th			-	-		-	-	_	-	-	_		-	2	-	-	800* -	-	-	1†	10	2	-	-
,,	19th				_	The same	-	1.000	-	-	-	_	_	-	-		-	210* -	-		-	8	-	-	-
,,	27th			-	-	-	_	775	-	-	-	-	-			-			-	-	-	6	-	-	_
April	2nd	20		-	-	-	-	-	-	-	-		_		_				-	-	-		-		-
,,	11th	20			1100	_	-	-	-	-	_	40	-	-	7	***	-	632* -		1		7	-		-
,,	16th			_	-		-	-	-		-	20	1000	-	20	-	-	190* -	- 3	_	-	1		_	
,,	24th			_		-		1	_	_	-	140	-	-	7	_	-	450* -	-	6		_	-	-	10
,,	29th			10	-	-	to the same of	3	10	330	_	140	40		10	_	-	370* 1	Ĺ	1		10		-	_
May	7th			1	_			8	24	10	_	386	_		_					2	-	193		3	4
,,	15th			20	_	320	_	40	_	140	1,680	760	100		40			2,460* -		1	_	520		_	41
,,	22nd	- 60	10	_	_	150	_	93	5	5	40	100	10	_	90	2		760* -		î	_	520	-	1	_
June	10th	- 100			-	60	10	430	_	_	80	50	11		850	-			-	-	_	100	-	_	5
"	19th				1	-	10	179	10	-	260	140	1	-	1,000	1	00000			3		68	-	20	1
77	26th			-	60	261	-	280	20	250	1,025	1,340	î	-	5,736	-	-	1,640* -	-	1		180	-	2	$\frac{1}{2}$

July	4th			-	_	161	-	129	-	180	800	920	-	-	1,932	des	_	-	-	-	-	80	20	4	2	
,,	9th			-	120		-	270	-	600	43,800	3,780	90	-	6,150	1	-	30	-	-	-	_	120	15	-	
	14th			-	_	-	1,110	180	_	180	24,540	2,970	90	-	11,040	-	-	-	-	_	-	-	30	94	_	
,,	23rd			-	_	_	330	180	-		1,080	5,790	150	1	28,440	-	-		-	2			21	127	_	
"	29th			_	_	_	990	60	_	870	810	3,750		-	23,040	-	-	****	-	3	-	-	39	35		
Aug.						_	-	150	300	_	330	1,620	/	1	40,020	_	-	4000		-	****	-	26	-		
	14th				122	-	1	44	120	60	160	4,020	40	_	33,220	446		-	_	-	1	-	137	-	-	
,,	21st			1000	6	_	_	9	500	-	_	3,240	28	_	8,360	17,442	20	641*	_	-		_	33	-	-	
,,	28th			-	_	1000		120	3,240	_	_	2,540	20	_	3,540	7,440	20	2*	_	-		_	32	-	-	
Sont					1		_	92	9,800	_	80	180	80		1,080	11,600	20	40*	-	-	_	1	8	17	_	
Sept.	11th			20	1		_	21	4,160		100	4,300	-	_	340	4,280	_	_	1	_	_	161	33	55	-	
,,	16th			20	_	-		53	2,310	10	30	100	_	-	51	2,800	30	_	_	_	-	220	6	20	-	
,,	24th			_	_			77	10	-	- 50	640		-	180	3,760	20	_	-	_		800	4	_	-	
Oct	1st	•		_	_	-		18	10	-		60			50	180	_	-	_	_	- 1	.370	3		_	
Oct.				-	_		-	3	_	_		340		_	20	660	_	_	_	1	_	500	4	-	_	
,,	7th	•		-	-	_	_	0			20	510		-	_	4,100	20		_	-	-	-	_	_	_	
,,	14th			_	_	-		_	_	_	10	330	-	_	-	1,680		_	-	-	-	-	-	_	_	
N	16th 6th			-		_		_		_	10	20		_	_	1,650	_	_	_	_	_	-		_	-	
Nov.				_	_	_	_	10	_	_	_	20			_	980	50	_	_		_	20	_	_	_	
,,	13th			_	_	_	-	10	_	_		10	359	_	10	1,390	10	_	_	_	-	_	_	_	_	
,,	20th			_			100		-	_		10	_	227	-	710	10	_	_	_	_	_	-	_	_	
Ď	26th			****	-		7/15			_	15	100			_	1,115	_	-	_	_	_	10	_	_	-	
Dec.	3rd				_	_	_	_	_	_	~	20	_		_	1,020	20	_	_			1		-	-	
,,	10th			_	_	_	_	-	_	_	-	20	_			690	50	_	_		_	_	_	_	_	
,,	17th			office.		_	_		-	_	_	_	_	1000	8 5 7	140	10	_		_	_	_		_		
,,	22nd		•	etics.	_	_	-	_	-	_			-			110	10									
109																										
193																310	130	-	_	_	-	-	_	_	_	
Jan.	1st	•	•	-	_	-	-		_		-	_			_	230	170	_	_	-	-	_		-	_	
,,	5th			_	-	7.50	95905	_	_	-		70		_	_	970	1,166	_	_	_	-	-	-	-	_	
2.2	15th			-	-	_	-		_	_		30		_		104	130	_	_	_	_		_	-	-	
,,	22nd			-		_		_	_		_	90			_	10	10	_	_	_	_	_	-	-	-	
77.1	26th			-	_	_	_	_	-	_	_	_	_			10	30			_	_	_	_		-	
Feb.	6th			-	-	-	-	_				30	-	_			-				1†	_	_	_	_	
,,	12th			-	-	-		-	_	-	- ST	10	_		-	_	30		1000		-1	_	220	_	-	
,,	20th			-	-	_	-	1	_	_	_	80	_		-	2	- 50				022	2200		_	_	
435	23rd			-		_	_	20	-	-	_	1,820		-	_		_			20		_			_	
‡Mar				_	-		_	2	- 20	_	70		_	-	_	_	180			16	0.77	_		_		
,,,	26th	•		-	_	. —	_	10	30		70	4,520	_	_	-	_	20	_	_	3	_	20	1000		_	
April				-	-	-	_	2	_	20	-	220	-	-	_	-	20	_	-	23	_	40			1,480	
,,	16th	*			-	-	_	-	-	660	_	12,120	_	-	_	_	-	-	-	40		30	-		1,100	
		ο,	ъ.	1.1						4 31	1		1			* Noo	tophoros				+ 1	Enhvre				

 $^{^{\}circ}$ Possibly a few $A.\ rugosa.$

[‡] Medusæ only counted.

^{*} Nectophores.

[†] Ephyra.

TABLE I—continued.

			Tomopteris helgolandica.	Terebellid larvæ.	Pœcilochætus serpens larvæ.	Sagitta elegans.	Sagitta setosa.	Calanus finmarchicus.	Euchæta hebes.	Metridia lucens.	Candacia armata.	Anomalocera patersoni.	Labidocera wollastoni.	Caligus sp.	Mysid sp.	Nyctiphanes couchii adult.	Euphausiid larvæ.	Meganyctiphanes norvegica.	Apherusa sp.	Hyperia galba.	Themisto gracilipes.	Idotea sp.	Cumacea.
1930															25.22	1,000,000							
Feb.	4th		140	2	-	11,360	_	1,200	-	-	20		-	-	100	4,218	-	150	-	-	112	-	-
,,	12th		770	-	-	16,270	130	540	-	_	_	-	-	1	_	-	380	-	-	-	-	-	-
,,	19th		251	-	-	10,411	389	1,280	-	_	60		-	1	-	20	-	-	-	-	1	-	_
	26th		3	-	-	452	110	580	-	-	-	-	-	_	-		-		-	-	_	-	-
Mar.	5th		165	-		1,517	93	450	-	-	-		-	_	_	4		_	_	-	1	-	-
,,	12th		33			455	5	260	_	-	-		-	-	-	_	_	_	-		_	-	-
,,	19th		-	_		8	_	-	-		-	-	_	1	-		-	-	-	-	-	-	-
,,,	27th		3	-		1,860	-	420	-	-	_	-		2	-	2	_	-	-	-	_	-	
April			-	-		24	1	10	-	aio.	-	38	****	Ť	_	-		-	-	-	1	-	-
,,	11th		3	_	_	240	-	440	-	60	-	-	_	Ţ	-	2	210	-			_	-	-
,,	16th		_	_	7.0	21	-	16,890	-	70	60	30	-	1	_	1	310	1	-	-	130	-	-
,,	24th		-	-	10	860	-	7,630	-	-	510	_	_	1	_	- 1	1,140 540	_	-	_	2	-	_
3.5	29th		_	200		39	1	960	-	$\frac{60}{193}$	80 772	-	-	-	_	2	2,702	_	-	_	2	9	_
May	7th		7	386 60	-	77	2	318,450	-	80	260	-	_	_	_	4	580	_	_	_	_	0	
,,	15th		10 10	60	_	243 101		24,720	-		50	20	_	1	77.70	_	450	_	7	_	-	_	
,,, T.,,,,	22nd		63	_	110		_	11,210 1,968	_	-	200	4	_	1	1	$\frac{-}{4}$	40	_					-
June			25	_	110	$\frac{1,324}{637}$	6	6,592	_	_	220	22	_	2	1	*	20	_		5			_
,,	19th 26th	•	95	_		760		3,560	_	40	320	13	_	2	2	_	40	-	20	0	20		_
July	4th	•	10	_	_	2,218	_	2,023	_	40	200	62	_	2	2		40		20		1		_
	9th		664	_		4,377	8	6,132	_		2,160	60		2	360	_			60	-	1	_	-
,,	14th		741	_		3,942	4	8,370	_	_	930	+	_	_	152	_	30	_	-	_	_	-	_
,,	23rd		759	_	30	2,142	10	2,910	-	30	1,860	10	_	2	60	8	30		750	_	-	**	_
"	29th		376		-	1,423	12	1,007	_	30	270	-		4	6	-	-	-	240	_	1		~4
"	20011		010			1,120	14	1,001		30	210				4						-		

Aug.	7th		1,602	-	_	7,800	36	2,730	-	_	450	-	_	5	94	102	_	-	30	-	1	-	-
,,	14th		222	1*	-	5,210	5	1,041	-		220	1	_	8	17	_	140		200	-	1		
,,	21st		42	1†	-	2,038	_	803	-	_	200	_	-	4	-	_	20	-	240	-	21	-	20‡
,,	28th		4	-	_	1,018	_	402	_	-	80	_	_	1	-	-	-	_	140		22	-	-
Sept.	3rd		179	-	-	1,222	47	363	_	-	140	1	-	_	20	-	_		360	-	1	_	-
,,	11th		-		_	4,398	760	612	-	-	560	3	-	7	_	1	20	_	100	-	25	-	-
,,	16th		26	1°	_	558	46	233	-	_	40	1	10	-	-	-	-	_	_	-	3		_
,,	24th		4	-		270	87	35	-	-	-	53		-	-	-	7	-	-	-	2	1	-
Oct.	1st		8	-	-	102	45	70	-	-	20	-	-	1	-		_		_	-	2	-	-
,,	7th		_	_	_	74	369	263	_	-	30	-	-	1	_	-	-	-	-	1	-	-	-
,,	14th		3	-	_	303	683	_	_		150	-	_	2	1	-		_	-	-	-	-	-
,,,	16th		1	_	-	229	258	118	-		10	10	-	_	_	-		-	-	-	-	-	-
Nov.				_	_	441	300	_	_	-	10	_	_	1	1	_	_	-	-	-		$\overline{}$	5
,,	13th			-	10	159	120	99	_	_	40	-		-	_	1	10	-	_	-	_	_	-
"	20th			1 ****	_	32	73	7	_	-	_	_	_	1	_	_	-	-	_	-	_	-	-
"	26th	0	1	_	_	61	84	14	_	_	20	_	_	_	_	_	_	_	_	-	_	_	-
Dec.	3rd	0	_	-	-	730	1,080	107	_		40	-	-	1	_	_		-	_	-	-	_	-
,,	10th		_	_	-	360	162	129	-	_	100	_	_	1	_	_	121	_		_	1	-	_
	17th		-	-	-	262	372	169	-	-	70	_	-	-	_	-	30	_		_	1	_	_
,,	22nd		_		-	72	78	98	-	-	90	-	_	_	_	_	10	-	_	_	_	_	-
,,	22110	•																					
193							12101201	100						-									
Jan.	1st		1	_	-	. 210	249	20	_	-	60	-	-	2	_	-	-	_		-	-	-	_
,,	5th		-			109	59	80	10		120	-	-	1	_	_	10	-	-	-	_		10 -10
,,,	15th		1	-	-	1,765	339	190	_		150	_	_	-	10	3	_	-		-	_	_	-
,,			_	_	-	172	270	56	-	-	20		-	1	-	-	8	-	-	-	_	_	_
,,	26th		-	-		117	101	19	10		20	-	-	-	= 2	-	7	_	_	-	_	_	-
Feb.	6th			-	-	51	2	_	-	_	20	-	_	-		-	_	-	_	-	_	-	_
,,	12th			_	-	83	17	40	-	_	10	-	_	1	_	_		-	-	_	-	-	-
,,	20th		-	-	_	816	239	446	-	-	220	-	_	1		_	_	_		-	_	-	
,,	23rd		_	-		474	134	343	_	-	420	-	-	1	_	_	20	-	_	-	-	-	-
Mar.	17th		7			7,502	221									36		Re	st of sa	mpl	e not	cour	nted
,,	26th		25	321††	-	491	333	110	-	-	50	_	100	3	-	-	-	_	-	-	-	-	_
April				60	-	935	1,048	320	-	-	220	-	-	-	-	-	-	-	_			-	
,,	16th		30	200	-	764	385	1,260	-	-	180		-	1	-	-		_	40	-	-	-	
77		9																					

^{*} Young Syllid. \dagger Glycera. \ddagger Gravel in catch, near bottom. \parallel 1 S. serratodendata. $^{\circ}$ 1 large polychæte, fair amount of gravel. ** 1 Autolytus. \dagger † 1 Autolytus.

TABLE I—continued.

1930. Feb. 4th	Toursdon on Jones	Leanuer sp. larvæ.	Alphæid larvæ,	Crangonid larvæ.	Pontophilus spinosus larvæ,	Homarus vulgaris larvæ.	Phyllosoma larvæ.	Galatheid larvæ.	Galatheid post-larvæ.	Upogebia sp. larvæ.	Upogebia sp. post-larvæ.	Callianassa subterranea larvæ.	Axius styrhinchus larvæ.	Pagurid larvæ.	Pagurid glaucothoë larvæ.	Porcellana sp. zoea.	Porcellana sp. post-larvæ.	Ebalia sp. zoea.	Ebalia sp. megalopa.	Crab zoea.	Crab megalopa.	Squilla sp. larvæ.	Larval gastropods.	Echinospira larvæ.	Limacina retroversa.	Clione limacina.	Cephalopod young.	Young starfish.	Oikopleura.	Tornaria larvæ.	Fish eggs.	Total.
,, 12th ,, 19th		10	-	-	-	_	-	40	_	_	_	10	_	20 40	_	_	_	-	_	30 1,150	_	-	=	-	_	$\frac{3}{17}$ $\frac{10}{10}$	-	-	_	_	$\frac{20}{160}$ $\frac{110}{110}$	17,787 19,280 13,792
Mar. 5th	: -	_	- 2	200	_	_	$\frac{1}{2}$	90		_		-	-	40	-	_	-	-	-	2,540 4,850	_	-	-	_	-	5	-		_	-	_	3,844
,, 12th		_	-	40	-	_	_	10	_	_	_	_	_	20	_	_	_	_	_	910	_	_	_	_	_	2	1	_	10	-	4,020	11,888
,, 19th ,, 27th		20	-	20	-	-	- 1	40		-	-	-	-	580	-	_	-	-	-	16,780	-	-	-	_	_	_	_	_	_	_	2,500 5,620	4,041 23,177
April 2nd	:]	_	_	50	_	_	2 1	160	-	-	-	-	-	270	-	-	-	-	-	8,270	-	-	-	-	-	5	-	-	_	-	2,750	13,799
" 11th		30	_	_	1	_	ĩ '	10		_	-	-	-	10	_	_	_	-	_	930 130	_	-	_	_	_	-		_	140	-	4,590	5,987
" 16th		110	-	60	90	- 1	0 8	370	_	_	-	_	_	480	-	40	_	-	_	3,650	110	_	_	_	_	-	_	_	90	_	740	2,445
,, 24th		30		390	30	-		300	-	-	_	-	-	260	-	50	-	-	-	12,810	650	-	-	_	70		_	_	_	_	3,850 800	26,945 26,419
,, 29th May 7th	: -		-	90	40	-		330 737	-	\rightarrow		-	-	110	-	190	-	-	-	1,420	61*	-	-		-	-	-	-	100	_	900	5,888
" 15th	: =			40	120		-3.5		_	-	-		100	440	3	386	\leftarrow	-	-	7,334	386	-	-	-	-	-	-	-	579	-	1,737	335,784
,, 22nd				80	30	- 3		890	_	10		1770	-	440 540	_	960 810	-	-		- 19010	280	-	+	-	140		-	-	20	-	860	53,302
June 10th				50	_	- 2		40		60				150	_	1,120	-	_	_	8,430 9,530	550	Print.	+	10	90	-	-	-	_	-	190	27,671
" 19th				00	8			80	100	180	-	_		80		220		20	-	4,580	400 520	-	80	20	280 280	-	-	-	10	***	350	19,291
,, 26th		200		20	21	- 4		20	-	440	-	-	-	860	_	880	_	940		8,440	140	_	80	40	4,320	_	100	-	_	440	300	16,696
July 4th	. 2	80		20	-	- 1		80		-	-	-	-	140	40	800	-	-	-	1,540	180	_	40	-	600	_	1	_	_	120	960 1,120	37,052 13,830
,, 9th ,, 14th	. 31 . 35	300 510		20	1		- 2,1		- ,	,170	-	90	-	1,560	60	300	-	420			900	. Stee	300	-	1,710	-	1	_		4,890	120	96,342
99nd	. 33			40 90		- 7	8 2,7				30	60	-	360		3,060	120	780	_	12,900	1,230	-	+	_	23,310	_	1	-		3,030	180	105,037
,, 29th		maa		90	_	_ ′				360 930	1	$\frac{60}{120}$		1,710		1,740	120	360		17,220	2,280	-	+	-	8,790	-	1	-	-	720	30	85,176
,,		.00						20 0		000	1	120	12	750	-	2,550	-	30	90	4,920	1,500	-	+	-	2,250	-	4	-	30	930	30	48,554

S	", 1 2 ept. ", 1 1 2 ept. ", 1 ept. ", 2 ept. ", 2 ept. ", 1 ept. ", 2	7th 4th 21st 28th 1th 6th 4th 1st 7th 4th 6th 3th 6th 3rd 0th 7th		1 1 1 2 -	330 600 160 100 320 40 30 	-	80	-		1 5 2 - - -	98 8 14 6 6 2	0 - 0 - 0 - 0 20	- 1,1660 - 80 - 40 - 30 - 50 - 50 - 190 	5) 5) 5) 5) 5) 7	300 860 1,380 280 160 20		360 380 160 60 10 50 20 10 60	1 21 - 2 41	810 2,840 440 940 120 140 - 300 60 170 210 440 10	100 - 120 60 - - - - - - 1	180 20	30 40 - 40 100 20 - - - - - - -	8,880 2,620 580 3,220 1,840 470 900 250 210 440 60 30 10 70 40 20	1,650 980 400 680 540 100 40 80 30 40 20 20 10 10	30 5 - 10 2 1 - -	+ 1 40 40 		660		4 111 3 45 - 90 1 51 - 167 2 2 - 3 62 10	20 20 100 80 10 240 - 190 40 - 20 40	20	120 60 	76,736 56,799 36,010 26,397 30,658 21,279 7,587 7,369 2,196 2,930 6,316 4,044 4,044 2,183 2,070 1,180 3,274 2,185
	1,			-	_	-	10		_	-					_	-	120	_	_	_	_				_	_	_			_	100	_	20	1,958
,	96	2nd		-	100	Pine	10	-	-	\rightarrow			-	-	_	_	20	-	_	_	-	-	40	_	_	-	-	_		_	70	_	70	708
1	931.																																	
_		lst		_	_	_					10) -					00						00	90							00		0.0	
		5th	•	-	_	_	1				1	, -				_	70	_	-	_	_	_	90 210	20 10	_	-	_	-		-	60	-	90	1,342
,	1.0	5th	•	_	10		_			-		0 100		993	- 5	_	80			_	_		80	10	-		-			_	40	-	100	1,220
,	96	2nd		-	10	Free	_				5) -		_		_	20	_	_	_			140	10	_	_	_	-			590	-	1,270	6,704
,	90	6th	•	-	-	_	_			_		_		_	_	_	50		_	-	_		80	-	_	_	_	-		_	110 70	_	1,300	2,431
É		6th		_	_	-	10		1000		90					_	20	_	_	-	_	_	370	_	-	_	_	-		-	50	-	640	1,134
,	16	2th	0	_	_	_	-		_	_	90		-			_	90	_	-		_	_	240	_	-	-	_	-		_	50	_	910	1,563
,	90	Oth		-		-	_	20		_	810					_	260	_		_	_	_	6,840	30	_		_			_	110	_	530	1,182
,	99	3rd		-	30	-	10	10		-	750				_	_	710		_	-	_	_	7,170	90	_		_	- 5				-	7,140	16,973
,	96	6th		-	-	-	20	10		_	90				_	_	40	_	_	-	_	_	4,840	10	-		_	-	-	_	_	_	7,090 780	17,354
		lst		-	80	-	20	-		_	80				_	_	600	_	_	_	_		14,480	20	_	_	-	_		140	_	_	80	11,949
,	10	6th		_	140		60	_		-	200		00		20		1,460	-	360	-	_		11,060	140	_	-	_	-		140	_	_	1,040	18,368 31,683
																	,						,										1,010	01,000

[†] Pilchard.

