



# MarLIN

## Marine Information Network

Information on the species and habitats around the coasts and sea of the British Isles

### Common Skate (*Dipturus batis*)

MarLIN – Marine Life Information Network  
Marine Evidence-based Sensitivity Assessment (MarESA) Review

Ken Neal & Paolo Pizzolla

2006-12-13

A report from:

The Marine Life Information Network, Marine Biological Association of the United Kingdom.

**Please note.** This MarESA report is a dated version of the online review. Please refer to the website for the most up-to-date version [<https://www.marlin.ac.uk/species/detail/1436>]. All terms and the MarESA methodology are outlined on the website (<https://www.marlin.ac.uk>)

This review can be cited as:

Neal, K.J. & Pizzolla, P.F 2006. *Dipturus batis* Common Skate. In Tyler-Walters H. and Hiscock K. (eds) *Marine Life Information Network: Biology and Sensitivity Key Information Reviews*, [on-line]. Plymouth: Marine Biological Association of the United Kingdom.  
DOI <https://dx.doi.org/10.17031/marlin.sp.1436.1>



The information (TEXT ONLY) provided by the Marine Life Information Network (MarLIN) is licensed under a Creative Commons Attribution-Non-Commercial-Share Alike 2.0 UK: England & Wales License. Note that images and other media featured on this page are each governed by their own terms and conditions and they may or may not be available for reuse. Permissions beyond the scope of this license are available [here](https://www.marlin.ac.uk). Based on a work at [www.marlin.ac.uk](https://www.marlin.ac.uk)

(page left blank)

*Dipturus batis*.

Photographer: Davy Holt

Copyright: Davy Holt

See online review for  
distribution map

Distribution data supplied by the Ocean  
Biogeographic Information System (OBIS). To  
interrogate UK data visit the NBN Atlas.

<b>Researched by</b>	Ken Neal & Paolo Pizzolla	<b>Refereed by</b>	This information is not refereed.
<b>Authority</b>	(Linnaeus, 1758)		
<b>Other common names</b>	-	<b>Synonyms</b>	<i>Raja batis</i> (Linnaeus, 1758)

## Summary

### 🔍 Description

A large ray with a long pointed snout. Males growing up to 2 m in length, while females may reach up to 3 m in length. The leading edge of the wings is slightly concave and the small dorsal fins near the tip of the tail almost touch. The young have large thorns near the eyes and one row of thorns along the back of the tail, while older specimens lack the thorns near the eyes but have two rows of along the tail. The upper (dorsal) side is brownish-green with lighter spots and the underside dark grey, sometimes with black stripes, spots or marbling. Immature skate under 40lb in weight are jet black underneath which fades to grey as they get larger (Davy Holt, pers. comm.).

### 📍 Recorded distribution in Britain and Ireland


Populations of *Dipturus batis* are found off the coasts of Isles of Scilly, western British Channel, west and north Ireland and west Scotland.

### 📍 Global distribution

Atlantic coasts from Madeira and northern Morocco northward to Iceland including the North Sea. Also in western parts of the Baltic and western and northern Mediterranean.

 **Habitat**

The skate lives on sandy and muddy bottoms. The adults live in depths of 10 to 600 m while younger specimens prefer shallower waters.

 **Depth range**  
down to 600 m **Identifying features**

- Up to 3 m in length.
- Long, pointed snout.
- Juveniles have large thorns near their eyes and one row of thorns along the back of tail.
- Adults lack the eye thorns but have two rows of thorns along the tail.
- Dorsal surface is brownish green with lighter spots, the underside is dark grey, sometimes with spots, stripes or marbling.

 **Additional information**

Following a report in recent taxonomic literature, this species now belongs to the genus *Dipturus*, although may still be referred to as *Raja* in some texts, (see McEachran & Dunn, 1988).

 **Listed by** **Further information sources**

Search on:



## Biology review

### ☰ Taxonomy

Phylum	Chordata	Sea squirts, fish, reptiles, birds and mammals
Class	Elasmobranchii	Sharks, rays and skates
Order	Rajiformes	
Family	Rajidae	
Genus	Dipturus	
Authority	(Linnaeus, 1758)	
Recent Synonyms	Raja batis (Linnaeus, 1758)	

### 🌿 Biology

Typical abundance	
Male size range	22 - 200cm
Male size at maturity	150cm
Female size range	180cm
Female size at maturity	
Growth form	Pisciform
Growth rate	0.9 - 14kg/year
Body flexibility	Not relevant
Mobility	
Characteristic feeding method	Predator
Diet/food source	
Typically feeds on	Bristle worms, sand eels, crabs and flatfish
Sociability	
Environmental position	Demersal
Dependency	No text entered.
Supports	Host the copepod <i>Acanthochondrites annulatus</i> which attaches to the gills of the skate.
Is the species harmful?	No

### 🏛️ Biology information

The growth rate listed above may seem quite rapid but if weight at maturity is taken into consideration, 54 kg for males (Muus & Dahlstrom, 1974) and 94 kg for females (Walker & Hislop 1998) it can be seen that it takes many years to reach maturity. The data for growth rate came from tag and release studies off the west coast of Scotland (Sutcliffe, 1994; Little, 1995, 1998) by comparing weight change of skate between captures. Skates seem to have a start-and-stop growth pattern, where they have rapid growth for a short period and then remain at that weight for some time with no growth until they have another episode of rapid growth (Sutcliffe, 1994).

### 🖼️ Habitat preferences

<b>Physiographic preferences</b>	Open coast, Offshore seabed
<b>Biological zone preferences</b>	Lower circalittoral, Lower infralittoral, Upper circalittoral, Upper infralittoral
<b>Substratum / habitat preferences</b>	Coarse clean sand, Fine clean sand, Mixed, Mud, Muddy gravel, Muddy sand, Sandy mud
<b>Tidal strength preferences</b>	Moderately Strong 1 to 3 knots (0.5-1.5 m/sec.), Strong 3 to 6 knots (1.5-3 m/sec.), Very Weak (negligible), Weak < 1 knot (<0.5 m/sec.)
<b>Wave exposure preferences</b>	Exposed, Moderately exposed, Sheltered, Very exposed, Very sheltered
<b>Salinity preferences</b>	Full (30-40 psu)
<b>Depth range</b>	down to 600 m
<b>Other preferences</b>	None known
<b>Migration Pattern</b>	Non-migratory / resident

### Habitat Information

*Dipturus batis* was found around all British and Irish coasts except the south east. However, it has disappeared from much of its former range due to fishing pressure (Brander, 1981, Walker & Hislop, 1998; Jennings *et al.*, 1999; Rogers & Ellis, 2000). Remnant populations occur in areas unsuitable for commercial fishing (Shark Trust pers. comm.).

## Life history

### Adult characteristics

<b>Reproductive type</b>	Gonochoristic (dioecious)
<b>Reproductive frequency</b>	See additional information
<b>Fecundity (number of eggs)</b>	11-100
<b>Generation time</b>	10-20 years
<b>Age at maturity</b>	11 years
<b>Season</b>	Insufficient information
<b>Life span</b>	20-100 years

### Larval characteristics

<b>Larval/propagule type</b>	-
<b>Larval/juvenile development</b>	Oviparous
<b>Duration of larval stage</b>	Not relevant
<b>Larval dispersal potential</b>	No information
<b>Larval settlement period</b>	Not relevant

## Life history information

Females breed every other year (Little, 1997) and produce up to 40 eggs (Walker & Hislop, 1998) which are laid in the spring and summer (Whitehead *et al.*, 1984). The egg case is large, 15-25 cm

long and 8-15 cm wide (Dipper, 2001). It is rectangular and similar to 'mermaids purses' that are often found on the strandline. Eggcases are laid on the seabed and have been reported as being 'loose' on the seabed and perhaps very vaguely 'wedged' in between rocks (Paul Kay, pers. comm.). The young hatch after 2-5 months (depending on temperature) (Muus & Dahlstrom, 1974) at about 22 cm in length (Brander, 1981).

*Dipturus batis* is vulnerable to overfishing because of its slow growth, late maturity and low fecundity (Brander, 1981; Jennings *et al*, 1999). Only about 40 eggs are laid every other year and each generation takes 11 years to reach maturity, therefore populations cannot recover quickly from large mortalities. It has been estimated that a mortality of greater than 38% per year will lead to continual decline in the population and recovery is unlikely to occur until mortality is relaxed (Walker & Hislop, 1998). Numbers of common skate caught in trawls began to decline in the 1920s and again in the 1950s after a recovery period during the second World War and disappeared from the North Sea between the mid 1950s and early 1980s (Walker & Hislop, 1998). However, it has been shown that *Dipturus batis* can survive being trawled if it is released after capture (Little, 1995). *Dipturus batis* was recorded as 'not uncommon in trawls' in the Marine fauna of the Isle of Man (Bruce *et al.*, 1963) and was regarded as a common species by Hureau & Monod (1979). However the common skate had become all but extinct by the late 1970s (Brander, 1981). Similarly, between 1901 and 1907, the common skate made up 4% of all elasmobranchs caught in trawls in southwest England but between 1989 and 1997 none were caught (Rogers & Ellis, 2000). A tag and recapture program has been implemented in northeastern Scotland. Of 147 recaptured individuals, only 5 had travelled more than 20 km (Little, 1998), which suggests that *Dipturus batis* is vulnerable to local extinction by fishing with little chance of re-population from adjacent areas.

## Sensitivity review

This MarLIN sensitivity assessment has been superseded by the MarESA approach to sensitivity assessment. MarLIN assessments used an approach that has now been modified to reflect the most recent conservation imperatives and terminology and are due to be updated by 2016/17.

### A Physical Pressures

	Intolerance	Recoverability	Sensitivity	Confidence
<b>Substratum Loss</b>	Low	High	Low	Very low
<p>As <i>Dipturus batis</i> is highly motile, it would move away from an area that lost a suitable substratum and return when the area was back to normal. A certain amount of stress maybe caused by loss of food items with the substratum and because of the need to find new foraging/spawning areas. Therefore an intolerance of low has been recorded. Recoverability is likely to be high, resulting in a sensitivity assessment of low. Substrate removal is likely to destroy egg cases, but the location of important breeding and nursery grounds is poorly understood.</p>				
<b>Smothering</b>	Low	High	Low	Very low
<p><i>Dipturus batis</i> would move away from an area that was being smothered but with some stress due to loss of food and energetic costs of migrating to new foraging areas. Therefore an intolerance of low has been recorded. Recoverability is likely to be high, resulting in a sensitivity assessment of low. Egg cases on the sea bed are likely to be more sensitive.</p>				
<b>Increase in suspended sediment</b>	Not relevant	Not relevant	Not relevant	Not relevant
<p>It is not known whether an increase in suspended sediment will have an effect on <i>Dipturus batis</i>. Not relevant has been recorded because the skate is mobile enough to avoid local adverse effects.</p>				
<b>Decrease in suspended sediment</b>	Not relevant	Not relevant	Not relevant	Not relevant
<p>It is not known whether an decrease in suspended sediment will have an effect on <i>Dipturus batis</i>. Not relevant has been recorded because the skate is mobile enough to avoid local adverse effects.</p>				
<b>Dessication</b>	Not relevant	Not relevant	Not relevant	Not relevant
<p><i>Dipturus batis</i> is a sublittoral species unlikely to be subject to exposure to air. Therefore desiccation is not relevant.</p>				
<b>Increase in emergence regime</b>	Not relevant	Not relevant	Not relevant	Not relevant
<p><i>Dipturus batis</i> is a sublittoral species unlikely to be subject to exposure to air. Therefore an increase in emergence is not relevant.</p>				
<b>Decrease in emergence regime</b>	Not relevant	Not relevant	Not relevant	Not relevant
<p><i>Dipturus batis</i> is a sublittoral species unlikely to be subject to exposure to air. Therefore a decrease in emergence is not relevant.</p>				
<b>Increase in water flow rate</b>	Low	High	Low	Low
<p><i>Dipturus batis</i> has been recorded from sites around the UK with varying hydrodynamic conditions (see adult distribution) and therefore is unlikely to be affected by changes in flow</p>				



rate (JNCC, 1999).

**Decrease in water flow rate**      **Low**      **High**      **Low**      **Not relevant**

*Dipturus batis* has been recorded from sites around the UK with varying hydrodynamic conditions (see adult distribution) and therefore is unlikely to be affected by changes in flow rate (JNCC, 1999).

**Increase in temperature**      **Intermediate**      **Moderate**      **Moderate**      **Very low**

Sudden changes in temperature are unlikely to affect adults because they can move away but developing young may be affected. A study on a related species, *Raja eglanteria*, found that the embryos of this species do not develop at temperatures over 24 °C (Whitehead *et al.*, 1984). As such intolerance is assessed as intermediate. Chronic changes in temperature would also have little effect as the adults experience large temperature changes when moving between deep and shallow water. In addition, the world distribution of *Dipturus batis* is from the coasts of north-western Africa to the North coast of Norway so it is unlikely to experience temperatures outside of its tolerance range in British and Irish waters. Therefore recoverability of adults is likely to be high, however due to a lack of information on the effects on developing young, recoverability is assessed as moderate, resulting in a moderate sensitivity rating.

**Decrease in temperature**      **Intermediate**      **Moderate**      **Moderate**

Sudden changes in temperature are unlikely to affect adults because they can move away but developing young may be affected. As such intolerance is assessed as intermediate. Chronic changes in temperature would have little effect as the adults experience large temperature changes when moving between deep and shallow water. In addition, the world distribution of *Dipturus batis* is from the coasts of north-western Africa to the North coast of Norway so it is unlikely to experience temperatures outside of its tolerance range in British and Irish waters. Therefore recoverability of adults is likely to be high, however due to a deficit of information on the effects on developing young, recoverability is assessed as moderate, resulting in a moderate sensitivity rating.

**Increase in turbidity**      **Tolerant**      **Not relevant**      **Not sensitive**      **Very low**

An increase in turbidity could potentially interfere with foraging by inhibiting visual location of prey. However, *Dipturus batis* is a bottom feeder that is probably adapted to murky, silty water and utilize smell and electromagnetic cues to locate prey. Therefore the species is considered tolerant, and not sensitive has been recorded.

**Decrease in turbidity**      **Tolerant**      **Not relevant**      **Not sensitive**

A decrease in turbidity may aid predators. However, since man is the main threat to *Dipturus batis*, a decrease in turbidity is unlikely to increase the predation rate on this species. A decrease in turbidity may however, influence foraging success, either because prey gain an earlier warning of the skates' approach or because the skates' visual acquisition of prey is improved. Therefore the species is considered tolerant, and not sensitive has been recorded.

**Increase in wave exposure**      **Tolerant**      **Not relevant**      **Not sensitive**      **Low**

*Dipturus batis* has been recorded at sites all over the UK which vary from very sheltered to very exposed (see adult distribution) and anyway can swim to deeper areas if wave action increases to the extent that oscillatory movements on the seabed become excessive. As a result *Dipturus batis* is unlikely to be affected by an increase or decrease in wave exposure, so it is considered tolerant, and not sensitive has been recorded.

**Decrease in wave exposure**      **Tolerant**      **Not relevant**      **Not sensitive**      **Low**

*Dipturus batis* has been recorded at sites all over the UK which vary from very sheltered to very exposed (see adult distribution). Therefore the species is unlikely to be affected by an increase or decrease in wave exposure, so is considered tolerant, and not sensitive has been recorded.

**Noise** Tolerant Not relevant Not sensitive Not relevant

Fish with swimbladders have been reported to be the most sensitive to noise (Vella *et al.*, 2001). *Dipturus batis* is an elasmobranch and therefore does not have a swimbladder so is among the fish that are less sensitive to noise. However, sudden loud noises of low frequency have been shown to elicit an avoidance response in most fish (Vella *et al.* 2001). Noise rarely has a physiological affect on fish so *Dipturus batis* has been deemed tolerant, and therefore not sensitive to noise.

**Visual Presence** Low Immediate Not sensitive Very low

Adult *Dipturus batis* can be found at depths from the shallow sublittoral down to 600 m so are unlikely to be disturbed by boats or divers, although divers might disturb young skate in shallow water. Recoverability is likely to be immediate, however, since the skate can swim away from the disturbance and return when it has gone. Therefore an intolerance of low has been recorded, and the species is deemed not sensitive.

**Abrasion & physical disturbance** Intermediate Moderate Moderate Very low

*Dipturus batis* has a high resilience when trawled or caught by rod and line and then released again (Little, 1995). Therefore adults are probably tolerant of abrasion and physical disturbance at the benchmark level. Because of the shape of rays, they cannot escape trawl nets once they have been captured. A newborn skate is about 22 cm long and almost as wide, therefore is unable to pass through the mesh of fishing nets. Because of their small size there is a greater chance a juvenile skate will be damaged in a net than an adult skate. This could lead to high mortality/stress in the juveniles and affect the processes maintaining the population. Therefore an intolerance of intermediate has been recorded. Recoverability is probably moderate (see information below) hence sensitivity is assessed as moderate.

**Displacement** Low High Low Very low

*Dipturus batis* has a high resilience when trawled or caught by rod and line and then released again (Little, 1995). In addition, this species is found on a variety of substrata all around the UK and therefore is probably quite tolerant of displacement. Recoverability is likely to be high, resulting in a low sensitivity recording.

## Chemical Pressures

**Intolerance**      **Recoverability**   **Sensitivity**      **Confidence**

**Synthetic compound contamination** Not relevant Not relevant

No information found.

**Heavy metal contamination** Not relevant Very low

General information on the tolerance of fish to metal contamination reveals that part per billion concentrations are not lethal but may reduce gill activity, growth and hatching success of eggs. Copper was reported to be the most toxic of metals and suppressed egg hatching at concentrations of 10 parts per billion in certain teleost fish (Bryan, 1984). The leathery egg case of rays may make them less susceptible to metal contamination. However, in the absence of evidence on the effects in *Dipturus batis* no assessment can be made.

<b>Hydrocarbon contamination</b>					Not relevant	Not relevant
No information found.						
<b>Radionuclide contamination</b>					Not relevant	Not relevant
No information found.						
<b>Changes in nutrient levels</b>					Not relevant	Not relevant
No information was found on the effect of nutrient enrichment or algal blooms was found.						
<b>Increase in salinity</b>		Low	High	Low		Very low
As with many of the other factors, the high motility of this species allows it to escape adverse changes in salinity. Therefore intolerance is recorded as low and recoverability is likely to be high, resulting in a sensitivity assessment of low.						
<b>Decrease in salinity</b>		Low	High	Low		
As with many of the other factors, the high motility of this species allows it to escape adverse changes in salinity. Therefore intolerance is recorded as low and recoverability is likely to be high, resulting in a sensitivity assessment of low.						
<b>Changes in oxygenation</b>					Not relevant	Not relevant
No information was found on the effects of hypoxia on <i>Dipturus batis</i> .						

## Biological Pressures

	Intolerance	Recoverability	Sensitivity	Confidence		
<b>Introduction of microbial pathogens/parasites</b>					Not relevant	Not relevant
No information found.						
<b>Introduction of non-native species</b>					Not relevant	Not relevant
No non-native species are known to compete with the common skate.						
<b>Extraction of this species</b>	High	Low	High	High		High
<p>The slow growth rate, late maturity and low fecundity make <i>Dipturus batis</i> vulnerable to overfishing and it has disappeared from much of its former range due to fishing pressure (Brander; 1981; Walker &amp; Hislop, 1998; Jennings <i>et al.</i> 1999; Rogers &amp; Ellis, 2000). Only about 40 eggs are laid every other year and each generation takes 11 years to reach maturity, therefore populations cannot recover quickly from large mortalities. It has been estimated that a mortality of greater than 38% per year will lead to continual decline in the population and recovery is unlikely to occur until mortality is relaxed (Walker &amp; Hislop, 1998). Numbers of common skate caught in trawls began to decline in the 1920s and again in the 1950s after a recovery period during the second world war and disappeared from the North Sea between the mid 1950s and early 1980s (Walker &amp; Hislop, 1998). <i>Dipturus batis</i> was recorded as 'not uncommon in trawls' in the Marine fauna of the Isle of Man (Bruce <i>et al.</i>, 1963) and was regarded as a common species by Hureau &amp; Monod (1979). However the common skate had become all but extinct by the late 1970s due to overfishing (Brander, 1981). Similarly, between 1901 and 1907, the common skate made up 4% of all elasmobranchs caught in trawls in southwest England but between 1989 and 1997 none were caught (Rogers &amp; Ellis, 2000). Therefore an intolerance of high and a recoverability of low has been recorded. The species is highly sensitive to this factor.</p>						

**Extraction of other species**

Low

Low

*Dipturus batis* feeds on flatfish and some of the larger individuals take cod, haddock and herring (Dipper, 2001), and fishing pressure on these species may affect the skates food supply. Therefore an intolerance of low has been recorded. Recovery is dependant on the recovery of prey stocks, for which insufficient information has been found to make an assessment. Hence a moderate sensitivity has been recorded.

**Additional information****Recoverability**

In general, the recoverability of *Dipturus batis* after a mortality event from any source is very slow. Skates live for at least twenty years, reach maturity at 11 years and the females produce a clutch of about 40 eggs every other year (see reproduction). This means that a female produces a minimum of about 160 eggs in its lifetime, a very low fecundity. Even if every juvenile born after a mortality event survived, it is evident that it would take many years for a population to recover to its original numbers.

## Importance review

### Policy/legislation

UK Biodiversity Action Plan Priority	<input checked="" type="checkbox"/>
Species of principal importance (England)	<input checked="" type="checkbox"/>
Species of principal importance (Wales)	<input checked="" type="checkbox"/>
Scottish Biodiversity List	<input checked="" type="checkbox"/>
OSPAR Annex V	<input checked="" type="checkbox"/>
IUCN Red List	Critically Endangered (CR)
Features of Conservation Importance (England & Wales)	<input checked="" type="checkbox"/>
Priority Marine Features (Scotland)	<input checked="" type="checkbox"/>

### Status

<b>National (GB) importance</b>	Not rare/scarce	<b>Global red list (IUCN) category</b>	Critically Endangered (CR)
---------------------------------	-----------------	--	----------------------------

### Non-native

<b>Native</b>	-		
<b>Origin</b>	-	<b>Date Arrived</b>	-

### Importance information

When it was common, skate used to be taken as bycatch in trawls and the 'wings' were sold for human consumption.

*Dipturus batis* is listed under the UK Biodiversity Action Plan (Anon, 1999vii) and on the OSPAR Annex V list of threatened and declining species and habitats. Although listed as 'endangered' in the IUCN Red list, it is considered to be 'critically endangered' in coastal waters (IUCN, 2003). In addition, *Dipturus batis* has been proposed for protection under Schedule 5 of the Wildlife & Countryside Act 1981.

## Bibliography

- Anonymous, 1999vii. Common skate (*Raja batis*). Species Action Plan. <http://www.ukbap.org.uk/asp/UKPlans.asp?UKListID=543>, 2004-01-20
- Brander, K., 1981. Disappearance of common skate *Raja batis* from Irish Sea. *Nature*, **290**, 48-49.
- Bruce, J.R., Colman, J.S. & Jones, N.S., 1963. *Marine fauna of the Isle of Man*. Liverpool: Liverpool University Press.
- Bryan, G.W., 1984. Pollution due to heavy metals and their compounds. In *Marine Ecology: A Comprehensive, Integrated Treatise on Life in the Oceans and Coastal Waters*, vol. 5. *Ocean Management*, part 3, (ed. O. Kinne), pp.1289-1431. New York: John Wiley & Sons.
- Dipper, F., 2001. *British sea fishes* (2nd edn). Teddington: Underwater World Publications Ltd.
- Dolgov, A.V., Drevetnyak, K.V. & Gusev, E.V., 2005. The Status of Skate Stocks in the Barents Sea *Journal of Northwest Atlantic Fisheries Science*, **35**, 249-260
- Hayward, P.J. & Ryland, J.S. (ed.) 1995b. *Handbook of the marine fauna of North-West Europe*. Oxford: Oxford University Press.
- Howson, C.M. & Picton, B.E., 1997. *The species directory of the marine fauna and flora of the British Isles and surrounding seas*. Belfast: Ulster Museum. [Ulster Museum publication, no. 276.]
- Hureau, J.C. & Monod, T., (ed.) 1973. *Check-list of the fishes of the north-eastern Atlantic and of the Mediterranean*. Paris: Unesco
- IUCN, 2006. *2006 IUCN Red List of Threatened Species*. [On-line] <http://www.redlist.org>, 2003-01-01
- Jennings, S., Greenstreet, S.P.R. & Reynolds, J.D., 1999. Structural change in an exploited fish community: a consequence of different fishing effects on species with contrasting life histories. *Journal of Animal Ecology*, **68**, 617-627.
- JNCC (Joint Nature Conservation Committee), 1999. *Marine Environment Resource Mapping And Information Database (MERMAID): Marine Nature Conservation Review Survey Database*. [on-line] <http://www.jncc.gov.uk/mermaid>
- Little, W., 1995. Common skate and tope: first results of Glasgow museum's tagging study. *Glasgow Naturalist*, **22**, 455-466.
- Little, W., 1997. Common skate in the Sound of Mull. *Glaucus*, **8**, 42-43.
- Little, W., 1998. Tope and skate tagging off west Scotland. *Glaucus*, **9**, 36-38.
- McEachran, J.D. & Dunn, K.A., 1988. Phylogenetic analysis of skates, a morphologically conservative clade of elasmobranchs (Chondrichthyes: Rajidae). *Copeia*, **2**, 271-290.
- Mitchell, S. & Gallagher, M., 2000. The parasitic copepod *Acanthochondrites annulatus* (Olsson, 1869) from the blue skate *Raja batis*. *Irish Naturalists' Journal*, **26**, 323-323.
- Muus, B.J. & Dahlstrom, P., 1974. *Collins guide to the sea fishes of Britain and North-Western Europe*. Wm Collins Sons & Co. Ltd: London.
- Rogers, S.I. & Ellis, J.R., 2000. Changes in demersal fish assemblages of British coastal waters during the 20th century. *ICES Journal of Marine Science*, **57**, 866-881.
- Sutcliffe, R., 1994. Twenty years of tagging common skate and tope off the west coast of Scotland. In *Tag and release schemes and shark and ray management plans. Proceedings of the second European Shark and Ray Workshop, Natural History Museum, London, 15-16 February 1994* (ed. R.C. Earll & S.L. Fowler), pp. 14-16., Peterborough, Joint Nature Conservation Committee
- Vella, G., Rushforth, I., Mason, E., Hough, A., England, R., Styles, P., Holt, T & Thorne, P., 2001. Assessment of the effects of noise and vibration from offshore windfarms on marine wildlife. *Department of Trade and Industry (DTI) contract report, ETSU W/13/00566/REP*. Liverpool: University of Liverpool., *Department of Trade and Industry (DTI) contract report, ETSU W/13/00566/REP*. Liverpool: University of Liverpool.
- Walker, P.A. & Hislop, J.R.G., 1998. Sensitive skates or resilient rays? Spatial and temporal shifts in ray species composition in the central north-western North Sea between 1930 and the present day. *ICES Journal of Marine Science*, **55**, 392-402.
- Whitehead, P.J.P., Bauchot, M.-L., Hureau, J.-C., Nielson, J. & Tortonese, E. 1986. *Fishes of the North-eastern Atlantic and the Mediterranean. Vol. I, II & III*. Paris: United Nations Educational, Scientific and Cultural Organisation (UNESCO).

## Datasets

- Centre for Environmental Data and Recording, 2018. Ulster Museum Marine Surveys of Northern Ireland Coastal Waters. Occurrence dataset <https://www.nmni.com/CEDaR/CEDaR-Centre-for-Environmental-Data-and-Recording.aspx> accessed via NBNAAtlas.org on 2018-09-25.
- Isle of Wight Local Records Centre, 2017. IOW Natural History & Archaeological Society Marine Records. Occurrence dataset: <https://doi.org/10.15468/7axhcv> accessed via GBIF.org on 2018-09-27.
- Kent & Medway Biological Records Centre, 2017. Fish: Records for Kent.. Occurrence dataset <https://doi.org/10.15468/kd1utk> accessed via GBIF.org on 2018-09-27.
- Manx Biological Recording Partnership, 2018. Isle of Man historical wildlife records 1990 to 1994. Occurrence dataset: <https://doi.org/10.15468/aru16v> accessed via GBIF.org on 2018-10-01.
- Merseyside BioBank., 2018. Merseyside BioBank (unverified). Occurrence dataset: <https://doi.org/10.15468/iou2ld> accessed via

GBIF.org on 2018-10-01.

NBN (National Biodiversity Network) Atlas. Available from: <https://www.nbnatlas.org>.

OBIS (Ocean Biogeographic Information System), 2019. Global map of species distribution using gridded data. Available from: Ocean Biogeographic Information System. [www.iobis.org](http://www.iobis.org). Accessed: 2019-03-12

Outer Hebrides Biological Recording, 2018. Vertebrates (except birds, INNS and restricted records), Outer Hebrides. Occurrence dataset: <https://doi.org/10.15468/dax3tf> accessed via GBIF.org on 2018-10-01.

Scottish Shark Tagging Programme, 2018. Capture Mark Recapture Data for Scottish Elasmobranchs: 2009-2018. Occurrence dataset: <https://doi.org/10.15468/zno14z> accessed via GBIF.org on 2018-10-02.