

1 Environmental regulation of individual body size contributes to geographic variation in clonal
2 life cycle expression

3

4 Will H. Ryan^{1,2}, Leoni Adams³, Guido Bonthond⁴, Nova Mieszkowska^{3,5}, Kathryn E. Pack^{3,6},
5 and Stacy A. Krueger-Hadfield^{1,3}

6 ¹University of Alabama at Birmingham, Birmingham, AL, USA

7 ²Department of Biological Science, Florida State University, Tallahassee, USA

8 ³The Marine Biological Association of the United Kingdom, Plymouth, United Kingdom

9 ⁴Benthic Ecology, GEOMAR Helmholtz Centre for Ocean Research Kiel, Germany

10 ⁵School of Environmental Sciences, University of Liverpool, Liverpool, United Kingdom

11 ⁶Ocean and Earth Science, National Oceanography Centre Southampton, University of
12 Southampton, Southampton, United Kingdom

13

14 Corresponding author: wryan@uab.edu (661) 904-8649 ORCID: 0000-0002-4294-5454

15

16 **Abstract**

17 Clonal behavior has been hypothesized to provide an escape from allometric metabolic
18 scaling that limits the maximum mass achieved by a single individual. Here we demonstrate the
19 capacity of a wide-spread, non-native sea anemone to buffer its colony biomass accumulation
20 rate across environments by modulating ramet body size through environmentally-dependent
21 growth, fission, and catabolism. In 2015, thermal reaction norms for growth and fission behavior
22 were constructed using clonal lines of the sea anemone *Diadumene lineata*. In 2018, variation in
23 growth patterns under a factorial cross of temperature level and oxygen availability were
24 examined to test the hypothesis that individual ramet size is regulated by oxygen limitation in
25 accordance with optimal size theory. Across a wide range of temperatures, colonies accumulated
26 a similar amount of biomass despite a radical shift from unitary to clonal growth, supporting
27 fission as a mechanism to buffer growth rates over a range of conditions. Individual body size
28 appears to be regulated by the environment with increased temperature and reduced oxygen
29 modifying fission and mass-specific growth patterns, leading to the production of smaller-bodied
30 ramets in warm conditions. However, whether anemones in common garden conditions reduce
31 individual body size through catabolism or fission depends on the region of origin and may relate
32 to differences in seasonal temperature patterns among coastlines, which influence the energetic
33 benefits of fission rate plasticity.

34 35 **Introduction**

36 Despite the relative simplicity of the cnidarian body plan, an astounding diversity of life
37 cycle and growth patterns are achieved through variation in the timing and extent of investment
38 in clonal growth (Fautin 2002; Geller et al. 2005). While many hypotheses have been put forth to
39 explain an adaptive advantage of clonality under specific ecological conditions (see Francis

40 1979, 1988; Jackson 1977; Sebens 1979, 1987; Hughes 1987,1989), traits that influence fission
41 rate (Ferretti and Geraudie 1998; Geller et al. 2005), and how selection acts on these traits
42 (Reitzel et al. 2011), remain poorly understood. Many studies have looked for interspecific
43 patterns, comparing the distribution of clonal and aclonal taxa with ecological factors to
44 illuminate putative selective pressures (e.g., Chia 1976; Sebens 1979; Jackson 1985; Jackson and
45 Coates 1986; Francis 1988). Where intraspecific comparisons have been made, there appears to
46 be ample variation among genotypes in many aspects of growth, physiology, and fission
47 behavior (e.g., Shick et al. 1979; Sebens 1980; Ayre 1985; McManus et al. 1997; Edmunds 2007;
48 Reitzel et al. 2013), suggesting that traits governing clonality are labile. Phylogenetic
49 comparisons in anemones suggest that clonality has evolved independently several times (Geller
50 and Walton 2001). Consequently, life cycle traits are expected to respond readily to selection in
51 this group of cnidarians. However, our knowledge of the mechanistic pathways underlying
52 fission behavior in cnidarians remains rudimentary despite more than a century of interest in the
53 lives of clonal marine animals (but see Mire and Venable 1999; Geller et al. 2005; Reitzel et al.
54 2011). To evaluate hypotheses about which demographic and environmental factors favor the
55 evolution of clonality, more information is needed about the nature of variation in clonal
56 behavior among genotypes and how traits are shaped by the environment (McManus et al. 1997).

57 In both unitary and clonal animals, body size is a fundamental property that governs
58 metabolic rate and reproductive potential, as well as ecological relationships (e.g., competition
59 and predation risk). Thus, it is expected to be a key target of selection. Theories of metabolic
60 scaling suggest that smaller-bodied individuals with large respiratory surface area to biomass
61 ratios are more energetically efficient under diffusion limiting conditions than large-bodied
62 individuals (e.g. Atkinson 1994; Koojiman 2010; Glazier 2014). Similar logic has been used to

63 explain the observation that for a diverse array of taxa, warmer conditions lead to smaller adult
64 body sizes, even for species where reproductive fitness is positively correlated with body size
65 (Kingsolver and Huey 2008; but see Audzijonyte et al. 2018). The risk of oxygen limitation is
66 particularly important in aquatic organisms where the strong negative correlation between
67 temperature and dissolved oxygen availability acts as a fundamental constraint on body shape,
68 size, and performance (Pörtner 2001; Forster et al. 2012; Horne et al. 2015).

69 Under conditions that limit individual body size (e.g. hypoxia), clonal animals can have an
70 energetic advantage. Genets (the collective term for a group of descendants produced through
71 an asexual process) that are capable of dividing biomass into smaller units (known as ramets) can
72 grow indefinitely (i.e., iso- versus allo-metric growth), where unitary animals typically reach a
73 maximum size and cease somatic growth (Cancino and Hughes 1985; Sebens 1987).

74 Accumulating evidence suggests that the link between ramet body size and metabolic
75 performance may be critical for explaining the evolution of clonal and colonial life histories
76 (Burgess et al. 2018). However, the degree to which clonal animals can regulate growth and
77 fission rates to achieve energetically optimal ramet sizes is not well understood.

78 The role of fission in body size regulation is difficult to characterize, as metabolic, growth, and
79 fission rates often vary with the environment (e.g., Johnson and Shick 1977; Buss and
80 Blackstone 1991; Geller et al. 2005; Reitzel et al. 2013). Such effects may be an unavoidable
81 consequence of physical or chemical properties, leading to non-adaptive variation in the
82 expressed phenotype across environments (Gotthard and Nylin 1995). Alternatively, growth and
83 fission rate plasticity may be adaptive, allowing genetically identical clones to express a locally
84 optimal body size, and investment in asexual reproduction, across a range of environmental
85 conditions (Edmunds 2007). Indeed, for species that live in fluctuating environments, body size

86 plasticity through shrinkage (e.g., Levitan 1988; Chomsky et al. 2004) or a variable fission rate
87 (Ryan 2018) may be essential for tracking changing size optima through time.

88 To better understand the role of temperature and oxygen availability in regulating individual
89 body size and shaping clonal life histories, we examined growth, fission, and body size patterns
90 produced by the clonal sea anemone *Diadumene lineata* (Verrill) under a variety of experimental
91 conditions. We also explored variation in these patterns among individuals collected from
92 different parts of the extant range that differ in seasonal temperature patterns to look for evidence
93 of variation in asexual behavior that can inform predictions about the role of thermal regime on
94 the evolution of life cycle plasticity. Phenotypic plasticity can be described as a “reaction norm,”
95 or a linear relationship between an environmental gradient and a phenotype produced by a
96 particular genotype (Bradshaw 1965). Small differences in the curvature or slope of reaction
97 norms can lead to large differences in expressed phenotypic patterns, particularly in fluctuating
98 environments. We used this conceptual framework to understand how individuals may modulate
99 growth patterns to match energetically optimal patterns dictated by the local environment and
100 discuss the evolutionary implications of variation in reaction norms that govern clonal behavior.

101 *Diadumene lineata* is a small-bodied, clonal sea anemone that occurs in the high intertidal
102 zone. The species likely originated in East Asia, but has been spread around the world through
103 anthropogenic activity for more than 100 years (Uchida 1932; Cohen and Carlton 1995). Like
104 many non-native species, *D. lineata* persists under a broad range of abiotic conditions and occurs
105 across a range of habitat types. The success of this species has been attributed, in part, to a
106 prolific schedule of binary fission (Uchida 1932). Fission rate is also known to increase with
107 temperature (Miyawaki 1952; Minasian 1979) which may allow individuals to modulate body
108 size and reproductive effort in response to novel or fluctuating conditions and likely structures

109 seasonal and geographic patterns of sexual and clonal reproduction across its distribution (Ryan
110 2018; Ryan and Miller 2019).

111 Seasonal and geographic temperature patterns vary across the species' North American range
112 (Table 1), which may impose differential selection on life cycle expression. Highly seasonal
113 temperature patterns across the Atlantic coast mirror conditions along the species' native
114 distribution in Japan. Nearshore waters of the Gulf of Mexico are similarly seasonal, but on
115 average much warmer than the Atlantic coast. Across the Pacific coast, there are smaller seasonal
116 fluctuations and cooler average sea surface temperatures. As residents of the high intertidal zone,
117 these anemones can also experience large daily temperature fluctuations through tidal cycles,
118 though the species is often found in tidal pools, under rocks, or embedded among co-occurring
119 organisms that can help buffer short-term abiotic fluctuations. We expected highly seasonal
120 environments to favor genotypes that rapidly increased fission rate and reduce ramet body size
121 under high temperatures as such a mechanism would allow genets to track predictable changes
122 in the optimal body size on the scale of weeks to months. Conversely, for anemones in
123 environments where daily temperature fluctuations were larger than seasonal fluctuations (i.e.,
124 Pacific US), we expected fission to be less responsive to temperature, as engaging in fission in
125 response to short-term peaks in temperature could cause ramets to be perpetually below the
126 optimal body size for the environment, thus be maladaptive.

127 In this study, a series of experiments examine both the short-term (4 week) and longer-term
128 (12 week) responses of *D. lineata* to environmental manipulation. The short-term manipulation
129 of temperature and oxygen helps illuminate mechanisms of environmental body size regulation,
130 whereas the longer-term manipulation demonstrates the variation in growth and fission patterns
131 that emerge from such regulation. We also explore geographic variation in the shape of reaction

132 norms of *D. lineata* using individuals collected from the Atlantic, Gulf, and Pacific coasts of the
133 US, which vary substantially in their seasonal temperature regimes. We discuss our findings in
134 the context of potential local adaptation, although remain conservative in these conclusions as
135 patterns of genetic differentiation within and among sites are currently unknown. Specifically,
136 we ask:

- 137 (1) What are the general shapes of the thermal reaction norms of fission rate, individual body
138 size, and clonal biomass accumulation in this species?
- 139 (2) How do temperature, oxygen, and coastline of origin contribute to ramet body size
140 regulation via changes in fission and growth rate?
- 141 (3) Does basal metabolic rate differ among coastlines of origin?
- 142 (4) Are reaction norm differences among anemones from different coastlines consistent with
143 the expected effects of seasonality?

144

145 **Methods**

146 Experiment 1: Characterizing reaction norms across five levels of temperature

147 *Producing clonal replicates*

148 Twenty individual anemones were collected from each three field sites across the US
149 Atlantic and Gulf coasts (Nahant, MA [Atlantic, 42° N]; St. Simon, GA [Atlantic, 31° N]; St.
150 Teresa, FL [Gulf, 30° N]) in January 2013. Within sites, individual anemones were collected
151 from points spaced one to two meters apart in an effort to represent the range of genetic diversity
152 available at each site. No clonal replicates were knowingly included in the initial collection,
153 however the individuals used to initiate lab cultures reflected the genetic structures of the field
154 populations, which were unknown. Single-strand conformation polymorphism data from a single

155 locus performed on individuals haphazardly collected from sites along the Atlantic and Pacific
156 coasts suggests that most sites host multiple genotypes, and that repeated genotypes are common
157 within, but not among sites (Ting and Geller, 2000). Given the geographic distance among sites,
158 we are confident that the sample of individuals included in the experiment contained multiple
159 genotypes, however, none of our conclusions require this assumption.

160 Collected individuals were used to establish isolated clonal lines under common garden
161 conditions that exposed them to a seasonally adjusted range of temperatures between 15-29° C,
162 mimicking field measured conditions from St. Teresa, FL (see Ryan 2018). Between March and
163 September 2015, clonal lines were kept at 20° C and fed *Artemia* nauplii (Brine Shrimp Direct,
164 Ogden, UT, USA) two times per week.

165

166 *Temperature treatments*

167 In September of 2015, individual ramets were randomly selected from each of twelve
168 genets (FL: 5, GA: 4, MA: 3) that had sufficiently large clonal populations. Individual anemones
169 from each putatively distinct genotype were measured for pedal disk area, then isolated in a 50ml
170 Falcon tube of artificial seawater (Instant Ocean; salinity 32 ppt) and randomly assigned to one
171 of five temperature chambers (including three refridgerators (see Ryan 2018), a bench top
172 incubator (PR205075G Thermo Scientific, Waltham, MA, USA), and a climate control chamber
173 (I-36VL Percival Scientific, Perry, IA, USA). Chambers differed in make and size (ranging from
174 an interior volume of 0.05 to 0.85 cubic meters), so they were left dark to standardize light
175 conditions and prevent algal fouling. This species is not known to harbor photosymbionts and
176 often occurs under rocks in total darkness, therefore does not require light for growth or
177 nutrition. All of the chambers maintained static cultures within 1°C of the target temperature.

178 The water in each tube was exchanged and anemones fed to repletion on 3 day-old *Artemia*
179 nauplii (Brine Shrimp Direct, Ogden, UT, USA) twice per week. Five temperature treatments
180 were used (6, 9, 14, 21.5, and 29°C) spanning the range of average monthly water temperatures
181 experienced by this species on the east coast of North America. Because the experiment required
182 five temperature levels, we were unable to use more than one environmental chamber per
183 temperature treatment. Thus, chamber and temperature treatment level are unfortunately
184 confounded. However, as anemones were isolated in sealed vials within chambers, we have no *a*
185 *priori* reason to suspect systematic bias due to chamber identity.

186 Each genet was initially represented by 1 to 4 ramets in each temperature condition,
187 depending on replicate availability. Eight of 12 genets had at least two replicate anemones
188 assigned to each treatment. Replicate ramets were limited for the other four genets such that
189 some temperature treatments only had one anemone assigned (see Table S1). Of these genets,
190 only two were retained in analysis (see results). The probability of survival of each genet in each
191 temperature treatment was calculated as the number of initial replicates still represented by at
192 least one ramet at the end of the experiment (week 12) divided by the initial number of genotypic
193 replicates in each treatment. Fission rate was quantified as the number of daughter clones
194 produced by each individual over the experimental period. Body size was measured by tracing
195 the outline of the attached pedal disk onto a sheet of acetate, scanning the drawings into a
196 computer and using Image J software (Rasband 1997) to calculate the area of the pedal disk
197 (mm²). Initial pedal disk area was used to estimate dry mass using the regression measured in
198 Ryan (2018). All individuals were then rinsed in freshwater to remove extraneous salt, separated
199 into pre-tared foil boats and dried at 70° C for 72 hours. Dried tissue (µg) was weighed using a
200 microbalance. Final tissue density was calculated as dry mass per pedal disk area (µg/mm²).

201 Change in colony weight was calculated as the difference between final and estimated initial
202 colony dry weight. Mean individual mass was calculated as the mean dry mass (μg) of all ramets
203 within a replicate.

204

205 *Analysis*

206 The individuals used were biased toward those genets that had enough ramets available.
207 Mortality during the experiment left both GA and MA with complete final data for only two and
208 one genets, respectively (see details in Table S1). Thus, no attempt was made to characterize
209 variation among sites of origin in this experiment. Genet survival, fission rate, and change in
210 colony dry weight were genet-level traits, so genet ID was used as a random factor in all models
211 (Table 3). Ramet body size and tissue density were calculated as ramet-level traits, so both genet
212 ID and replicate ID within genet were used as random factors to account for variation among
213 ramets derived from independent cultures of clonal replicates derived from the same genet. No
214 genotype by treatment interactions were considered.

215 The effect of temperature on the probability of genet survival was estimated with
216 generalized linear mixed model (GLMER) with a logit-linked binomial distribution. The effect
217 of temperature on the number of clonal descendants (fission rate) was analyzed with GLMER
218 using a negative binomial distribution. The effects of temperature on change in colony dry
219 weight, ramet dry weight, and tissue density were analyzed with GLMERs using log-linked
220 Gaussian error distributions. In all cases, temperature was initially fitted with the highest order
221 polynomial supported by the levels of temperature (4th degree) to elucidate the shape of the
222 relationship. Model selection using AICc was then used to find the best-fit model using the
223 dredge function in the R (R Core Team, 2018) package MuMIn (Barton 2018). The model with

224 the lowest AICc value was chosen, except where a model with fewer parameters had a similar
225 AICc value ($dAICc < 2$) (see model details in Supplement 1). The significance of the contribution
226 of each retained parameter in the best fit model was evaluated using the type II Anova function
227 in the R package car (Fox and Weisberg 2011).

228 All analyses were done in R ver. 3.5.1 (R Core Team, 2018)

229

230 Experiment 2A: Characterizing growth and fission variation among individuals collected from
231 the Pacific, Gulf, and Atlantic coasts of the United States

232

233 *Anemone collection*

234 Between June and November 2017, *Diadumene lineata* individuals were collected from
235 ten sites in the species' US range (Table 2). At each site, from one to twenty individuals were
236 collected from each of five 0.25m² quadrats along a 25 meter transect through the high intertidal
237 zone, parallel to the water line, except at ESL where quadrats were placed at 5 meter intervals
238 along a floating dock. This resulted in the collection of between 19 and 100 individuals per site.

239 The number collected from each quadrat at each site varied, as this species has a patchy
240 distribution. For the experiments, samples were drawn randomly from anemones available from
241 each site without respect to quadrat of origin. To reduce the effects of prior environmental
242 variation, collected individuals were maintained separately in the laboratory in 50 ml tubes of
243 artificial seawater at 30ppt, 15° C temperature, on a 12:12 hour light:dark cycle from the time of
244 collection until use in the experiment, a period ranging from one to six months. Anemones were
245 fed weekly with Microvert liquid invertebrate food (Kent Marine, Franklin, WI, USA) until the
246 experiment began. Size variation naturally occurred among populations and persisted in collected

247 samples until the start of the experiment (Table 2). However, within site of origin, the mean
248 initial size of individuals assigned to each treatment group did not differ significantly when
249 compared with ANOVA (see results).

250

251 *Experimental design*

252 In December 2017, 20 - 25 individuals were randomly selected from among the common
253 garden-maintained cultures from each site and were assigned to one of four treatment conditions
254 representing a factorial cross of temperature (15° and 25° C) by dissolved oxygen level (50% and
255 100% of normoxia). As above, no clonal replicates were knowingly included in the experiment,
256 however the underlying genetic structure of these populations was unknown. Individuals used to
257 initiate lab cultures reflected a random sample of the genetic structures of the field populations.
258 For the purposes of this experiment, each individual was treated as an independent replicate,
259 however we were careful to avoid drawing conclusions with regard to the role of genetic
260 diversity underlying the observed variation in phenotypes.

261 Each individual was wet weighed, photographed for pedal area measurements (see
262 method in Experiment 1), then placed individually into a tube with 15 ml of artificial seawater
263 (30 ppt). Live anemones in tubes of seawater were shipped with ice packs from the University of
264 Alabama at Birmingham (UAB; Birmingham, AL, USA) to the Marine Biological Association of
265 the United Kingdom (MBA; Plymouth, UK) and arrived within 36 hours of shipment. Upon
266 arrival, 20 anemones from each site (except CFP, where only 19 individuals were available) were
267 transferred individually into the wells of twenty 12-well plates (Corning Inc., NY USA) (N = 5
268 per treatment) filled with filtered natural seawater (salinity 30 ppt), which were then divided

269 among four, 4 L sealable plastic tanks fitted with air stones. Remaining anemones were set aside
270 for experiment 2B (see below).

271 To facilitate water exchange with a surrounding tank, holes were pre-drilled into the lids
272 of the 12-well plates and then were lined with a fine mesh to prevent anemones from escaping or
273 moving among wells. The ability for water to exchange freely between the tank and each well
274 was confirmed by observing the ability of food dye to diffuse easily across the mesh when a
275 prototype plate was submerged in water. The plastic tanks, each containing five plates, were then
276 set in water baths to control their temperature. All anemones were maintained at approximately
277 10° C and aerated for two weeks to allow them to acclimate to the growth chambers with
278 minimal mass change and no fission. On January 2, 2018 a factorial cross of temperature and
279 oxygen level manipulations were initiated. Five individuals from each site, randomly positioned
280 across plates, were subjected to each of the four treatments. The temperature of the room was
281 raised to 15°C and then submersible aquarium heaters were added to water baths surrounding
282 half of the plates, raising their temperature to 25° C over the course of two days. Temperature
283 was monitored at five minute intervals with Hobo loggers (Onset, Bourne, MA, USA) in each
284 plastic tank and by daily checks with an infrared thermometer. To manipulate the availability of
285 dissolved oxygen, half of the tanks were aerated with ambient air fed from outside the building
286 (100% ambient oxygen), the remaining tanks were aerated with a pre-mixed gas of 10.5%
287 oxygen and 89.5% nitrogen (BOC, Plymouth, UK), equivalent to 50% of the ambient oxygen
288 treatments. Because pre-mixed gas was used, oxygen level was not monitored during the
289 experiment. While maintaining an independent environmental manipulation for each replicate
290 plate would have been ideal for statistical independence, it would have required a prohibitively

291 large volume of mixed gas to run the experiment. Thus, each treatment consisted of one
292 treatment tank that housed replicate plates.

293 Through the duration of the experiment, anemones were fed to repletion on a diet of two-
294 day old *Artemia* nauplii (Brine Shrimp Direct, Ogden, UT, USA) every other day. Plates were
295 removed from treatment tanks, the water in wells discarded (with care not to dislodge
296 individuals), and a 3 ml aliquot of a well-stirred culture of nauplii was pipetted into each well.
297 Plates were then returned to treatment tanks. This feeding protocol resulted in an 18% water
298 replacement in treatment tanks per week.

299 After four weeks, anemones were returned to 15 ml tubes of filtered seawater and
300 shipped back to UAB on ice where all anemones were wet weighed, photographed for pedal
301 area, and then dried at 72° C for 72 hours and dry weighed. Fission rate, change in colony mass,
302 change in mass-specific growth rate, and mean individual mass were calculated for each genet as
303 in Experiment 1.

304

305 *Analysis*

306 Variation in initial size among treatments and site of origin was assessed using a two-way
307 ANOVA. Since initial size varied among sites of origin, initial body size was considered in all
308 full models. Initial models also used site as a random variable to account for variation within
309 coastline. However, in no case did including site as a random variable improve the fit, thus it was
310 dropped for all analyses. In all cases, an initial model containing all predictor variables
311 (temperature, oxygen, and coastline of origin) and interactions as well as a polynomial series of
312 the natural log of initial wet weight, was constructed using GLMER in the lme4 package (Bates
313 et al. 2015) for R (R Core Team, 2018). To determine the shape of the relationship with initial

314 size, the initial model used the highest order polynomial supported by available degrees of
315 freedom (typically 5 degrees) (see supplement 2 for details). Stepwise model selection using
316 AICc was then used to determine the best fit models (Table 3) as described for experiment 1. The
317 probability of genet survival and probability of fission were modeled with binomial distributions.
318 The mass-specific change in mass, calculated as the natural log of the final wet weight minus the
319 natural log of initial wet weight, was normally distributed. The total change in colony mass,
320 calculated as the final wet weight minus the initial wet weight, was also modeled to provide a
321 visualization of biomass change patterns. But, given the highly leptokurtic distribution of this
322 metric, statistical inferences are best drawn from mass-specific analyses above. The final body
323 size of individual anemones (grams wet weight) was modeled with a log-linked gaussian
324 distribution. Genet ID was included as a random factor to account for the non-independence of
325 ramets produced by each genet. Analysis of Variance tables were constructed for each model to
326 aid in interpreting the contribution of each predictor. Goodness of fit values for all models were
327 calculated with the `r.squaredGLMM` function in the R package MuMIn or `rsquared` function in
328 the R package `piecewiseSEM` (Lefcheck 2018) for GLM(M)s.

329

330 Experiment 2B: Estimating relative differences in basal metabolic rate through starvation.

331 *Experimental design*

332 Thirty-three individuals representing seven sites of origin on three coastlines (Pacific: 4
333 sites (n = 5), Gulf: 2 sites (n = 4), Atlantic: 1 site (n = 5); Table 1) were used to measure loss of
334 body mass through starvation. Once at the MBA, these individuals were kept sequestered at 15°
335 C in individual tubes with 14 mL of artificial seawater, leaving a small headspace of air in each
336 tube. This temperature was chosen to minimize the likelihood of fission which would complicate

337 the interpretation of the results. Approximately once per week, tubes were agitated gently, but no
338 food was provided. After four weeks, anemones were returned to UAB on ice and were weighed
339 and measured as in Experiment 2A.

340

341 *Analysis*

342 Weight loss through starvation was calculated as the final wet weight (g) minus initial
343 wet weight. To test whether the rate of biomass catabolism depended on initial body size, weight
344 loss was regressed on the natural log of initial wet mass with log-linked gaussian linear
345 regression (GLM) (Table 3). Coastline of origin was treated as a fixed factor to evaluate
346 differences in starvation-induced shrinkage as a proxy for basal metabolic rate (Sebens 1981).

347

348 **Results**

349 *Experiment 1: Characterizing reaction norms across five levels of temperature*

350 Of the 125 individuals originally included in the experiment, 23 died soon after being
351 moved into the experiment. This transplant mortality was heavily concentrated among two
352 genets, which were both removed from all subsequent analyses (see Table S1). Among the
353 remaining ten genets (FL:5, GA:3, MA: 3) mortality was low during the experiment; the average
354 probability of replicate survival was 0.93. Genet identity was the major factor contributing to
355 variance in survival (conditional $r^2 = 0.23$). Temperature was not retained as a significant
356 predictor of mortality in the best-fit binomial GLMM (Figure 1A, see analysis details in S1). For
357 two genets, all replicates in 29° C died, precluding the construction of growth and fission
358 reaction norms. Thus, only the eight genets with complete data (FL: 5, GA: 2, MA: 2) were used
359 in subsequent analyses.

360 The number of clonal descendants produced ranged from one (no fission) to 13 ramets
361 and showed a significant, monotonic increase with temperature (GLMM, $\chi^2(1,73) = 67.75$, p
362 <0.001 ; Figure 1B). All surviving colonies accumulated biomass over the 12-week experiment.
363 Change in colony dry weight (mass accumulation) across temperature was best described by a
364 third order polynomial (GLMM, $\chi^2(3,71) = 66934$, $p <0.001$; Figure 1C). Mass accumulation
365 was lowest at low temperatures and highest at intermediate temperatures, peaking in the 14° C
366 treatment. Mass accumulation remained intermediate to high across the warmer temperatures
367 despite a rapid increase in fission; though, many genets showed a dip in mass accumulation at
368 21.5° C relative to 14 and 29° C. At the end of the experiment, individual ramet dry mass also
369 differed significantly across temperatures, which was best described by a third order polynomial
370 (GLMM, $\chi^2(3,168) = 51.89$, $p <0.001$; Figure 1D). Ramet body size was unimodal, peaking at
371 14° C. Individuals in the coldest treatment grew, but stayed small without dividing, whereas
372 individuals in warmer treatments ($\geq 21.5^\circ$ C) accumulated colony mass through the production of
373 daughter clones which were smaller-bodied than the founding individual. Tissue density showed
374 a significant, monotonic decline with increasing temperature (GLMM, $\chi^2(1,169) = 17.11$, p
375 <0.001 ; Figure S1). See Supplement 1 for model selection details and parameter estimates for all
376 analyses above.

377

378 *Experiment 2A: Characterizing growth and fission variation among genotypes collected from the*
379 *Pacific, Gulf, and Atlantic coasts of the United States*

380 Of the 199 individuals included in the initial experiment, data from 5 individuals were
381 exclude from analysis. Two individuals from CFP were too small for initial wet weights to be
382 measured confidently (< 0.0001 grams). Two individuals from WAS were an order of

383 magnitude smaller any others from the site (> 1.9 standard deviation units below the mean). One
384 individual from WAS was an order of magnitude larger than any others from the site (> 2.5
385 standard deviation units above the mean). These replicates were removed to prevent statistical
386 estimations from being extrapolation over a size range for which not all treatment levels were
387 represented. Statistical inferences were not altered by the inclusion or exclusion of these data.

388 There was variation in initial body size (the natural log of wet weight) among sites of
389 origin (Two-way ANOVA, $F(9,154) = 29.79$, $p < 0.001$), but no systematic variation among
390 assigned treatment levels (Two-way ANOVA, $F(3,154) = 0.862$, $p = 0.46$). There was also no
391 significant difference in initial size between treatment by site of origin (Two-way ANOVA,
392 $F(27, 154) = 0.52$, $p = 0.52$). The median initial wet weight was highest for Pacific coast
393 individuals, followed by Atlantic and Gulf Coast individuals (0.020, 0.018, 0.012 grams,
394 respectively) See Table 2 for the median initial wet weights by site for individuals used in
395 experiments 2A.

396 Over 4 weeks, survival was high (94%) among the 194 individuals included in the
397 experimental analysis. Exposure to low oxygen conditions significantly reduced individual
398 survival to 90% compared to 98% of individuals in high oxygen conditions (GLM, $r^2 = 0.18$, χ^2
399 $(1,193) = 6.00$, $p = 0.014$; Table S2). The lowest genet survival rate (83%) occurred for
400 anemones of Pacific origin experiencing both high temperature and low oxygen conditions,
401 though neither temperature nor coastline of origin were retained as significant predictors in the
402 best-fit GLM model. Likewise, initial body size was not retained in the final model. Site of
403 origin as a random factor was removed through model selection.

404 Among the individuals that survived, the probability of undergoing fission was
405 influenced by initial body size, coastline of origin, temperature, and oxygen treatments (GLM, r^2

406 = 0.49, Figure 2). As expected, high temperature significantly increased the probability of fission
407 (GLM, χ^2 (1, 173) = 47.77, $p < 0.001$). Coastline of origin had a significant effect (GLM, χ^2 (2,
408 173) = 17.95, $p < 0.001$); Gulf Coast individuals had the highest probability of dividing across all
409 treatments, followed by the Atlantic, then Pacific individuals. The same order was reflected in
410 the mean number of clonal descendants produced across treatments (Gulf: 1.40 +/- 0.08 se,
411 Atlantic: 1.32 +/- 0.10 se, Pacific: 1.08 +/- 0.03 se). The influence of initial body size differed
412 among coastline (GLM, initial size X coastline χ^2 (2, 173) = 6.74, $p = 0.034$). The probability of
413 fission declined with initial body size for Pacific genets and increased with body size for Gulf
414 genets. Initial body size showed no average effect for Atlantic genets, partly due to the
415 significant interactive effect between oxygen and initial size. In all cases, exposure to low
416 oxygen increased the probability of fission at large initial body sizes relative to the high oxygen
417 treatment (GLM, initial size X oxygen χ^2 (1, 173) = 4.89, $p = 0.027$) (see analysis details
418 Supplement 2).

419 Mass-specific weight change, or the growth rate per unit initial biomass, show a clear
420 monotonic decline with initial body size (GLM $r^2 = 0.60$, $F(1, 176) = 258.46$, $p < 0.001$; Figure
421 3A). High temperature reduced average growth (GLM, $F(1, 176) = 6.58$, $p = 0.011$) and caused a
422 marginally significantly steeper slope in the decline of growth with body size (GLM, initial size
423 x temperature $F(1, 176) = 3.78$, $p = 0.053$). Oxygen level did not change the average mass-
424 specific weight change (GLM, $F(1,176) = 0.115$, $p = 0.115$), but did have a significant
425 interaction with initial body size (GLM, initial size x oxygen $F(1, 176) = 5.06$, $p = 0.026$).
426 Individuals exposed to low oxygen showed a trend of dampened growth among small individuals
427 where growth rates were highest, but did not alter the threshold size above which anemones lost

428 mass. Coastline of origin was not retained in the best-fit model suggesting that all regions of
429 origin showed similar treatment responses (see analysis details in Supplement 2).

430 When the raw change in colony wet weight was plotted against initial wet weight (Figure
431 3B), both the energetic benefit of optimal size and the high cost of being too large are evident.
432 The best-fit model describing the change in colony wet weight is a fourth degree polynomial
433 (GLM, $r^2 = 0.67$), which demonstrates the peak in growth at an intermediate initial size, and
434 precipitous loss of mass for larger individuals (Model details are provided in supplement 2,
435 however, the influence of temperature and oxygen is best understood from the patterns of mass-
436 specific growth described above).

437 The final individual wet weight (ramet size) was influenced by initial size and
438 temperature (GLM, $r^2 = 0.56$, Figure 3C), and varied among genets within coastline (variance =
439 0.18, $sd = 0.43$). Final size was significantly, positively correlated with initial wet weight (GLM,
440 $\chi^2(1, 223) = 59.49$, $p < 0.001$); though, the slope of the relationship was consistently less than
441 one suggesting a tendency for body size to converge on a similar size within treatment over time
442 regardless of initial size. High temperature led to significantly smaller body sizes on average
443 (GLM, $\chi^2(1, 223) = 14.49$, $p < 0.001$) (median wet weight: 0.010 vs. 0.022 grams in low
444 temperature treatment). The rank order in body size among coastlines persisted (median wet
445 weight: 0.021, 0.015, 0.009 grams for Pacific, Atlantic, Gulf individuals, respectively), but
446 coastline of origin was not retained as an explanatory variable in the best-fit model (see
447 supplement 2). Likewise, there was little effect of oxygen treatment and this factor was not
448 retained in the final model. Interestingly, ramets from Gulf Coast genets tended to become
449 smaller through fission, whereas Pacific Coast ramets tended to shrink through catabolism
450 without undergoing fission (Figure 4). Atlantic Coast genets showed a mix of individual

451 shrinkage and fission to reduce body size. In most cases, fission produced two similarly sized
452 daughter clones (i.e., binary fission), though pronounced asymmetry in final ramet size was
453 observed particularly among Gulf individuals with large initial sizes (Figure 4). In one case, a
454 Pacific Coast individual produced a pedal lacerate.

455

456 *Experiment 2B: Estimating relative differences in basal metabolic rate among coastlines of*
457 *origin through starvation.*

458 All individuals survived the duration of the experiment and none underwent fission. All
459 individuals lost weight over four weeks, but the rate of weight loss, which is inversely
460 proportional to resting metabolic rate, was significantly influenced by both initial body size and
461 coastline of origin (GLM, $r^2 = 0.93$, Figure 5). The natural log of mass loss increased
462 significantly with the natural log of initial body size (GLM, $F(1, 22) = 288.29$, $p < 0.001$).
463 Individuals from the Gulf and Atlantic coasts lost significantly more mass than those from the
464 Pacific coast (GLM, $F(2, 22) = 5.99$, $p = 0.008$), but showed a similar slope (see Supplement 3
465 for statistical details). Because there was not complete overlap in initial body size distribution
466 available from the coasts, inferences about the performance of very large or very small
467 individuals are limited in this data set.

468

469 **Discussion**

470 Thermal growth curves for most ectothermic animals take the shape of a left skewed
471 distribution, with a gradual increase to a peak followed by a rapid decline in performance (Shulte
472 et al. 2011). The first experiment shows that the thermal reaction norms constructed for
473 *Diadumene lineata* over 12 weeks were different. They revealed the capacity of a clonal animal

474 to maintain high biomass accumulation rates across a wide range of temperatures while
475 modulating the size of ramets through changes in fission and growth. The second set of
476 experiments showed that individual body size is regulated by the abiotic environment via
477 changes in the slope of size-dependent growth curves. Over four weeks, individual growth,
478 catabolism, and fission all contributed to the pattern of individuals converging toward
479 environment-specific body sizes. Together these results support the hypothesis that life cycle
480 plasticity in clonal animals can stabilize growth across variable environments and encourages
481 more exploration into how reaction norms for clonal behavior are shaped by local environmental
482 patterns.

483 The interaction of temperature and dissolved oxygen with metabolic rate is complex. As
484 temperature increases, metabolic rate increases, which increases the demand for oxygen. At the
485 same time, higher temperatures reduce the capacity for water to hold dissolved oxygen. Thus,
486 both a direct reduction in oxygen input and temperature increase can increase the risk of oxygen
487 limitation. Metabolic demand for oxygen also increases with body size. Thus, the optimal body
488 size for avoiding oxygen limitation depends on both water temperature and rate that oxygen
489 flows into the environment.

490 In the longer-term experiment, fission rate increased linearly while individual ramet size
491 was unimodal over a broad range of temperatures. There was a transition from large and unitary
492 to small-bodied and clonal as temperature increased, though the exact slopes and inflection
493 points appeared to vary among genets. Notably, the rate of total colony mass accumulation was
494 similar for many genets between 14° to 29° C despite the major transition in growth pattern. Over
495 the 12 weeks of the experiment, fission served to stabilize tissue growth rates across a span of
496 temperatures over which oxygen consumption could reasonably be expected to triple ($Q_{10} \sim 2$;

497 Sassaman and Mangum 1970). The reduction observed for many genets in biomass accumulation
498 at 21.5° C may suggest that asymmetry in the energetic costs and benefits of fission may lead to
499 uneven growth across temperatures.

500 In the four-week experiment (2A), fission increased with temperature but showed a
501 complicated relationship with body size, oxygen level and coastline of origin. Fission appeared
502 to be stimulated when anemones were too large for the abiotic conditions, resulting in lost mass.
503 However, fission also occurred when anemones were growing rapidly, so were presumably well-
504 suited to the environment. Both patterns are consistent with oxygen limitation acting as a cue for
505 fission. These alternate roles for fission behavior may help account for the observed interaction
506 between oxygen treatment and body size in predicting the likelihood of fission in high
507 temperature conditions.

508 Overall, Gulf Coast anemones showed the highest propensity toward fission, consistent with
509 previous observations of high fission rates and small body sizes for Gulf versus Atlantic genets
510 of the species under a seasonal temperature cycle (Ryan 2018). The role of fission in growth also
511 varied among coastlines. For Gulf Coast genets, fission occurred most frequently where
512 individuals were initially large-bodied and played a role in reducing ramet size. Despite being
513 initially larger-bodied on average, Pacific Coast genets rarely engaged in fission during the four
514 week experiment. Pacific ramets that ended up smaller than the initial size mostly did so through
515 catabolism rather than fission, similarly to how unitary anemones perform under high
516 temperatures (Chomsky et al. 2004). Along with evidence of lower basal metabolic rates in
517 Pacific genets, these results are consistent with patterns of reduced temperature sensitivity found
518 for other organisms evolving under weak versus strong seasonal temperature fluctuation
519 (Baumann and Conover 2011).

520 Recent theory suggests that non-fluctuating environments favor thermal performance curves
521 that match optima based on mean temperature, whereas strongly seasonal environments tend to
522 favor higher resting metabolic rates and strategies that minimize the risk of stress during summer
523 high temperatures due to asymmetry in the energetic cost of being warmer rather than cooler
524 than optimal (Amarasekare and Johnson 2017). Because body size influences metabolic rate,
525 traits that modulate body size (such as growth and fission) are expected to be more responsive to
526 temperature in seasonal environments, where the energetic cost of environmental mismatch is
527 likely much higher (Scranton and Amarasekare 2017).

528 In anemones, an increased risk of exposure to heat-related hypoxia in predictably varying
529 environments may drive the evolution of fission rates that are more responsive to temperature.
530 Gulf anemones are at the highest risk of predictable periods of persistent hypoxia due to high
531 mean water temperatures coupled with strong seasonal fluctuations. In contrast, Pacific
532 anemones, which may encounter bouts of high temperature and hypoxia during low tide
533 (Helmuth et al. 2002), do not experience sustained and predictable exposure to warm, hypoxic
534 water to the same degree. As a consequence, Pacific anemones may use fission primarily to
535 maintain colony growth through the production of uniformly sized ramet whereas Gulf and
536 Atlantic populations use fission for both growth and as a means to rapidly modulate body size to
537 avoid hypoxic stress during seasonal flux. As a point of comparison, the larger-bodied clonal
538 species, *Anthoplura elegatissima* occurs across a similar latitudinal distribution on the Pacific
539 coast of the US. Fission in this species has been described primarily as a mechanism of asexual
540 growth driven by food availability, rather than as a mechanism for modulating body size through
541 temperature cycles (Sebens 1980,1982). However, we currently lack both the local-scale
542 temperature data and depth of sampling for *D. lineata* to evaluate the merits of this explanation.

543 Genets from all three coastlines showed size-dependent growth, where mass-specific
544 growth rate declined steeply with increased body size resulting in a unimodal pattern of total
545 colony mass change. As predicted, the energetic cost of being smaller than optimal (reduced
546 growth rate) is much lower than the cost of being too large (mass loss), perhaps favoring fission
547 behavior when the risk of hypoxia is high. Optimal size theory extends the predictions of
548 metabolic scaling theory to suggest that the body size that maximizes energetic efficiency
549 decreases monotonically from cold to warm conditions (Sebens 2002, Kingsolver and Huey 2008
550 Forster et al. 2011, Sheridan et al. 2011). Consistent with these predictions, a combination of
551 fission, growth, and shrinkage behavior led ramets to converge toward a environment-specific
552 body size. The size onto which ramets converged differed between temperature treatments and
553 mirrored the body size that led to the highest colony biomass accumulation, consistent with
554 regulation of growth and body size via allometric metabolic scaling (reviewed by Glazier 2014).
555 These results all support the existence of energetic advantage for genets that can modulate the
556 size of ramets produced in variable environments. Thus, the patterns we observed suggest a
557 degree of adaptation in the shape of the underlying reaction norms that govern fission and
558 growth behavior in *D. lineata*. However, there is little known about how such plasticity is
559 expressed under field conditions in spatially or temporally heterogenous environments. Here we
560 have interpreted patterns through the lens of oxygen limitation, but there are likely many
561 additional effects of temperature (e.g. changes in gene expression, tissue damage, etc.)
562 influencing growth and fission in ways that we have not yet investigated. Likewise, there is
563 much to explore about the role of invasion dynamics and habitat filtering in the geographic
564 patterns observed for this species, whose range has been expanded through human activity to
565 encompass three coastlines with very different climatic patterns. Differences in the invasion

566 history of non-native populations across the US could provide an alternate explanation for the
567 observed variation among coastlines. Future work elucidating patterns of genetic diversity and
568 connectivity within and among sites will likely provide critically needed context for interpreting
569 the eco-evolutionary significance of such variation.

570 These findings also contribute to a growing pool of observations in need of synthesis on
571 how environmental parameters influence asexual reproduction in sea anemones. For example,
572 several authors have concluded that asexual reproduction is a hallmark of growth in favorable
573 conditions, such as reporting increases in asexual behavior with increased food consumption
574 (*Diadumene lineata*: Minasian 1979; *Nematostella vectensis*: Reitzel et al. 2007), or in
575 temperatures associated with high survival and growth rates (*Metridium senile* (L.): Glon et al.
576 2019). While others have suggested fission to be a stress response, for example during starvation
577 (*Anthoplura elegantissima*: Sebens 1980,1982; *Exaptasia diaphana*: Bedgood et al. in revision).
578 There are a number of complications preventing a general explanation for asexual reproduction
579 in anemones, chief among them being the diversity of processes included in that description,
580 including binary fission, transverse fission, and pedal laceration (Fautin 2002). There is
581 intriguing variation even within *D. lineata*, where some populations in the native range are
582 known to asexually reproduce by pedal laceration rather than binary fission (Atoda 1973). In our
583 experiment, only one out of 194 individuals considered from across the US invaded range
584 displayed pedal laceration. The factors favoring this alternate asexual mode are unknown, and its
585 apparent rarity in invasive populations (it is previously undocumented outside of the native
586 range) remains unexplained. There is also some evidence that species differ in their capacity for
587 physiological acclimation (Zamer and Mangum 1979), though this has not been explored widely.
588 Another factor limiting a unified understanding, is the tendency to measure environmental

589 performance across a subset of conditions (e.g., two temperature treatments), using a narrow
590 range of potential body sizes. As is well appreciated in the plasticity literature, describing the
591 trend of a non-linear process depends heavily on which treatment levels are included (Murren et
592 al. 2014). The results of our study emphasize the complexity of the relationship between growth
593 and the abiotic environment, particularly for clonal organisms. It appears that fission can
594 increase in both favorable *and* stressful conditions. Reaction norm experiments are logistically
595 challenging, but necessary for understanding the true shape of environmental performance
596 curves. Likewise, concepts of metabolic scaling provide a strong framework for integrating
597 information about food and oxygen availability, temperature, body size, and shape, but require a
598 lot of data. Moreover, to understand the evolutionary basis of clonal life cycles, such
599 physiological knowledge needs to be combined with ecological and demographic data to capture
600 the multifaceted fitness effects of life cycle variation. Studies such as this one, however, suggest
601 that there are strong unifying mechanisms waiting to be characterized underneath the
602 overwhelming reproductive diversity in anemones.

603 In summary, the adaptive value of clonality depends on the degree to which the lifetime
604 fitness of a genet is increased by dividing its mass into multiple units as opposed to retaining all
605 biomass in a unitary body. For long-lived organisms in fluctuating environments, such as *D.*
606 *lineata*, the ability to manipulate body size through fission seems to offer an obvious advantage.
607 The fitness value of fission behavior must, however, be integrated across the environments a
608 genet experiences, which may be linked in different sequences, or on different time scales, all of
609 which influence the energetic costs and benefits of a given body size. Other size-dependent
610 phenomena combine with metabolic scaling dynamics to shape the adaptive landscape on which
611 organisms must “decide” if and when they should divide. Because sexual maturity and gamete

612 production depend on body size (Ryan and Miller 2019), fitness defined as the production of
613 sexual offspring might be very different between a single large or many small anemones. We
614 currently lack data to compare genet-level gamete or offspring production of replicates raised in
615 different temperature conditions, though this comparison is necessary to fully appreciate the
616 effects of temperature-dependent fission behavior on sexual fitness. Much future work is needed
617 to fully understand how and why species maintain clonal life cycles. But, with this study we add
618 to the collection of tantalizing observations that have long made this species a promising model
619 for the eco-evolutionary forces driving life cycle evolution.

620

621 **Compliance with ethical standards**

622 Data generated during the current study are available from the corresponding author on
623 reasonable request. The authors declare that they have no conflicts of interest. All applicable
624 international, national, and/or institutional guidelines for the care and use of animals were
625 followed.

626 **Acknowledgements**

627 We thank C.R. Hadfield and K. Atkins for logistics and help in the mesocosm at the Marine
628 Biological Association; M. Yant for help maintaining anemones at the University of Alabama at
629 Birmingham; J. Mutz and J. Imhoff for field assistance; J. Mutz and two anonymous reviewers
630 for comments on the manuscript; and B. Hughes and K. Wasson for access to Elkhorn Slough.
631 Funding was provided by the PADI Foundation (#21902) to WHR; the Ray Lankester
632 Investigatorship of the Marine Biological Association of the UK (2017-2019) to SAKH; Start-up
633 funds from the University of Alabama at Birmingham to SAKH; the Marine Biological

634 Association of the UK Fellowship to NM. WHR was supported as a UAB MERIT postdoctoral
635 fellow during the preparation of the manuscript.

636

637

638

639 **References**

- 640 Amarasekare P, Johnson C (2017) Evolution of Thermal Reaction Norms in Seasonally Varying
641 Environments. *Am Nat* 189:E31–E45
- 642 Atkinson D (1994) Temperature and Organism Size—A Biological Law for Ectotherms? *Adv*
643 *Ecol Res* 25:1–58
- 644 Atoda K (1973) Pedal laceration of the sea anemone, *Haliplanella Luciae*. *Publ Seto Mar Biol*
645 *Lab* 20:299-313
- 646 Audzijonyte A, Barneche DR, Baudron AR, Belmaker J, Clark TD, Marshall CT, Morrongiello
647 JR, van Rijn I (2018) Is oxygen limitation in warming waters a valid mechanism to explain
648 decreased body sizes in aquatic ectotherms? *Global Ecol Biogeogr* 28:64- 77
- 649 Ayre DJ (1985) Localized Adaptation of Clones of the Sea Anemone *Actinia tenebrosa*.
650 *Evolution* 39:1250–1260
- 651 Barton K (2018) MuMIn: multi-modal inference. R Package version 1.15.6
- 652 Bates D, Mächler M, Bolker B, Walker S (2015). Fitting linear mixed-effects models using lme4.
653 *Journal of Statistical Software* 67:1-48.
- 654 Baumann H, Conover DO (2011) Adaptation to climate change: contrasting patterns of thermal-
655 reaction-norm evolution in Pacific versus Atlantic silversides. *Proc R Soc B* 278:2265–2273
- 656 Bradshaw AD (1965) Evolutionary significance of phenotypic plasticity in plants. *Adv Genet*
657 13:115-155
- 658 Burgess SC, Ryan WH, Blackstone NW, Edmunds PJ, Hoogenboom MO, Levitan DR, Wulff JL
659 (2017) Metabolic scaling in modular animals. *Invertebr Biol* 136:456–472
- 660 Buss LW, Blackstone NW (1991) An Experimental Exploration of Waddington’s Epigenetic
661 Landscape. *Philos Trans R Soc B Biol Sci* 332:49–58

662 Cancino JM, Rodger N. Hughes (1985) An ecological overview of cloning in Metazoa. In:
663 Jeremy B. C. Jackson, Buss LW, Robert E. Cook (eds) Population Biology and Evolution of
664 Clonal Organisms. Yale University Press, New Haven, CT, p 153–186

665 Chia F (1976) Sea anemone reproduction: patterns and adaptive radiations. *Coelenterate Ecol*
666 *Behav*:261–270

667 Chomsky O, Kamenir Y, Hyams M, Dubinsky Z, Chadwick-Furman NE (2004) Effects of
668 temperature on growth rate and body size in the Mediterranean Sea anemone *Actinia*
669 *equina*. *J Exp Mar Bio Ecol* 313:63–73

670 Cohen AN, Carlton JT (1995) Nonindigenous aquatic species in a United States Estuary: A case
671 study of the biological invasions of the San Fransisco Bay and Delta. Washington D.C.: US
672 Fish and Wildlife Service.

673 Edmunds PJ (2007) Physiological ecology of the clonal corallimorpharian *Corynactis*
674 *californica*. *Mar Biol* 150:783–796

675 Fautin DG (2002) Reproduction of Cnidaria. *Can J Zool* 80:1735–1754

676 Ferretti P, Géraudie J (1998) Cellular and molecular basis of regeneration : from invertebrates to
677 humans. Wiley, Hoboken, NJ

678 Forster J, Hirst AG, Atkinson D (2012) Warming-induced reductions in body size are greater in
679 aquatic than terrestrial species. *Proc Natl Acad Sci* 109:19310–19314

680 Forster J, Hirst AG, Atkinson D (2011) How do organisms change size with changing
681 temperature? The importance of reproductive method and ontogenetic timing. *Funct Ecol*
682 25:1024–1031

683 Fox J, Weisberg S (2011) An R companion to applied regression, second edition. Sage, Thousand
684 Oaks, CA

685 Francis L (1979) Contrast Between Solitary and Clonal Lifestyles in the Sea Anemone. *Am Zool*
686 19:669–681

687 Francis L (1988) Cloning and aggression among sea anemones (Coelenterata: Actiniaria) of the
688 rocky shore. *Biol Bull* 174:241–253

689 Geller JB, Walton ED (2001) Breaking up and getting together: evolution of symbiosis and
690 cloning by fission in sea anemones (Genus *Anthopleura*). *Evolution* 55:1781–1794

691 Geller JB, Fitzgerald LJ, King CE (2005) Fission in sea anemones: Integrative studies of life
692 cycle evolution. *Integr Comp Biol* 45:615–622

693 Glazier DS (2014) Metabolic scaling in complex living systems. *Systems* 2:451–540

694 Glon H, Haruka Y, Daly M, Nakaoka M (2019) Temperature and salinity survival limits of the
695 fluffy sea anemone, *Metridium senile* (L.), in Japan. *Hydrobiologia* 830:303–315

696 Gotthard K, Nylin S (1995) Adaptive plasticity and plasticity as an adaptation: a selective review
697 of plasticity in animal morphology and life history. *Oikos* 74:3

698 Helmuth B, Harley CDG, Halpin PM, O’Donnell M, Hofmann GE, Blanchette CA (2002)
699 Climate change and latitudinal patterns of intertidal thermal stress. *Science* 298:1015–1017

700 Horne CR, Hirst AG, Atkinson D (2015) Temperature-size responses match latitudinal-size
701 clines in arthropods, revealing critical differences between aquatic and terrestrial species.
702 *Ecol Lett* 18:327–335

703 Hughes RN (1987) The functional ecology of clonal animals. *Funct Ecol* 1:63–69

704 Hughes RN (1989) *Functional Biology of Clonal Animals*, 1st edn. Springer Science and
705 Business Media, Suffolk

706 Jackson JBC (1977) Competition on Marine Hard Substrata: The Adaptive Significance of
707 Solitary and Colonial Strategies. *Am Nat* 111:743–767

708 Jackson JBC (1985) Distribution and ecology of clonal and aclonal benthic invertebrates. In:
709 Jackson JBC, Buss LW, Cook RE (eds) Population Biology and Evolution of Clonal
710 Organisms. Yale University Press, p 297–355

711 Jackson JBC, Coates AG (1986) Life Cycles and Evolution of Clonal (Modular) Animals. Philos
712 Trans R Soc B Biol Sci 313:7–22

713 Johnson LL, Shick JM (1977) Effects of fluctuating temperature and immersion on asexual
714 reproduction in the intertidal sea anemone *Haplanella luciae* (Verrill) in laboratory
715 culture. J Exp Mar Bio Ecol 28:141–149

716 Kingsolver JG, Huey RB (2008) Size, temperature, and fitness: Three rules. Evol Ecol Res
717 10:251–268

718 Kooijman SALM (2010) Dynamic Energy Budget Theory for Metabolic Organization.
719 Cambridge University Press, Cambridge

720 Lefcheck J (2018) Piecewise structural equation modeling. R Package version 2.0.2

721 Levitan DR (1988) Density-dependent size regulation and negative growth in the sea urchin
722 *Diadema antillarum* Philippi. Oecologia 76:627–629

723 McManus MG, Place AR, Zamer WE (1997) Physiological Variation Among Clonal Genotypes
724 in the Sea Anemone *Haliplanella lineata* : Growth and Biochemical Content. Biol Bull
725 192:426–443

726 Minasian LL (1979) The effect of exogenous factors on morphology and asexual reproduction in
727 laboratory cultures of the intertidal sea anemone, *Haliplanella luciae* (Verrill) (Anthozoa:
728 Actiniaria) from Delaware. J Exp Mar Bio Ecol 40:235–246

729 Mire P, Venable S (1999) Programmed Cell Death during Longitudinal Fission in a Sea
730 Anemone. 118:319–331

731 Miyawaki M (1952) Temperature as a factor influencing upon the fission of the orange-striped
732 sea-anemone, *Diadumene Luciae* (with 2 text-figures). J Fac Sci Hokkaido Univ, Ser 6 Zool
733 11:77–80

734 Murren CJ, Maclean HJ, Diamond SE, Steiner UK, Heskell MA, Handlesman CA, Ghalambor
735 CK, Auld JR, Callahan HS, Pfenning DW, Relyea RA, Schlichting CD, Kingsolver J (2014)
736 Evolutionary change in continuous reaction norms. Am Nat 183:453–467

737 Pörtner H (2001) Climate change and temperature-dependent biogeography: Oxygen limitation
738 of thermal tolerance in animals. Naturwissenschaften 88:137–146

739 R core team (2014) R: A language and environment for statistical computing.

740 Rasband WS (1997) Image J. National Institutes of Health, Bethesda, Maryland, USA.

741 Reitzel AM, Burton P, Krone C, Finnerty J (2007) Comparison of developmental trajectories in
742 the Starlet Sea Anemone *Nematostella vectensis*: embryogenesis, regeneration, and two
743 forms of asexual fission. Invertebr Biol 126:99–112

744 Reitzel AM, Chu T, Edquist S, Genovese C, Church C, Tarrant AM, Finnerty JR (2013)
745 Physiological and developmental responses to temperature by the sea anemone
746 *Nematostella vectensis*. Mar Ecol Prog Ser 484:115–130

747 Reitzel AM, Stefanik D, Finnerty JR (2011) Asexual reproduction in Cnidaria: Comparative
748 developmental processes and candidate mechanisms. In: Flatt T, Heyland A (eds)
749 Mechanisms of Life History Evolution: The Genetics and Physiology of Life. Oxford
750 University Press, Oxford, p 101–113

751 Ryan WH (2018) Temperature-dependent growth and fission rate plasticity drive seasonal and
752 geographic changes in body size in a clonal sea anemone. Am Nat 191:211–219

753 Ryan WH, Miller TE (2019) Reproductive strategy changes across latitude in a clonal sea
754 anemone. *Mar Ecol Prog Ser* 611:129-141

755 Sassaman C, Mangum CP (1970) Patterns of temperature adaptation in North American Atlantic
756 coastal actinians. *Mar Biol* 7:123–130

757 Schulte PM, Healy TM, Fangué NA (2011) Thermal performance curves, phenotypic plasticity,
758 and the time scales of temperature exposure. *Integr Comp Biol* 51:691-702

759 Scranton K, Amarasekare P (2017) Predicting phenological shifts in a changing climate. *Proc*
760 *Natl Acad Sci* 114:201711221

761 Sebens KP (1979) The energetics of asexual reproduction and colony formation in benthic
762 marine invertebrates. *Am Zool* 19:683–697

763 Sebens K (1980) The regulation of asexual reproduction and indeterminate body size in the sea
764 anemone *Anthopleura elegantissima* (Brandt). *Biol Bull* 158:370–382

765 Sebens K (1981) The allometry of feeding, energetics, and body size in three sea anemone
766 species. *Biol Bull* 161:152-171

767 Sebens KP (1982) Asexual reproduction in *Anthopleura elegantissima* (Anthozoa: Actinaria):
768 seasonality and spatial extent of clones. *Ecology* 63: 434-444

769 Sebens KP (1987) The ecology of indeterminate growth in animals. *Annu Rev Ecol Syst* 18:371–
770 407

771 Sebens KP (2002) Energetic constraints, size gradients, and size limits in benthic marine
772 invertebrates. *Integr Comp Biol* 42:853–861

773 Sheridan JA, Bickford D (2011) Shrinking body size as an ecological response to climate
774 change. *Nat Clim Chang* 1:401–406

775 Shick JM, Hoffmann RJ, Lamb AN (1979) Asexual Reproduction , Population Structure , and
776 Genotype-Environment Interactions in Sea Anemones. *Am Zool* 19:699–713
777 Ting J, Geller JB (2000) Clonal diversity in introduced populations of an Asian sea anemone in
778 North America. *Biol Invasions* 2:23-32
779 Uchida T (1932) Occurrence in Japan of *Diadumene luciae*, a remarkable actinian of rapid
780 dispersal (with plate IV, 1 chart and 4 text-figures). *J Fac Sci Hokkaido Univ, Ser 6 Zool*
781 2:69–82
782 Zamer WE, Mangum CP (1979) Irreversible Nongenetic Temperature Adaptation of Oxygen
783 Uptake in Clones of the Sea Anemone *Haliplanella luciae* (Verrill). *Biol Bull* 157:536–547
784

Table 1. Approximate seasonal and latitudinal patterns in nearshore sea surface temperature across the known range of *Diadumene lineata* in Japan and the United States. Mean sea surface temperature ranges approximated for 2014 – 2017 using data from NOAA National Data Buoy Center (<https://www.ndbc.noaa.gov/>). Species occurrence based on WoRMS data base (marinespecies.org) and personal observation.

Coast	Known latitudinal range of species	Seasonal range of monthly mean water temperature	Latitudinal range of mean annual water temperature
Pacific Japan (native)	Hokkaido to Oita	10 to 20° C	9 to 25 °C
Atlantic US	Cobscook Bay, ME to Cape Canaveral, FL	13 to 20° C	9 to 25 °C
Gulf of Mexico US	Corpus Christi, TX to Tampa Bay, FL	12 to 17° C	22 to 24 °
Pacific US	Vancouver, BC to San Diego, CA	2 to 5 ° C	9 to 17 ° C

Table 2. *Diadumene lineata* collection sites for geographical comparison. Individuals collected from sites in Washington were pooled to achieve sufficient replication from this region. No individuals from sites north of Virginia on the Atlantic coast could be acquired at the time of the experiment. Bold site ID denotes use in both experiments 2A and 2B. Median initial wet weight (g) was calculated for all replicates assigned to experiment 2A. SAKH: S.A. Krueger-Hadfield; GB: G. Bonthond; WHR: W.H. Ryan.

Coast	Site ID	Site	Collection date	Collector	Median initial wet weight (g)
Atlantic	ESL	VIMS Eastern Shore Laboratory, Wachapraque, VA	Sep. 2017	SAKH, GB	0.006
Atlantic	STS	St. Simon Island, GA	Nov. 2017	WHR	0.033
Gulf	WAK	Wakulla beach, FL	Nov. 2017	WHR	0.020
Gulf	ESP	Eastpoint, FL	Nov. 2017	WHR	0.021
Gulf	SGI	St. George Island, FL	Nov. 2017	WHR	0.009
Gulf	CFP	Copano fishing pier, Rockville TX	Jul. 2017	WHR	0.001
Pacific	ROK	Morro Bay, CA	Nov. 2017	WHR	0.038
Pacific	AZO	Azevedo Pond, Elkhorn Slough, CA	Sep. 2017	SAKH, GB	0.017
Pacific	BRK	Berkeley, CA	Nov. 2017	WHR	0.050
Pacific	WAS	Pooled from Grey's Harbor, WA & Willapa Bay, WA	Jun. 2017	WHR	0.011

Table 3. Overview of experiments and related analyses performed. Model structure reflects best-fit model resulting from model selection procedures. The predictor initial weight is the natural log transformed wet weight (g) of individual anemones at the start of the experiment.

Exp	Response variable	Model structure*	Error distribution	Link function
1	genet survival	$\sim 1 + (1 \text{genetID})$	binomial	logit
	number of clonal descendants	$\sim \text{temp} + (1 \text{genetID})$	negative binomial	log
	Δ colony dry mass	$\sim \text{temp} + \text{temp}^2 + \text{temp}^3 + (1 \text{genetID})$	Gaussian	log
	ramet dry weight	$\sim \text{temp} + \text{temp}^2 + \text{temp}^3 + (1 \text{genetID}/\text{rep})$	Gaussian	log
	tissue density	$\sim \text{temp} + (1 \text{genetID}/\text{rep})$	Gaussian	log
2A	genet survival	$\sim \text{oxygen}$	binomial	logit
	occurrence of fission	$\sim \text{initial weight} * (\text{coast} + \text{oxygen}) + \text{temp}$	binomial	logit
	mass-specific Δ colony wet weight	$\sim \text{initial weight} * (\text{temp} + \text{oxygen})$	Gaussian	identity
	Δ colony wet weight	$\sim \text{temp} * \text{initial weight} + \text{initial weight}^2 + \text{initial weight}^3 + \text{initial weight}^4$	Gaussian	identity
	Individual wet weight	$\sim \text{initial weight} + \text{temperature} + (1 \text{genetID})$	Gaussian	log
2B	Δ wet weight	$\sim \text{initial weight} + \text{coast}$	Gaussian	log

*model structures are displayed following the syntax of the R package lme4 (Bates et al. 2015)




Figure 1. (Experiment 1) Reaction norms for four variables across five levels of temperature for 10 genotypes of *Diadumene lineata* grown for 12 weeks in common garden conditions (experiment 1), including (A) Probability of genet survival, (B) the number of clonal descendants produced, (C) the natural log of the change in estimated colony dry mass, and (D) ramet body size (ln dry mass (ug)). Colored lines show average pattern for each genet, black lines indicate the best-fit linear regression model. Points show either genet replicate-level (A,B,C) or ramet-level (D) values. Different point shapes of the same color designate independent replicates of a genet within temperature level. Genets 7 and 10 excluded from statistical analysis as all replicates raised at 29° C died. Points horizontally jittered to reduce overplotting.




Figure 2. (Experiment 2A) The probability that *Diadumene lineata* individuals engaged in fission over 4 weeks in treatment conditions (experiment 2A) depending on initial body size and coastline of origin (panels). Lines reflect the best-fit linear model for individuals exposed to a temperature of either 15 °C (solid line, dots) or 25 °C (dashed line, stars), and an oxygen level of either 50% (grey) or 100% (black) of normal. Points are vertically jittered to reduce overplotting.




Figure 3. (Experiment 2A) The mass-specific change in wet weight (A), change in total colony wet weight (g) (B), and ramet body size (ln wet weight [g]) (C) as a function of initial body size (ln wet weight (g) of *Diadumene lineata* individuals after 4 weeks in treatment conditions (experiment 2A). Lines reflect the best-fit linear model for individuals exposed to a temperature of either 15 °C (solid line, dots) or 25 °C (dashed line, stars), and an oxygen level of either 50% (grey) or 100% (black) of normal. Model fit does not differ statistically between oxygen levels for (B) and (C). Panels in (C) show coastline of origin. The light grey line in each panel denotes a zero-change isocline.




Figure 4. (Experiment 2A) Final versus initial size (ln wet weight (g)) of *Diadumene lineata* ramets after 4 weeks depending on fission activity (colors), treatment temperature (15 vs 25o C shown by dots vs. stars), and coastline of origin (panels). Lines connect ramets produced by the same genet. Points below the zero-change isocline (grey line) show ramets that are smaller than the initial individual, either due to fission or shrinkage. One case of pedal laceration was observed (pl).




Figure 5. (Experiment 2B) The relationship between initial mass and mass lost (ln wet weight [g]) through starvation over four weeks at 15^o C (experiment 2B). Weight loss through starvation is inversely proportional to resting metabolic rate. Lines reflect best fit linear model. Point shade and shape reflect coastline of origin. Regions of the initial size spectrum that did not have representatives from all three coasts available (i.e. regions beyond the best fit lines) were excluded from statistical analysis.