

## Production of methanol, acetaldehyde, and acetone in the Atlantic Ocean

Joanna L. Dixon,<sup>1</sup> Rachael Beale,<sup>1</sup> and Philip D. Nightingale<sup>1</sup>

Received 18 July 2013; revised 23 August 2013; accepted 29 August 2013; published 12 September 2013.

[1] The biogeochemistry of oxygenated volatile organic compounds (OVOCs) like methanol, acetaldehyde, and acetone in marine waters is poorly understood. We report the first in situ gross production rates for methanol, acetaldehyde, and acetone of 49–103, 25–98, and 2–26 nmol L<sup>-1</sup> d<sup>-1</sup> over contrasting areas of marine productivity, including oligotrophic gyres and eutrophic upwellings. Photochemical production estimates are mostly negligible for methanol, up to 68% for acetaldehyde and up to 100% of gross production rates for acetone. Microbial surface OVOC oxidation to CO<sub>2</sub> accounts for between 10–50% and 0.5–13% of the methanol and acetone losses, respectively, but largely control acetaldehyde concentrations (49–100%). Biological lifetimes in a coastal upwelling vary between ≤1 day for acetaldehyde, to approximately 7 days for methanol and up to ~80 days for acetone. In open oceanic environments, the lifetime of acetaldehyde ranges between 2 and 5 h, compared to 10–26 days for methanol and 5–55 days for acetone. **Citation:** Dixon, J. L., R. Beale, and P. D. Nightingale (2013), Production of methanol, acetaldehyde, and acetone in the Atlantic Ocean, *Geophys. Res. Lett.*, **40**, 4700–4705, doi:10.1002/grl.50922.

### 1. Introduction

[2] Oxygenated volatile organic compounds (OVOCs) including methanol, acetaldehyde, and acetone are ubiquitous in the atmosphere [e.g., Lewis *et al.*, 2005; Singh *et al.*, 1995, 2003] where they affect the tropospheric ozone budget, are precursors to peroxy acetyl nitrate and, in the remote marine environment, represent a significant sink of the hydroxyl radical and thus the oxidizing capacity of the lower atmosphere [Folkins and Chatfield, 2000; Lewis *et al.*, 2005]. In remote marine air, oceanic sources and sinks of OVOCs are assumed to be significant in controlling air concentrations [Read *et al.*, 2012], although the magnitude and direction of the OVOC air-sea fluxes are a matter of debate [Beale *et al.*, 2013; Carpenter *et al.*, 2004; Heikes *et al.*, 2002; Marandino *et al.*, 2005; Taddei *et al.*, 2009; Williams *et al.*, 2004] largely as a consequence of extremely limited OVOC measurements in oceanic surface waters. Knowledge of OVOC production and loss rates, and an appreciation of the mechanisms involved in our oceans are also lacking.

[3] The carbonyl compounds acetaldehyde and acetone are thought to be produced in surface waters by the photo-

degradation of colored dissolved organic matter (CDOM) [de Bruyn *et al.*, 2011; Kieber *et al.*, 1990; Zhou and Mopper, 1997]. Modeling studies have suggested that there must be a large marine in situ source of methanol in the ocean mixed layer [Millet *et al.*, 2008], which is speculated to be biological in nature. For example, methanol has been observed in the gaseous headspace above laboratory phytoplankton cultures and in water surrounding intact macroalgal cells [Nightingale, 1991; Riemer, 1998]. Bacterial consortia are also thought to transform algal carbohydrates to methanol in the upper aerobic ocean [Sieburth and Keller, 1989].

[4] In this paper, we present results of incubation experiments conducted on seawater samples collected from contrasting regions of the Atlantic Ocean; from oligotrophic gyres to productive upwelling locations. This work was conducted to test our hypothesis that biological activity plays a significant role in controlling measured seawater concentrations and production rates of OVOCs in marine waters.

### 2. Experiments and Techniques

[5] Seawater samples were collected from the Atlantic Ocean (Table 1) during two research cruises (a) SOLAS ICON (Surface Ocean Lower Atmosphere Study, UK “The Impact of Coastal upwellings On the production of climate active gases”) aboard the Royal Research Ship (RRS) Discovery (D338, 15 April to 27 May 2009) and (b) AMT19 (Atlantic Meridional Transect cruise number 19) aboard RRS James Cook (JC039, 13 October to 1 December 2009). Samples were collected with 20 L Niskin bottles deployed on a rosette equipped with a Seabird conductivity, temperature, and depth sensors. The seawater was immediately transferred using Tygon tubing into acid-washed quartz incubation vessels (internal diameter 20 mm, length 300 mm) with Teflon screw caps (~300 mL).

[6] The first incubation experiment (Table 1 and Figures 1 and 2a) was carried out using surface seawater and water collected from 200 m depth, with parallel incubation vessels incubated under in situ light and dark conditions. The subsequent five photochemical incubation experiments (Table 1 and Figures 1 and 2b–2d) were carried out with surface seawater only at in situ temperatures starting predawn and ending after sunset. Typically, each experiment had four to six time points from which net change in concentrations were derived. Quartz incubation vessels were placed in on deck incubators with flowing surface seawater during the natural light incubations, and for dark experiments, the quartz vessels were placed in temperature-controlled ThermoTote incubators. Seawater concentrations of methanol, acetaldehyde, and acetone were determined at each time point using a membrane inlet system coupled to a proton transfer reaction mass spectrometer [Beale *et al.*, 2011]. In situ initial ( $T_0$ ) concentrations

Additional supporting information may be found in the online version of this article.

<sup>1</sup>Plymouth Marine Laboratory, Plymouth, Devon, UK.

Corresponding author: J. L. Dixon, Plymouth Marine Laboratory, Prospect Place, West Hoe, Plymouth, Devon, PL1 3DH, UK. (jod@pml.ac.uk)

**Table 1.** Summary of Incubation Experiments Using Surface (~5 m) Seawater<sup>a</sup>

Data	ΣUVA+UVB (Wm <sup>-2</sup> ·d)	CDOM <sub>α<sub>350</sub></sub> (m <sup>-1</sup> )	Light Attenuation Coefficient <i>K<sub>d</sub></i> (340 nm, m <sup>-1</sup> )	In situ (T <sub>s</sub> ) [OVOC] (nM)		ΣOVOC Oxidation <sup>c</sup> (nmol L <sup>-1</sup> d <sup>-1</sup> )		Fractional OVOC Loss Due to Microbial Oxidation <sup>f</sup> (%)				Biological Lifetime <sup>g</sup> (d)						
				M <sup>b</sup>	AA <sup>c</sup>	AA <sup>c</sup>	A <sup>d</sup>	M <sup>b</sup>	AA <sup>c</sup>	A <sup>d</sup>	M <sup>b</sup>	AA <sup>c</sup>	A <sup>d</sup>	M <sup>b</sup>	AA <sup>c</sup>	A <sup>d</sup>		
Mauritanian upwelling																		
08/05/2009	231	0.23 <sup>h</sup>	0.30	33 (62)	15 (11)	9 (1)	ND	ND	ND	10 (100)	92 (81)	0.5 (0.7)	7 (3)	0.25 (0.25)	82 (20)			
22/05/2009	164	0.22	0.38	64	12	4	ND	ND	ND	10	49	8	7	1.0	3			
Open ocean																		
NAG 26/10/2009	154	0.04 <sup>i</sup>	0.07	304	7	20	18 (56%)	41 (53%)	0.5 (63%)	4	100	11	17	0.17	41			
EU 5/11/2009	60	0.05 <sup>i</sup>	0.15	272	7	8	11 (56%)	36 (47%)	0.4 (34%)	3	72	8	26	0.19	23			
SAG 16/11/2009	192	0.04 <sup>i</sup>	0.08	109	5	11	10 (38%)	65 (46%)	0.2 (45%)	50	66	9	11	0.08	55			
ST 25/11/2009	158	0.80 <sup>i</sup>	0.31	136	6	2	14 (38%)	52 (48%)	0.4 (45%)	13	60	13	10	0.12	5			

<sup>a</sup>ND, not determined. Values in bold and brackets represent Experiment 1 but waters from 200 m out of the euphotic zone; NAG: North Atlantic Gyre, EU: Equatorial Upwelling, SAG: South Atlantic Gyre, ST: Southern temperate region.

<sup>b</sup>M, methanol,

<sup>c</sup>AA, acetaldehyde,

<sup>d</sup>A, acetone,


<sup>e</sup>Combined integrated oxidation (12 h light plus 12 h dark) with % oxidation in light indicated in the brackets,

<sup>f</sup>Calculated as microbial OVOC oxidation divided by gross production rates, unless production is negative when calculated as oxidation divided by net loss (assuming steady state),

<sup>g</sup>Calculated as in situ surface concentration divided by the surface microbial oxidation rate which for Experiments 1 and 2 are estimated as described in section 4.

<sup>h</sup>CDOM  $\alpha_{350}$  data from 09 May 2009 as not available from 08 May 2009.

<sup>i</sup>CDOM  $\alpha_{350}$  from a comparable date and location but from AMT18 in 2008 as data not available from AMT19 in 2009.




**Figure 1.** Location of incubation experiments (1–6) where the circle color represents surface temperature and the size of the circle the concentration of surface chlorophyll a (data in Table S1 in the supporting information). The background is a remotely sensed Moderate Resolution Imaging Spectroradiometer Aqua composite chlorophyll a image from October 2009 covering the start of AMT19 (courtesy of Natural Environment Research Council Earth Observation Data Acquisition and Analysis Service).

of OVOCs were also determined using the same analytical system (Table 1).

[7] Microbial oxidation rates of methanol, acetaldehyde, and acetone were determined using <sup>14</sup>C-labeled low nanomolar additions (<10% of in situ concentrations) and incubations of typically 1 h either in quartz microtubes placed in the light incubators or in the dark [Dixon *et al.*, 2011a]. OVOC uptake rates in nmol L<sup>-1</sup> h<sup>-1</sup> were calculated by multiplying the sample counts (nCimL<sup>-1</sup> h<sup>-1</sup>) by the specific activity of the <sup>14</sup>C compound (methanol 57.1 mCi mmol<sup>-1</sup>, acetaldehyde 50 mCi mmol<sup>-1</sup>, and acetone 30 Ci mmol<sup>-1</sup>). In order to calculate the total loss of OVOCs over 24 h due to microbial oxidation, rates were integrated over 12 h in light and dark experiments and combined (Table 1). Microbial oxidation was assumed to be the dominant biological removal pathway [Dixon *et al.*, 2011b], and OVOC uptake into microbial biomass was not determined during the experiments. Microbial loss rates calculated for methanol (as methanol oxidation rates) in the coastal upwelling station should be considered minimum values, as up to 57% of methanol can be assimilated into microbial biomass (rates are not known for acetaldehyde and acetone) [Dixon *et al.*, 2012].

### 3. Results

[8] Results from Experiment 1 in the Mauritanian coastal upwelling region (U1) are shown in Figure 2a. Methanol



**Figure 2.** The change in concentration of OVOCs over time ( $n \geq 4$ ) for (a) surface and 200 m water in Experiment 1 in the Mauritanian upwelling (Table 1) and (b) methanol, (c) acetaldehyde, and (d) acetone for surface waters of Experiments 2–6 (Table 1). The legend for Figures 2b–2d is shown in Figure 2b.

showed a net production of  $44 \text{ nmol L}^{-1}$  in surface waters over 12 h incubation in the light, but zero change under dark incubations conditions giving a daily net production of  $44 \text{ nmol L}^{-1} \text{ d}^{-1}$ . Methanol showed no overall change in concentration in 200 m samples under light and dark conditions, i.e.,  $0 \text{ nmol L}^{-1} \text{ d}^{-1}$ . Surface acetaldehyde showed net production of 7.6 in the light, but a loss of  $2.7 \text{ nmol L}^{-1}$  in the dark, giving a daily net production rate of  $4.9 \text{ nmol L}^{-1} \text{ d}^{-1}$ . Acetaldehyde from 200 m when exposed to the light for 12 h showed a net increase of  $9.4 \text{ nmol L}^{-1}$ , with no overall change in concentration in the dark resulting in a daily net production rates of  $9.4 \text{ nmol L}^{-1} \text{ d}^{-1}$ . However, water sampled from 200 m would not receive any natural light; thus, in situ acetaldehyde production rates are assumed to be  $\sim 0 \text{ nmol L}^{-1} \text{ d}^{-1}$ . Acetone always showed net production in surface waters of 18.5 and  $7.5 \text{ nmol L}^{-1}$  in light and dark incubations, respectively, resulting in daily net production rates of  $26 \text{ nmol L}^{-1} \text{ d}^{-1}$ . Acetone production in 200 m water was 12.8 and  $3.5 \text{ nmol L}^{-1}$  in light and dark incubations, respectively, resulting in a daily net production rate of  $7.0 \text{ nmol L}^{-1} \text{ d}^{-1}$ , i.e., twice the 200 m dark rate. The difference in production rates between 200 m water incubated in light and dark conditions of 9.4 and  $9.3 \text{ nmol L}^{-1}$  for acetaldehyde and acetone respectively suggests that these deeper upwelling waters contain acetaldehyde and acetone precursors; possibly from sinking phytoplankton detritus derived from the above highly productive waters. Integrated microbial OVOC oxidation rates were not determined during Experiment 1.

[9] OVOC net production rates from Experiments 2–6 are shown in Figures 2b–2d. In the low chlorophyll a ( $< 0.1 \mu\text{g L}^{-1}$ ), surface waters of the Atlantic gyres and equatorial upwelling (EU, Experiments 3–5, Figure 1, and Table S1) methanol shows net daily losses of between 22 and  $428 \text{ nmol L}^{-1} \text{ d}^{-1}$ . For higher chlorophyll stations

( $> 1.0 \mu\text{g L}^{-1}$ ) of the southern temperate region (ST) and Mauritanian coastal upwelling (U2), methanol shows net daily production rates of 89 and  $93 \text{ nmol L}^{-1} \text{ d}^{-1}$ , respectively. Acetaldehyde net production rates ranged between 13 and  $35 \text{ nmol L}^{-1} \text{ d}^{-1}$  in all regions sampled, except the North Atlantic gyre (NAG), which showed no overall change in concentration over 24 h. Acetone net daily production rates were relatively modest compared to acetaldehyde and ranged between 2 and  $4 \text{ nmol L}^{-1} \text{ d}^{-1}$  in open ocean low chlorophyll regions (Experiments 3–5, Table 1). In higher chlorophyll waters of coastal upwellings, net acetone production rates up to  $26 \text{ nmol L}^{-1} \text{ d}^{-1}$  were found (Figure 2).

[10] Daily integrated methanol loss rates due to microbial oxidation ranged between 10 and  $18 \text{ nmol L}^{-1} \text{ d}^{-1}$  for Experiments 3–6 (Table 1) and were highest in NAG. Microbial acetaldehyde oxidation rates were relatively higher at  $36\text{--}65 \text{ nmol L}^{-1} \text{ d}^{-1}$  with highest loss rates in South Atlantic gyre water (SAG, Table 1). In contrast, microbial acetone oxidation rates were modest at  $0.2\text{--}0.5 \text{ nmol L}^{-1} \text{ d}^{-1}$ .

#### 4. Discussion

[11] Methanol showed a daily net production of  $44 \text{ nmol L}^{-1} \text{ d}^{-1}$  in surface waters of U1 (Figure 2a). We do not have concurrent rates of microbial methanol oxidation but estimate that rates were  $\sim 5 \text{ nmol L}^{-1} \text{ d}^{-1}$  (based on surface water rates on 9th May 2009). Surface methanol gross production rates were therefore estimated at  $\sim 49 \text{ nmol L}^{-1} \text{ d}^{-1}$ , with microbial loss processes accounting for  $\sim 10\%$  reduction in methanol production rates at U1. Methanol production and consumption rates balanced in seawater samples from 200 m (dark and below the mixed layer). Microbial methanol oxidation rates of  $\sim 20 \text{ nmol L}^{-1} \text{ d}^{-1}$  from a day later at 200 m were much larger than surface values, which suggests that gross methanol production was  $\sim 20 \text{ nmol L}^{-1} \text{ d}^{-1}$  ( $\sim 41\%$

of surface rates). The ICON experiment was conducted in a Lagrangian framework with SF<sub>6</sub> and <sup>3</sup>He measurements confirming that we sampled the same water mass on the 8th and 9th May 2009. The biological lifetime of methanol in seawater was 7 days in surface waters and ~3 days in 200 m water from U1 (Table 1, calculated by comparing in situ concentrations (T<sub>0</sub>) with daily integrated microbial oxidation rates).

[12] Microbial acetaldehyde oxidation rates were much higher than for methanol at 60 and 44 nmol L<sup>-1</sup> d<sup>-1</sup> in surface and 200 m waters, respectively (on 9th May 2009). Assuming these rates are comparable to those of Experiment 1 (8th May 2009), we estimate gross surface acetaldehyde production rates of approximately 65 and 54 nmol L<sup>-1</sup> d<sup>-1</sup> for surface and 200 m water, respectively, with microbial oxidation largely controlling acetaldehyde concentrations. The biological lifetime of acetaldehyde in U1 is ~6 h in surface and 200 m water.

[13] Microbial acetone oxidation rates (also determined on 9th May 2009) were comparatively low at 0.11 and 0.05 nmol L<sup>-1</sup> d<sup>-1</sup> in surface and 200 m water, respectively. We therefore estimate gross surface acetone production rates to be very similar to net rates, at fractionally over 26 and 7 nmol L<sup>-1</sup> d<sup>-1</sup>, for surface and 200 m, respectively. Thus, for acetone, microbial losses due to oxidation are minor. The biological lifetime of acetone at U1 is >80 days in surface waters and ~20 days at 200 m, where in situ acetone concentrations were relatively low at 9 and 1 nM, respectively (Table 1). The microbial acetone loss rates reported in this study are substantially lower than those of a coastal station in the Pacific Ocean, where biotic losses were estimated at ~2.7 d<sup>-1</sup> [de Bruyn *et al.*, 2013], which if multiplied by our in situ acetone concentrations (Table 1) suggest biologically driven loss rates between 2.7 and 24 nmol L<sup>-1</sup> d<sup>-1</sup>. This large difference could represent differences in location. Alternatively, the large spike of fully deuterated (d-6) acetone (4–26 nM or 44–288% of our in situ acetone surface concentrations determined at station 1, Table 1) used by de Bruyn *et al.* [2013] may overestimate losses of acetone at in situ concentrations.

[14] The large net losses of methanol found in surface waters of NAG and EU cannot be fully explained (Figure 2b). Microbial methanol oxidation accounts for 3–4% of the total daily loss of methanol at these stations (Table 1). This suggests other removal mechanisms, perhaps via microbial uptake of methanol (given the high surface concentrations of 272–304 nM, Table 1) and subsequent excretion of overflow metabolites as other organic intermediates. However, in the SAG, where surface methanol concentrations are approximately threefold lower, measured microbial losses account for ~50% of methanol loss. There is net production of methanol in higher chlorophyll regions of the ST and U2. For ST waters, integrated microbial oxidation rates were 14 nmol L<sup>-1</sup> d<sup>-1</sup> (Table 1). Thus, gross methanol production in this biologically active region is estimated at 103 nmol L<sup>-1</sup> d<sup>-1</sup>, with a biological lifetime more similar to the coastal upwelling regions of 10 days. Methanol production rates in U2 can also be corrected for microbial oxidation estimated at 9.8 nmol L<sup>-1</sup> d<sup>-1</sup> (from 21st May 2009 in the same upwelling water mass) resulting also in methanol gross production rates of 103 nmol L<sup>-1</sup> d<sup>-1</sup>, which is over double the previous surface rate estimate in U1, but the biological lifetime is the same at ~7 days (Table 1).

This increase in methanol production rates between U1 and U2 could be due to an increase in rates of primary and bacterial productivity of over 80% (Table S1), if the main source of methanol is either from phytoplankton cells [Nightingale, 1991; Riemer, 1998] or from bacterial breakdown of algal products [Sieburth and Keller, 1989]. It is also interesting that net production of methanol negatively correlates with surface temperature ( $r = -0.854$ ,  $P > 0.05$ ).

[15] Acetaldehyde net production rates can be corrected for microbial losses due to oxidation giving estimated gross production rates (assuming microbial oxidation is the major loss mechanism in seawater) of 41, 50, 98, and 87 nmol L<sup>-1</sup> d<sup>-1</sup> for the NAG, EU, SAG, and ST waters, respectively. Estimated microbial oxidation rates of acetaldehyde for U2 of 12 nmol L<sup>-1</sup> d<sup>-1</sup> (on 21st May 2009) result in a relatively lower gross production of 25 nmol L<sup>-1</sup> d<sup>-1</sup>. This is lower than observed for acetaldehyde in U1, principally due to a higher microbial acetaldehyde oxidation rate of 60 nmol L<sup>-1</sup> d<sup>-1</sup>. This trend between the two upwelling filaments, i.e., U1 and U2 for acetaldehyde is opposite to that found for methanol, and could reflect decreasing microbial requirement for acetaldehyde as a preferential energy source in U2, perhaps due to an elevated range or quantity of organic sources from enhanced microbial activity. Overall, the biological acetaldehyde lifetimes in open oceanic waters are 2–5 h (Experiments 3–6 in Table 1), but up to 24 h in upwelling filaments.

[16] Microbial acetaldehyde oxidation rates in surface waters are always equal to, or significantly greater than net production rates, implying strong microbial control of acetaldehyde concentrations in seawater (Table 1). High atmospheric acetaldehyde at Cape Verde in the Atlantic Ocean [Read *et al.*, 2012] compared to the Indian [Wisthaler *et al.*, 2002] and Pacific Oceans [Singh *et al.*, 2003] have been attributed to high photoproduction of acetaldehyde in the biologically active upwelling regions of the West African coast. The fraction of acetaldehyde gross production attributed to photochemical production ranges between 16% and 68% for the coastal upwelling and NAG locations, closest to the Cape Verde (Table S2). Our results also demonstrate the significance of surface ocean microbes in reducing and controlling oceanic acetaldehyde concentrations and could account for a lower global oceanic source of 17 Tgy<sup>-1</sup> based on in situ seawater concentrations [Beale *et al.*, 2013], compared to estimates of 57–175 Tgy<sup>-1</sup> based largely on modeled air data [Millet *et al.*, 2010; Singh *et al.*, 2004].

[17] In the open ocean, acetone losses due to microbial oxidation (Table 1) are relatively small resulting in gross production rates approximately 9–13% higher than net rates (Table 1, assuming microbial oxidation is the dominant loss pathway). Although microbial acetone oxidation rates from U2 (on 21st May 2009) were higher at 1.2 nmol L<sup>-1</sup> d<sup>-1</sup> (gross production of 15.9 nmol L<sup>-1</sup> d<sup>-1</sup>), elevated net production rates suggest a reduced role for bacteria in removing acetone from surface seawater. Our results contrast with those from coastal Pacific experiments [de Bruyn *et al.*, 2013] as discussed previously. Comparison of in situ acetone concentrations (T<sub>0</sub>, Table 1) with microbial oxidation rates suggests that acetone has a biological lifetime of ~41–55 days in the oligotrophic gyres, ~23 days in EU water, ~5 days in ST waters, and 3–82 days in the highly productive coastal upwelling waters. Acetone production is always greater when exposed to light compared to dark conditions (Figure 2d) and photochemical production is estimated at 48–100% of gross production

(Table S2), except in the NAG where production in the dark is approximately equal to that in the light. This could be due to elevated microbial acetone oxidation rates in the light (Table 1). Acetone oxidation rates correlate with bacterial production ( $r=0.856$ ,  $P>0.05$ ), and Beale *et al.* [2013] report a negative relationship between acetone seawater concentrations and bacterial production. Thus, although acetone oxidation rates are low, these relationships suggest that as bacterial production increases, so does the rate of microbial acetone oxidation, leading to a reduction in the in situ concentration of acetone. Net acetone production rates also correlate with the numbers of picoeukaryotic cells ( $r=0.926$ ,  $P>0.05$ ), while gross production rates normalized to CDOM ( $\alpha_{350}$ , Table 1) explain approximately 67% of the observed spatial variability (reducing it from ninefold to threefold). Thus, we suggest that acetone production is mainly photochemical [de Bruyn *et al.*, 2011; Kieber *et al.*, 1990; Zhou and Mopper, 1997] and seems to be related to the UV breakdown of CDOM originating from the number of picoeukaryotic cells.

[18] Spatial differences in the daily UV dose, CDOM ( $\alpha_{350}$ ) and the diffuse light attenuation coefficient  $K_d$  (340 nm) are shown in Table 1. However, for methanol and acetaldehyde, normalizing the gross production rates by any of the aforementioned parameters in isolation or combination does not help explain the spatial variability.

[19] Microbial oxidation rates (integrated to 1 m) are compared to air-sea flux estimates from the same cruises [Beale *et al.*, 2013; Beale, 2011] (Table S3). Comparisons suggest that for methanol, microbial oxidation (loss) is of the same order of magnitude as the air-sea flux. For acetaldehyde, the biological loss is an order of magnitude higher than the air-sea flux, and for acetone biological losses due to oxidation are an order of magnitude lower than the air-sea flux.

[20] We conclude that in productive coastal upwelling filaments, the gross production of methanol, acetaldehyde, and acetone is 49–103, 25–65, and 16–26  $\text{nmol L}^{-1} \text{d}^{-1}$ , respectively. Microbial oxidation reduces net surface production rates by 10%, 50–92%, and 0.5–8% for methanol, acetaldehyde, and acetone, respectively. Biological lifetimes vary between  $\leq 1$  day for acetaldehyde, to approximately 7 days for methanol and up to  $\sim 80$  days for acetone.

[21] In oceanic regions, methanol largely showed a net loss in the incubations, of which only 3–4% could be attributed to microbial oxidation rates in the NAG and EU regions, although this increased to 50% for the SAG. Gross methanol production of 103  $\text{nmol L}^{-1} \text{d}^{-1}$  in the ST eutrophic region (methanol biological lifetime  $\sim 10$  days) was highly comparable to higher chlorophyll *a* waters of the Mauritanian upwelling filaments. Acetaldehyde gross production rates in open ocean environs varied between 41 and 98  $\text{nmol L}^{-1} \text{d}^{-1}$  and were highest in the SAG. In agreement with coastal upwelling experiments, surface concentrations were controlled by microbial loss processes (60–100%) with a biological lifetime of 2–5 h. In contrast, acetone gross production rates were relatively low between 2.2 and 4.5  $\text{nmol L}^{-1} \text{d}^{-1}$ , with microbial oxidation reducing production rates by 8–13% (Table 1). The biological lifetime of surface acetone in remote low chlorophyll *a* ( $<0.1 \mu\text{g L}^{-1}$ ) regions was 23–55 days, which reduced to a minimum of 3–5 days in productive waters (chlorophyll *a*  $>0.5 \mu\text{g L}^{-1}$ ).

[22] Our results suggest that methanol photochemical production is relatively insignificant; and concentrations are controlled by microbial oxidation and overflow metabolism,

surface seawater acetaldehyde concentrations are largely controlled by microbial losses and photochemical production, and acetone production is mainly photochemical with relatively low microbial loss rates. We have thus highlighted the importance of the ocean in both the production and consumption of these atmospherically important OVOC compounds and have highlighted significant compound and spatial differences.

[23] **Acknowledgments.** We thank all the scientists, officers, and crew of the RRS Discovery and James Cook during D338 and JC029. We also thank C. Widdicombe for primary production rates and chlorophyll *a* concentrations, S. Sargeant for bacterial production rates (for Experiments 3–6), T. Smyth for UV and  $K_d$  (AMT) data, V. Kitidis for  $K_d$  and CDOM data (ICON), and G. Tilstone for CDOM (AMT) data. This study was supported by the UK Natural Environmental Research Council (NERC) via UK SOLAS and Oceans 2025 funding for Plymouth Marine Laboratory. This is contribution 239 of the AMT program.

[24] The Editor thanks two anonymous reviewers for their assistance in evaluating this paper.

## References

- Beale, R. (2011), Quantification of oxygenated volatile organic compounds in seawater, PhD thesis, University of East Anglia: Norwich, UK.
- Beale, R., P. S. Liss, J. L. Dixon, and P. D. Nightingale (2011), Quantification of oxygenated volatile organic compounds in seawater by membrane inlet-proton transfer reaction/mass spectrometry, *Anal. Chim. Acta*, 706, 128–134, doi:10.1016/j.aca.2011.08.023.
- Beale, R., J. L. Dixon, S. R. Arnold, P. S. Liss, and P. D. Nightingale (2013), Methanol, acetaldehyde and acetone in the surface waters of the Atlantic Ocean, *J. Geophys. Res. Oceans*, doi:10.1002/jgrc.20322.
- Carpenter, L. J., A. Lewis, J. R. Hopkins, K. A. Read, I. D. Longley, and M. W. Gallagher (2004), Uptake of methanol to the North Atlantic ocean surface, *Global Biogeochem. Cycles*, 18, GB4027, doi:10.1029/2004GB002294.
- De Bruyn, W. J., C. D. Clark, L. Pagel, and C. Takehara (2011), Photochemical production of formaldehyde, acetaldehyde and acetone from chromophoric organic matter in coastal waters, *J. Photochem. Photobiol. A: Chem.*, 226, 16–22.
- De Bruyn, W. J., C. D. Clark, L. Pagel, and H. Singh (2013), Loss rates of acetone in filtered and unfiltered coastal seawater, *Mar. Chem.*, 150, 39–44, doi:10.1016/j.marchem.2013.01.003.
- Dixon, J. L., R. Beale, and P. D. Nightingale (2011a), Microbial methanol uptake in northeast Atlantic waters, *ISME J.*, 5, 704–716.
- Dixon, J. L., R. Beale, and P. D. Nightingale (2011b), Rapid biological oxidation of methanol in the tropical Atlantic: Significance as a microbial carbon source, *Biogeosciences*, 8, 2707–2716, doi:10.5194/bg-8-2707-2011.
- Dixon, J. L., S. Sargeant, P. D. Nightingale, and J. C. Murrell (2012), Gradients in microbial methanol uptake: Productive coastal upwelling waters to oligotrophic gyres in the Atlantic Ocean, *ISME J.*, 7, 568–580, doi:10.1038/ismej.2012.130.
- Folkens, I., and R. Chatfield (2000), Impact of acetone on ozone production and OH in the upper troposphere at high NO<sub>x</sub>, *J. Geophys. Res.*, 105, 11,585–11,599.
- Heikes, B. G., et al. (2002), Atmospheric methanol budget and ocean implication, *Global Biogeochem. Cycles*, 16(4), 1133, doi:10.1029/2002GB001895.
- Kieber, R. J., X. L. Zhou, and K. Mopper (1990), Formation of carbonyl compounds from UV-induced photodegradation of humic substances in natural waters: Fate of riverine carbon in the sea, *Limnol. Oceanogr.*, 35, 1503–1515.
- Lewis, A. C., J. R. Hopkins, L. J. Carpenter, J. Stanton, K. A. Read, and M. J. Pilling (2005), Sources and sinks of acetone, methanol, and acetaldehyde in North Atlantic marine air, *Atmos. Chem. Phys.*, 5, 1963–1974.
- Marandino, C. A., W. J. De Bruyn, S. D. Miller, M. J. Prather, and E. S. Saltzman (2005), Oceanic uptake and the global atmospheric acetone budget, *Geophys. Res. Lett.*, 32, L15806, doi:10.1029/2005GL023285.
- Millet, D. B., et al. (2008), New constraints on terrestrial and oceanic sources of atmospheric methanol, *Atmos. Chem. Phys.*, 8, 6887–6905.
- Millet, D. B., et al. (2010), Global atmospheric budget of acetaldehyde: 3-D model analysis and constraints from in situ and satellite observations, *Atmos. Chem. Phys.*, 10, 3405–3425.
- Nightingale, P. D. (1991), Low molecular weight halocarbons in seawater, PhD thesis, University of East Anglia: Norwich, UK.
- Read, K. A., L. J. Carpenter, S. R. Arnold, R. Beale, P. D. Nightingale, J. R. Hopkins, A. C. Lewis, J. D. Lee, L. Mendes, and S. J. Pickering

- (2012), Multiannual observations of acetone, methanol, and acetaldehyde in remote tropical Atlantic air: Implications for atmospheric OVOC budgets and oxidative capacity, *Environ. Sci. Technol.*, *46*, 11,028–11,039.
- Rierner, D. (1998), Marine and terrestrial sources of reactive volatile organic compounds and their impact on the tropospheric ozone chemistry of the Earth, PhD thesis, Univ of Miami, Miami, Florida, USA.
- Sieburth, J. M., and M. D. Keller (1989), Methylaminotrophic bacteria in xenic nanoalgal cultures: Incidence, significance, and role of methylated algal osmoprotectants, *Biol. Oceanogr.*, *6*, 383–395.
- Singh, H. B., M. Kanakidou, P. J. Crutzen, and D. J. Jacob (1995), High concentrations and photochemical fate of oxygenated hydrocarbons in the global troposphere, *Nature*, *378*, 51–54.
- Singh, H. B., A. Tabazadeh, M. J. Evans, B. D. Field, D. J. Jacob, G. Sachse, J. H. Crawford, R. Shetter, and W. H. Brune (2003), Oxygenated volatile organic chemicals in the oceans: Inferences and implications based on atmospheric observations and air-sea exchange models, *Geophys. Res. Lett.*, *30*(16), 1862, doi:10.1029/2003GL017933.
- Singh, H. B., et al. (2004), Analysis of the atmospheric distribution, sources and sinks of oxygenated volatile organic chemicals based on measurements over the Pacific during TRACE-P, *J. Geophys. Res.*, *109*, D15S07, doi:10.1029/2003JD003883.
- Taddei, S., P. Toscano, B. Gioli, A. Matese, F. Miglietta, F. P. Vaccari, A. Zaldei, T. Custer, and J. Williams (2009), Carbon dioxide and acetone air – sea fluxes over the Southern Atlantic, *Environ. Sci. Technol.*, *43*, 5218–5222.
- Williams, J., R. Holzinger, V., Gros, X. Xu, E. Atlas, and D. W. R. Wallace (2004), Measurements of organic species in air and seawater from the tropical Atlantic, *Geophys. Res. Lett.*, *31*, L23S06, doi:10.1029/2004GL020012.
- Wisthaler, A., A. Hansel, R. R. Dickerson, and P. J. Crutzen (2002), Organic trace gas measurements by PTR-MS during INDOEX 1999, *J. Geophys. Res.*, *107*(D19), 4402, doi:10.1029/2001JD000567.
- Zhou, X., and K. Mopper (1997), Photochemical production of low molecular-weight carbonyl compounds in seawater and surface microlayer and their air – sea exchange, *Mar. Chem.*, *56*, 201–213.