



MarLIN

Marine Information Network

Information on the species and habitats around the coasts and sea of the British Isles

Corallina officinalis and *Mastocarpus stellatus* on exposed to moderately exposed lower eulittoral rock

MarLIN – Marine Life Information Network
Marine Evidence-based Sensitivity Assessment (MarESA) Review

Dr Heidi Tillin & Dr Harvey Tyler-Walters

2018-09-18

A report from:

The Marine Life Information Network, Marine Biological Association of the United Kingdom.

Please note. This MarESA report is a dated version of the online review. Please refer to the website for the most up-to-date version [<https://www.marlin.ac.uk/habitats/detail/1180>]. All terms and the MarESA methodology are outlined on the website (<https://www.marlin.ac.uk>)


This review can be cited as:

Tillin, H.M. & Tyler-Walters, H., 2018. [*Corallina officinalis*] and [*Mastocarpus stellatus*] on exposed to moderately exposed lower eulittoral rock. In Tyler-Walters H. and Hiscock K. (eds) *Marine Life Information Network: Biology and Sensitivity Key Information Reviews*, [on-line]. Plymouth: Marine Biological Association of the United Kingdom. DOI <https://dx.doi.org/10.17031/marlinhab.1180.1>



The information (TEXT ONLY) provided by the Marine Life Information Network (MarLIN) is licensed under a Creative Commons Attribution-Non-Commercial-Share Alike 2.0 UK: England & Wales License. Note that images and other media featured on this page are each governed by their own terms and conditions and they may or may not be available for reuse. Permissions beyond the scope of this license are available [here](#). Based on a work at www.marlin.ac.uk

(page left blank)



Researched by Dr Heidi Tillin & Dr Harvey Tyler-Walters

Refereed by This information is not refereed.

Summary

☰ UK and Ireland classification

EUNIS 2008 A1.1221

Corallina officinalis and *Mastocarpus stellatus* on exposed to moderately exposed lower eulittoral rock

JNCC 2015 LR.HLR.FR.Coff.Coff

Corallina officinalis and *Mastocarpus stellatus* on exposed to moderately exposed lower eulittoral rock

JNCC 2004 LR.HLR.FR.Coff.Coff

Corallina officinalis and *Mastocarpus stellatus* on exposed to moderately exposed lower eulittoral rock

1997 Biotope

🔍 Description

Exposed lower eulittoral rock or moderately exposed lower eulittoral vertical rock that supports a dense turf of the red seaweed *Corallina officinalis*, often on wave surged rocky slopes. There is usually a low abundance of other turf-forming red seaweeds such as *Lomentaria articulata*,

Mastocarpus stellatus, *Palmaria palmata* and *Osmundea pinnatifida*. Other seaweeds that occur in low abundance includes the wrack *Himanthalia elongata* and the kelp *Laminaria digitata*, while the brown seaweed *Leathesia difformis* can be found growing on and around the other seaweeds. Green seaweeds such as *Ulva intestinalis*, *Ulva lactuca* and *Cladophora rupestris* are also present. The coralline turf creates a micro-habitat for small animals such as the colonial tube-building polychaete *Spirobranchus* sp. and the barnacle *Semibalanus balanoides*. The mussel *Mytilus edulis* is often found in small cracks and crevices while the sponges *Halichondria panicea* and *Hymeniacion perleve* can be found in shaded areas or on overhangs. The limpets *Patella ulyssiponensis* and *Patella vulgata* can be found on the bedrock underneath the turf. The brown seaweed *Bifurcaria bifurcata* and the barnacle *Balanus perforatus* may occur in the extreme south-west.

This community usually forms a distinct band just above the kelp zone (Ala, Ala.Ldig or Ldig). It can be found below the barnacle and *Patella vulgata* dominated biotopes (Cht; Sem.Sem; Sem.FvesR). If there are many of the characteristic *Himanthalia elongata* buttons present in early spring careful comparison with the Him biotope should be made. (Information from Connor *et al.*, 2004; JNCC, 2015).

↓ Depth range

Lower shore

🏛️ Additional information

-

✓ Listed By

- none -

🔗 Further information sources

Search on:



Sensitivity review

Sensitivity characteristics of the habitat and relevant characteristic species

Corallina officinalis is the dominant characterizing species within this biotope LR.HLR.FR.Coff.Coff and LR.HLR.FR.Coff.Puly. *Corallina officinalis* forms a dense turf that provides substratum and refuges for a diverse epifauna. As the biotope characterization and many of the associated species are dependent on the presence of *Corallina officinalis* the sensitivity assessments specifically considers this species as both a key structuring and characterizing species. Other turf-forming algae such as *Lomentaria articulata*, *Mastocarpus stellatus*, *Palmaria palmata* and *Osmundea pinnatifida* occur in low abundances. The green seaweeds *Ulva intestinalis*, *Ulva lactuca* and *Cladophora rupestris* are present as well and assessments describe the sensitivity of the red and green species in general terms. *Himantalia elongata* also occurs but in lower abundances than the LR.HLR.FR.Him biotope, therefore, this species is considered specifically within the assessment where its sensitivity differs and it may come to dominate the biotope. Gastropods *Littorina littorea*, *Patella vulgata* and *Patella ulyssiponensis* are significant grazers in the eulittoral zone and, by preferentially grazing on foliose red and green algae, structure the biotope allowing *Corallina* sp. to dominate. They are, therefore, included as important structural species. A number of invertebrates are present on the bedrock underneath the coralline turf, including the barnacle *Semibalanus balanoides*, the mussel *Mytilus edulis*, the sponges *Halichondria panicea* and *Hymeniacidon perleve*, the anemone *Actinia equina*. These common rocky shore species contribute to species diversity and ecological function within the biotope but are not considered to be important structural or functional species and are only generally referred to within the assessments. Epiphytic grazers, such as amphipods, isopods and small gastropods probably keep the turf free of epiphytic algae and are important structural species. Due to lack of evidence the sensitivity of this group is considered only generally where the pressures may impact on this biotope. Temporal variation of the abundances of the characterizing species within this biotope may lead to biotope reversion between LR.HLR.FR.Coff.Coff, LR.HLR.FR.Coff.Puly and LR.HLR.FR.Him as these contain broadly similar species and occur in similar conditions (Connor *et al.*, 2004).

Resilience and recovery rates of habitat

Where this biotope is impacted by pressures, recovery of the key structuring and characterizing species *Corallina officinalis* will require either regrowth from surviving holdfast or basal crusts or recolonization by propagules. The crustose holdfast or base is perennial and grows apically (continuous growth at tips), similar to encrusting corallines such as *Lithothamnium* sp. The basal crust may grow continuously until stimulated to produce fronds (Littler & Kauker 1984; Colhart & Johanssen 1973). Littler & Kauker (1984) suggest that the crustose bases are an adaptation to resist grazing and desiccation whereas the fronds are adapted for higher primary productivity and reproduction. The basal crusts are tougher than the upright fronds (requiring a pressure of 94 g/mm² to penetrate, compared to 43 g/mm² respectively). Regeneration of the basal crusts provides a more rapid route to recovery than recolonization. Experiments in the intertidal in southern California found that areas scraped back to crusts recovered four times more rapidly than sterilised plots where the crusts were removed (Littler & Kauker, 1994).

In culture, *Corallina officinalis* fronds exhibited an average growth rate of 2.2 mm/month at 12 and 18°C. The growth rate was only 0.2 mm/month at 6°C and no growth was observed at 25°C (Colhart & Johanssen 1973). Similarly Blake & Maggs (2003) observed much higher growth rates

of 2 mm/month over 6 months starting from September in *Corallina officinalis* grown in Strangford Lough (Northern Ireland) at 5 and 10 m depth, these rates are similar to those observed by Andrade & Johansen, (1980) in winter in New Hampshire. The evidence for growth rate suggests that to achieve a height of 10 cm the turf would be at least 4 years old (probably older as higher temperatures appear to slow growth. A lower level turf of, for example 5 cm, could theoretically be achieved within 2 years.

Where the bases are removed, recovery will depend on recolonization. Areas that are cleared during the reproductive period have the potential to be rapidly colonized. *Corallina officinalis* was shown to settle on artificial substances within 1 week of their placement in the intertidal in New England summer (Harlin & Lindbergh, 1977). However, settlement plates laid out in the autumn were not recolonized until the next spring. In the lower rocky intertidal in southern California dominated by *Corallina officinalis* with foliose overstorey algae present, Littler & Kauker (1984) experimentally cleared plots and followed the recovery for 12 months. Some areas were scraped allowing the basal crusts to remain whereas others were completely sterilised (removal of all material and surfaces then scorched with a blow torch to remove bases). In scraped plots, up to 15% cover of *Corallina officinalis* fronds returned within 3 months after removal of fronds and all other epiflora/fauna (Littler & Kauker, 1984) while in sterilized plots (all basal crusts removed) appearance of articulated fronds occurred 6 months following clearance. At the end of the 12 month observation period, *Corallina officinalis* cover had increased to approximately 18% in plots where basal crusts remained and to approximately 10% in sterilised plots. Similarly, Bamber & Irving (1993) reported that new plants grew back in scraped transects within 12 months, although the resistant crustose bases were probably not removed.

Once established turfs of *Corallina* spp. can persist for a long time, surveys of rocky intertidal ledges at Hinkley point, Somerset in England have found that the patches mapped in the 1980s (Bamber & Irving, 1993) had not changed position when resurveyed 18 years later (Burdon *et al.*, 2009). It has been speculated but not definitively demonstrated that turf-forming algae and canopy-forming algae may represent alternate stable states on temperate rocky shores and a shift in balance to the alternate state may prevent recovery. Some potential mechanisms for inhibition of canopy-forming species are space pre-emption by turfs that prevent recruitment of taller algae (Perkol-Finkel & Airoidi, 2010, Kennelly, 1987) due to the coverage of suitable rock surfaces and the presence of sediments within the turf (Airoidi, 2003). Clearance experiments on rocky, intertidal shores in Southern California (Sousa, 1979) found that *Ulva* species which have a longer reproductive season could colonize cleared areas preventing the establishment of perennial red algae. However, grazing by crabs removed the green algae (Sousa, 1979), highlighting the potential importance of grazers, particularly littorinids, to the re-establishment of this biotope.

Resilience assessment. No direct evidence was found for the age of individual *Corallina* crusts, the longevity of turfs, or the time to recover from basal crusts or sterilised plots to a full dense cover. New crustose bases may recruit and develop quickly but the formation of new fronds from these bases and recovery of original cover may take longer. Once a coralline turf has developed it will probably be colonized by epiphytic invertebrates such as harpacticoids, amphipods and isopods relatively quickly from the surrounding area. Therefore, the biotope would be recognizable once the coralline turf has regrown, which is likely to be quite rapid if the resistant crustose bases remain. The clearance experiments by Littler & Kauker (1984) suggest that recovery of a dense turf cover whether basal crusts remained or were totally removed would require more than 2 years. Presumably, as crusts can grow in all directions percentage cover is not a linear function and that gap closure would speed up with greater cover. Recruitment of associated species of red algae is probably equally rapid, and once the algal turf has developed most of the epiphytic invertebrates

would colonize quickly. The ephemeral green algae associated with the biotope are opportunist colonizers of gaps and would be expected to recover within a year. Limpets and littorinids could recover through migration but where populations are removed over a larger area recolonization by larvae would be required. Recruitment through larvae may be episodic and recovery to former population structure may require > 2 years. More detailed information on recovery of associated species can be found in the information for biotopes where these are the key characterizing species.

Resilience of the biotope is assessed as '**High**' where resistance is 'High' (no significant impact) or 'Medium' (where <25 % of *Corallina officinalis* fronds are removed and bases remain) based on regrowth from the basal crusts and vegetative growth from surrounding turfs and repair, migration or recolonization of associated species. Where resistance is 'Low' or 'None' then resilience is assessed as '**Medium**', between 2 -10 years but towards the lower end of that range for *Corallina officinalis* and the associated species. Where perturbations have a large spatial footprint with the widespread removal of crusts over a large area then the development of an alternate state emerging with dominance by canopy-forming algae is a possibility. In such an instance recovery could take much longer and depend on active management or further perturbations. No evidence was found however to determine when such shifts might occur.

NB: The resilience and the ability to recover from human induced pressures is a combination of the environmental conditions of the site, the frequency (repeated disturbances versus a one-off event) and the intensity of the disturbance. Recovery of impacted populations will always be mediated by stochastic events and processes acting over different scales including, but not limited to, local habitat conditions, further impacts and processes such as larval-supply and recruitment between populations. Full recovery is defined as the return to the state of the habitat that existed prior to impact. This does not necessarily mean that every component species has returned to its prior condition, abundance or extent but that the relevant functional components are present and the habitat is structurally and functionally recognisable as the initial habitat of interest. It should be noted that the recovery rates are only indicative of the recovery potential.

Hydrological Pressures

	Resistance	Resilience	Sensitivity
Temperature increase (local)	High Q: High A: Medium C: High	High Q: High A: High C: High	Not sensitive Q: High A: Medium C: High

Species found in the intertidal are exposed to extremes of high and low air temperatures during periods of emersion. They must also be able to cope with sharp temperature fluctuations over a short period of time during the tidal cycle. In winter, air temperatures are colder than the sea, conversely in summer air temperatures are much warmer than the sea. Species that occur in this intertidal biotope are therefore generally adapted to tolerate a range of temperatures, although the timing of site-specific factors such as low tides will influence local acclimation. For intertidal species, increased temperatures may also result in desiccation when exposed (see changes in emergence pressure).

The key characterizing species, *Corallina officinalis* has a cosmopolitan distribution (Guiry & Guiry, 2015) and throughout its range experiences wide variation in temperatures (although local populations may be acclimated to the prevailing thermal regime). Littler & Kauker (1984) suggested that the crustose bases of *Corallina officinalis* are more resistant of desiccation or

heating than fronds. Severe damage was noted in *Corallina officinalis* fronds as a result of desiccation during unusually hot and sunny weather in summer 1983. An abrupt increase in temperature of 10°C caused by the hot, dry 'Santa Anna' winds (between January -and February) in Santa Cruz, California resulted in die back of several species of algae exposed at low tide (Seapy & Littler, 1982). Lüning (1990) reported that *Corallina officinalis* from Helgoland survived one week of exposure to temperatures between 0°C and 28°C.

In an exceptionally hot summer (1983, with an increase of between 4.8 and 8.5°C) Hawkins & Hartnoll (1985) observed no temperature bleaching of adult *Himanthalia elongata* (although some buttons were bleached) or other canopy forming species. However, understorey red algae showed more signs of damage with bleached *Corallina officinalis* and 'lithothamnia' observed around the edges of pools due to desiccation. Occasional damaged specimens of *Palmaria palmata*, *Osmundea pinnatifida* and *Mastocarpus stellatus* were observed.

Latham (2008) investigated the effects of temperature stress on *Corallina officinalis* through laboratory tests on samples collected in the Autumn in Devon, England from rock pools. Samples were kept at 1°C for three days and then exposed to temperatures of 5°C, 15°C, 20°C, 25°C and 30°C (the normal range of temperature experienced was suggested to be between 5 and 15°C). At 35°C, the *Corallina* was completely bleached after 3 days with a sample kept at 30°C beginning to bleach. After 7 days (the end of the experiment) the sample kept at 30°C was partially bleached. Samples kept at 5, 15, 20 and 25°C showed little change in chemicals produced in reaction to thermal stress and no bleaching suggesting the temperatures in that range had not induced stress reactions.

Most of the other species within the biotope are distributed to the north and south of Britain and Ireland and unlikely to be adversely affected by a chronic long-term temperature change. *Ulva* spp. are characteristic of upper shore rock pools, where water and air temperatures are greatly elevated on hot days. Empirical evidence for thermal tolerance to anthropogenic increases in temperature is provided by the effects of heated effluents on rocky shore communities in Maine, USA. *Ascophyllum* and *Fucus* were eliminated from a rocky shore heated to 27-30°C by a power station whilst *Ulva intestinalis* (as *Enteromorpha intestinalis*) increased significantly near the outfall (Vadas *et al.*, 1976). Barnacles, *Semibalanus balanoides*, limpets, *Patella vulgata* and littorinids also occur within this biotope. Laboratory studies suggest that adults of these species can tolerate temperature increases. The median upper lethal temperature limit in laboratory tests on *Littorina littorea*, *Littorina saxatilis* and *Semibalanus balanoides* was approximately 35°C (Davenport & Davenport, 2005). *Patella vulgata* can also tolerate high temperatures. The body temperature of *Patella vulgata* can exceed 36°C in the field, (Davies, 1970); adults become non-responsive at 37-3°C and die at temperatures of 42°C (Evans, 1948). The smaller species associated with the *Corallina officinalis* may be protected within fronds and accumulated sediments from changes in temperature although no direct evidence was found to assess the sensitivity of these to increased temperatures. Hiscock *et al.* (2004), suggest that a 1-2°C increase in temperature could increase the reproductive success of *Patella ulyssiponensis* potentially resulting in a northward expansion of the range.

Sensitivity assessment. Based on the global distribution of *Corallina officinalis* and the experiments by Latham (2008) which approximate to the pressure benchmark more than the observations of extreme events (Seapy & Littler, 1982, Hawkins & Hartnoll, 1985) it is suggested that *Corallina officinalis* would not be sensitive to either an acute or chronic increase in temperature at the pressure benchmark. Littler & Littler, (1984) suggest that the basal crustose stage is adaptive as resisters of sand scour and wave shearing as well as physiological stressors such as desiccation and

heating. Where these survive any increases in temperature above the pressure benchmark they would provide a mechanism for biotope recovery. The sensitivity of the biotope is based on the key characterizing *Corallina* turf but it should be noted that many of the associated species are considered to have 'High' resistance to changes in temperature at the pressure benchmark. It should be noted that the timing of acute increases would alter the degree of impact and hence sensitivity. An acute change occurring on the hottest day of the year and exceeding thermal tolerances would lead to mortality. The sensitivity of *Patella vulgata* and *Semibalanus balanoides* to longer-term, broad-scale perturbations would potentially be greater due to effects on reproduction but these changes may lead to species replacements (by *Patella depressa* or *Patella ulyssiponensis* and *Chthamalus* spp.) and are not considered to significantly affect the character of the biotope. An increase in *Patella ulyssiponensis* may lead to the conversion of some examples of this biotope to the sub-biotope variant characterized by this species (LR.HLR.FR.Coff.Puly).

Temperature decrease (local)

High

Q: High A: High C: High

High

Q: High A: High C: High

Not sensitive

Q: High A: High C: High

Many intertidal species are tolerant of freezing conditions as they are exposed to extremes of low air temperatures during periods of emersion. They must also be able to cope with sharp temperature fluctuations over a short period of time during the tidal cycle. In winter air temperatures are colder than the sea, conversely in summer air temperatures are much warmer than the sea. Species that occur in the intertidal are therefore generally adapted to tolerate a range of temperatures, with the width of the thermal niche positively correlated with the height of the shore (Davenport & Davenport, 2005).

Under extremely low temperatures, components of the community demonstrate tolerance. Lüning (1990) reported that *Corallina officinalis* from Helgoland survived 0°C when exposed for one week. New Zealand specimens were found to tolerate -4 °C (Frazer *et al.*, 1988). Lüning (1990) suggested that most littoral algal species were tolerant of cold and freezing. For example, the photosynthetic rate of *Chondrus crispus* recovered after 3hrs at -20°C but not after 6 hrs (Dudgeon *et al.*, 1990). The photosynthetic rate of *Mastocarpus stellatus* higher on the shore fully recovered from 24 hrs at -20°C.

The associated species are also likely to be tolerant of a decrease in temperature at the pressure benchmark. *Mytilus edulis* and *Ulva* spp. are eurytopic, found in a wide temperature range and in areas which frequently experience freezing conditions and are vulnerable to ice scour (Seed & Suchanek 1992). The tolerance of *Semibalanus balanoides* collected in the winter (and thus acclimated to lower temperatures) to low temperatures was tested in the laboratory. The median lower lethal temperature tolerance was -14.6°C (Davenport & Davenport, 2005). A decrease in temperature at the pressure benchmark is therefore unlikely to negatively affect this species. The same series of experiments indicated that median lower lethal temperature tolerances for *Littorina saxatilis* and *Littorina littorea* were -16.4 and -13°C respectively. Adults of *Patella vulgata* are also largely unaffected by short periods of extreme cold. Ekaratne & Crisp (1984) found adult limpets continuing to grow over winter when temperatures fell to -6°C, and stopped only by still more severe weather. However, loss of adhesion after exposure to -13°C has been observed with limpets falling off rocks and therefore becoming easy prey to crabs or birds (Fretter & Graham, 1994). However, in the very cold winter of 1962-3 when temperatures repeatedly fell below 0°C over a period of two months large numbers of *Patella vulgata* were found dead (Crisp, 1964). Periods of frost may also kill juvenile *Patella vulgata*, resulting in recruitment failures in some years (Bowman & Lewis, 1977). In colder conditions an active migration by mobile species may occur

down the shore to a zone where exposure time to the air (and hence time in freezing temperatures) is less. *Patella ulyssiponensis* may be sensitive to long-term decreases in temperature (Hiscock *et al.*, 2004)

Sensitivity assessment. Based on the characterizing and associated species, this biotope is considered to have 'High' resistance and 'High' resilience (by default) to this pressure and is therefore considered to be 'Not sensitive'. The timing of changes and seasonal weather could result in greater impacts on species. An acute decrease in temperature coinciding with unusually low winter temperatures may exceed thermal tolerances and lead to mortalities of the associated species although this would not alter the character of the biotope. A long-term decrease in temperature may lead to conversion of biotopes characterized by *Patella ulyssiponensis* to a similar sub-biotope.

Salinity increase (local)

Low	Medium	Medium
Q: Low A: NR C: NR	Q: High A: Low C: Medium	Q: Low A: Low C: Low

Local populations may be acclimated to the prevailing salinity regime and may, therefore, exhibit different tolerances to other populations subject to different salinity conditions and therefore caution should be used when inferring tolerances from populations in different regions. This biotope is found in full (30-35 ppt) salinity (Connor *et al.*, 2004). Biotopes found in the intertidal will naturally experience fluctuations in salinity where evaporation increases salinity and inputs of rainwater expose individuals to freshwater. Species found in the intertidal are therefore likely to have some form of behavioural or physiological adaptations to changes in salinity. The characterizing species *Corallina officinalis* and *Patella ulyssiponensis* are found in tide pools where salinities may fluctuate markedly during exposure to the air. Kinne (1971) cites maximal growth rates for *Corallina officinalis* between 33 and 38 psu in Texan lagoons.

The associated species are typically found in a range of salinities. *Ulva* species can survive hypersaline conditions in supralittoral rockpools subjected to evaporation and is considered to be a very euryhaline species, tolerant of extreme salinities ranging from 0 psu to 136 psu (Reed & Russell, 1979). In the laboratory, *Semibalanus balanoides* was found to tolerate salinities between 12 and 50 psu (Foster, 1970). Young *Littorina littorea* inhabit rock pools where salinity may increase above 35psu. Thus, the associated species may be able to tolerate some increase in salinity. *Mytilus edulis* is found in a wide range of salinities from variable salinity areas and mussels in rock pools are likely to experience hypersaline conditions on hot days. Newell (1979) recorded salinities as high as 42psu in intertidal rock pools, suggesting that *Mytilus edulis* can tolerate high salinities.

Sensitivity assessment. No direct evidence was found to assess sensitivity to this pressure. Although some increases in salinity may be tolerated by the associated species present these are generally short-term and mitigated during tidal inundation. This biotope is considered, based on the distribution of *Corallina officinalis* on the mid to lower shore to be sensitive to a persistent increase in salinity to >40 ppt. Resistance is therefore assessed as 'Low' and recovery as 'Medium' (following restoration of usual salinity). Sensitivity is therefore assessed as 'Medium'.

Salinity decrease (local)

Low	Medium	Medium
Q: Medium A: Low C: Low	Q: High A: Low C: Medium	Q: Low A: Low C: Low

Biotopes found in the intertidal will naturally experience fluctuations in salinity where evaporation increases salinity and inputs of rainwater expose individuals to freshwater. Species found in the

intertidal are therefore likely to have some form of behavioural or physiological adaptations to changes in salinity. As this biotope is present in full salinity, the assessed change at the pressure benchmark is a reduction in salinity to a variable regime (18-35 ppt) or reduced regime (18-30 ppt).

In the Baltic, *Corallina officinalis* is confined to deeper waters as surface salinity decreases (Kinne, 1971) suggesting that full salinity is required in the long-term although short-term fluctuations may be tolerated (although the thresholds of this tolerance are not clear). Kinne (1971) cites maximal growth rates for *Corallina officinalis* between 33 and 38 psu in Texan lagoons so that a decrease in salinity at the pressure benchmark would be predicted to lead to reduced growth.

Based on occurrence in estuaries, it is clear that some of the species associated with this biotope have a high tolerance for this pressure. However, it should be noted that local populations may be acclimated to the prevailing salinity regime and may, therefore, exhibit different tolerances to other populations subject to different salinity conditions so that caution should be used when inferring tolerances from populations in different regions. *Ulva* species are considered to be a very euryhaline species, tolerant of extreme salinities ranging from 0 psu to 136 psu, although some variation in salinity tolerance between populations of *Ulva intestinalis* has been found indicating that plants have some adaptation to the local salinity regime (Reed & Russell, 1979). *Littorina littorea* is found in waters of full, variable and reduced salinities (Connor *et al.*, 2004) and so populations are considered tolerant of decreases in salinity at the pressure benchmark. *Mytilus edulis* is found in a wide range of salinities from variable salinity areas (18-35ppt) such as estuaries and intertidal areas to areas of more constant salinity (Connor *et al.*, 2004). *Mytilus edulis* was recorded to grow in a dwarf form in the Baltic sea where the average salinity was 6.5psu (Riisgård *et al.*, 1993).

Prolonged reduction in salinity, e.g. from full to reduced (18-30 ppt), is likely to reduce the species richness of the biotope due to loss of some intolerant invertebrates from the assemblage associated with the *Corallina officinalis* turf.

Sensitivity assessment. Although some daily changes in salinity may be experienced these will be mitigated during tidal inundation. This biotope is considered, based on *Corallina officinalis* distribution and the evidence from Kinne, (1971), to be sensitive to a decrease in salinity at the pressure benchmark. Resistance is, therefore, assessed as '**Low**' and recovery as '**Medium**' (following restoration of usual salinity) Hence, sensitivity is assessed as '**Medium**'.

Water flow (tidal current) changes (local)

High

Q: High A: Low C: NR

High

Q: High A: High C: High

Not sensitive

Q: High A: Low C: Low

The biotope is found in a range of flow rates from 'moderately strong' (0.5-1.5 m/s) to very 'weak' (negligible) (Connor *et al.*, 2004, Dommasnes, 1969). Moderate water movement is beneficial to seaweeds as it carries a supply of nutrients and gases to the plants and removes waste products. However, if the flow becomes too strong, plants may become displaced. Additionally, an increase to stronger flows may inhibit settlement of spores and remove adults or germlings. However, *Corallina officinalis* has a compact, turf-forming growth form that reduces water flow through turbulence and friction and is probably resistant to displacement by an increase in water flow. Changes in water flow at the pressure benchmark may result in increased or decreased sediment deposition, these are not considered to alter the character of the biotope but may alter species richness of the small invertebrates associated with the turf.

Sensitivity assessment. The biotope is found across a range of flow rates, mid-range populations are considered to have 'High' resistance to a change in water flow at the pressure benchmark (although see sediment supply caveats). Resilience is assessed as 'High', by default, and the biotope is considered 'Not sensitive'.

Emergence regime changes

Low

Q: High A: Low C: Medium

Medium

Q: High A: Low C: Medium

Medium

Q: High A: Low C: Medium

Emergence regime is a key factor structuring this (and other) intertidal biotopes although it should be noted that *Corallina officinalis* may occur at a range of shore heights depending on local conditions such as the degree of wave action (Dommasnes, 1969), shore topography, run-off and degree of shelter from canopy-forming macroalgae. Increased emergence may reduce habitat suitability for characterizing and associated species through greater exposure to desiccation and reduced feeding opportunities for the barnacles, anemones, sponges and *Mytilus edulis* which feed when immersed. Changes in emergence may, therefore, lead to species replacement and the development of a biotope, more typical of the changed shore level may develop.

This biotope is considered sensitive to increased emergence as the key characterizing *Corallina officinalis* are sensitive to desiccation (Dommasnes, 1969) and are generally not found on open rock unless protected by algal canopies or where the surfaces are damp or wet. At Hinkley Point (Somerset, England), for example, seawater run-off from deep pools high in the intertidal supports dense turfs of *Corallina* spp. lower on the shore (Bamber & Irving, 1993). Fronds are highly intolerant of desiccation and do not recover from a 15 % water loss, which might occur within 40-45 minutes during a spring tide in summer (Wiedemann, 1994). Bleached corallines were observed 15 months after the 1964 Alaska earthquake which elevated areas in Prince William Sound by 10 m. Similarly, increased exposure to air caused by upward movement of 15 cm due to nuclear tests at Amchitka Island, Alaska adversely affected *Corallina pilulifera* (Johansen, 1974). During an unusually hot summer, Hawkins & Hartnoll (1985) observed damaged *Corallina officinalis* and other red algae. Littler & Littler, (1984) suggest that the basal crustose stage is adaptive, allowing individuals to survive periods of physical stress as well as physiological stress such as desiccation and heating. The basal crust stage may persist for extended periods with frond regrowth occurring when conditions are favourable.

Mobile epifauna are likely to relocate to more suitable habitats. Species such as *Patella vulgata* and *Littorina littorea* that are found throughout the intertidal zone are adapted to tolerate desiccation to some extent. For example, littorinids can seal the shell using the operculum while limpets clamped tightly to rock will reduce water loss. *Patella ulyssiponensis* is a key characterizing species for the variant biotope *coff.puly* and is generally restricted to the lower shore although it may inhabit tide pools on the upper shore (Delaney *et al.*, 1998). The green algae are also resistant to this pressure (although it may be bleached at higher shore levels during periods of high temperature) and are found throughout the intertidal including the high shore levels which may not be inundated every day. A significant, long-term, increase in emergence is therefore likely to lead to the replacement of this biotope with one more typical of the changed conditions dominated by limpets, barnacles and mussels or green algae for example.

Corallina officinalis and many of the associated species are found subtidally. Decreased emergence is likely to lead to the habitat the biotope is found in becoming more suitable for the lower shore species generally found below the biotope, leading to replacement by, for example, a kelp dominated biotope with red algae and *Corallina officinalis* surviving under the canopy.

Sensitivity assessment. Emergence is a key factor structuring the distribution of on the shore, resistance to **increased emergence** is assessed as **'Low'** as *Corallina officinalis* and associated red algae are intolerant of desiccation but basal crusts may allow individuals to persist in conditions that are unfavourable to frond development until the emergence regime is re-established. Resilience is assessed as **'Medium'** and sensitivity is therefore assessed as **'Medium'**. Pre-emption of space by *Corallina officinalis* and other red algae may reduce the establishment of lower shore species including kelps, resistance is therefore assessed as **'Medium'** to **decreased emergence** and resilience as **'Medium'** so that sensitivity is assessed as **'Medium'**. The more precautionary assessment for increased emergence is presented in the table.

Wave exposure changes (local)

High

Q: High A: Low C: NR

High

Q: High A: High C: High

Not sensitive

Q: High A: Low C: Low

This biotope is recorded from locations that are judged to range from very exposed, moderately exposed or exposed (Connor *et al.*, 2004), while Dommasnes (1969) recorded turfs from very wave sheltered areas in Norway. The degree of wave exposure influences wave height, as in more exposed areas with a longer fetch, waves would be predicted to be higher. As this biotope occurs across a range of exposures, this was therefore considered to indicate, by proxy, that biotopes in the middle of the wave exposure range would tolerate either an increase or decrease in significant wave height at the pressure benchmark. It should be noted that amounts of sediment accumulated within the turf and the associated fauna are influenced by the prevailing conditions but the biotope is still recognisable as a coralline turf.

Sensitivity assessment. The biotope is found across a range of wave exposures, mid-range populations are considered to have **'High'** resistance to a change in significant wave height at the pressure benchmark. Resilience is assessed as **'High'**, by default, and the biotope is considered **'Not sensitive'**.

Chemical Pressures

Resistance

Resilience

Sensitivity

Transition elements & organo-metal contamination

Not Assessed (NA)

Not assessed (NA)

Not assessed (NA)

Q: NR A: NR C: NR

Q: NR A: NR C: NR

Q: NR A: NR C: NR

This pressure is **Not assessed** but evidence is presented where available.

Contamination by non-synthetic chemicals, at levels greater than the pressure benchmark, may adversely impact the biotope. Little information was found concerning the effects of heavy metals on turf forming and encrusting coralline algae. However, Bryan (1984) suggested that the general order for heavy metal toxicity in seaweeds is: organic Hg > inorganic Hg > Cu > Ag > Zn > Cd > Pb. Most of the information available suggests that the associated adult gastropod molluscs are rather tolerant of heavy-metal toxicity (Bryan, 1984). Winkles may absorb metals from the surrounding water by absorption across the gills or from their diet, and evidence from experimental studies on *Littorina littorea* suggest that diet is the most important source (Bryan *et al.*, 1983). The species has been suggested as a suitable bioindicator species for some heavy metals in the marine environment. Bryan *et al.* (1983) suggested that the species is a reasonable indicator for Ag, Cd, Pb and perhaps As. In the Fal estuary *Patella vulgata* occurs at, or just outside, Restronguet Point, at

the end of the creek where metal concentrations are in the order: Zinc (Zn) 100-2000 µg/l, copper (Cu) 10-100µg/l and cadmium (Cd) 0.25-5µg/l (Bryan & Gibbs, 1983). However, in the laboratory, *Patella vulgata* was found to be intolerant of small changes in environmental concentrations of Cd and Zn by Davies (1992). At concentrations of 10µg/l, pedal mucus production and levels of activity were both reduced, indicating a physiological response to metal concentrations. Exposure to Cu at a concentration of 100 µg/l for one week resulted in progressive brachycardia (slowing of the heart beat) and the death of limpets. Zn at a concentration of 5500 µg/l produced the same effect (Marchan *et al.*, 1999).

Hydrocarbon & PAH contamination

Not Assessed (NA)

Q: NR A: NR C: NR

Not assessed (NA)

Q: NR A: NR C: NR

Not assessed (NA)

Q: NR A: NR C: NR

This pressure is **Not assessed** but evidence is presented where available.

Contamination by non-synthetic chemicals, at levels greater than the pressure benchmark, may adversely impact the biotope. Where exposed to direct contact with fresh hydrocarbons, encrusting coralline algae appear to have a high intolerance. Crump *et al.* (1999) described "dramatic and extensive bleaching" of '*Lithothamnia*' following the *Sea Empress* oil spill. Observations following the *Don Marika* oil spill (K. Hiscock, pers. comm.) were of rockpools with completely bleached coralline algae. However, Chamberlain (1996) observed that although *Lithophyllum incrustans* was affected in a short period of time by oil during the *Sea Empress* spill, recovery occurred within about a year. The oil was found to have destroyed about one third of the thallus thickness but regeneration occurred from thallus filaments below the damaged area.

Following the *Torrey Canyon* oil spill in 1967, oil and detergent dispersants affected high shore specimens of *Corallina officinalis* more than low shore specimens. Plants in deep pools were afforded some initial protection, although probably later affected by contaminated runoff. In areas of heavy spraying, however, *Corallina officinalis* was killed. (Smith 1968).

Synthetic compound contamination

Not Assessed (NA)

Q: NR A: NR C: NR

Not assessed (NA)

Q: NR A: NR C: NR

Not assessed (NA)

Q: NR A: NR C: NR

This pressure is **Not assessed** but evidence is presented where available.

Contamination by synthetic chemicals, at levels greater than the pressure benchmark, may adversely impact the biotope. Smith (1968) reported that oil and detergent dispersants from the *Torrey Canyon* spill affected high water plants of *Corallina officinalis* more than low shore plants and some plants were protected in deep pools. In areas of heavy spraying, however, *Corallina officinalis* was killed (Smith, 1968). Regrowth of fronds had begun within two months after spraying ceased (Smith, 1968). O'Brien & Dixon (1976) suggested that red algae were the most sensitive group of algae to oil or dispersant contamination, possibly due to the susceptibility of phycoerythrins to destruction. They also reported that red algae are effective indicators of detergent damage since they undergo colour changes when exposed to a relatively low concentration of detergent. However, Smith (1968) reported that red algae such as *Chondrus crispus*, *Mastocarpus stellatus* and *Laurencia pinnatifida* were amongst the algae least affected by detergents. Laboratory studies by Grandy (1984) on the effects of oil and dispersants on several red algal species concluded that they were all sensitive to oil/dispersant mixtures, with little difference between adults, sporelings, diploid or haploid life stages.

Cole *et al.* (1999) suggested that herbicides were (not surprisingly) very toxic to algae and macrophytes. Hoare & Hiscock (1974) noted that with the exception of *Phyllophora* species, all red algae including encrusting coralline forms were excluded from the vicinity of an acidified halogenated effluent discharge in Amlwch Bay, Anglesey. Intertidal populations of *Corallina officinalis* occurred in significant amounts only 600 m east of the effluent. Chamberlain (1996) observed that although *Lithophyllum incrustans* was quickly affected by oil during the *Sea Empress* spill, recovery occurred within about a year. The oil was found to have destroyed about one third of the thallus thickness but regeneration occurred from thallus filaments below the damaged area.

Most pesticides and herbicides were suggested to be very toxic for invertebrates, especially crustaceans (amphipods isopods, mysids, shrimp and crabs) and fish (Cole *et al.*, 1999).

Radionuclide contamination

No evidence (NEv)

Q: NR A: NR C: NR

Not relevant (NR)

Q: NR A: NR C: NR

No evidence (NEv)

Q: NR A: NR C: NR

No evidence was found to assess this pressure at the benchmark. Algae bioaccumulate radionuclides (with extent depending on the radionuclide and the algae species). Adverse effects have not been reported at low levels.

Introduction of other substances

Not Assessed (NA)

Q: NR A: NR C: NR

Not assessed (NA)

Q: NR A: NR C: NR

Not assessed (NA)

Q: NR A: NR C: NR

This pressure is **Not assessed**.

De-oxygenation

High

Q: Low A: NR C: NR

High

Q: High A: High C: High

Not sensitive

Q: Low A: Low C: Low

This biotope would only be exposed to low oxygen in the water column intermittently during periods of tidal immersion. In addition, in areas of wave exposure and moderately strong current flow, low oxygen levels in the water are unlikely to persist for very long as oxygen levels will be recharged by the incorporation of oxygen in the air into the water column or flushing with oxygenated waters.

No evidence was found to assess this pressure for the *Corallina* turfs. However, the associated species are unlikely to be impacted by this pressure, at the benchmark. Experiments have shown that thallus discs of *Ulva lactuca* plants can survive prolonged exposure to anoxia and hypoxia (Vermaat & Sand-Jensen, 1987; Corradi *et al.*, 2006). Following the resumption of normal oxygen conditions, gametes were produced. The associated invertebrate species also show high tolerances for reduced oxygen at levels that exceed the pressure benchmark. *Littorina littorea* can easily survive 3-6 days of anoxia (Storey *et al.*, 2013). *Semibalanus balanoides* can respire anaerobically, so they can tolerate some reduction in oxygen concentration (Newell, 1979). When placed in wet nitrogen, where oxygen stress is maximal and desiccation stress is low, *Semibalanus balanoides* have a mean survival time of 5 days (Barnes *et al.*, 1963). Limpets can also survive for a short time in anoxic seawater; Grenon & Walker, (1981) found that in oxygen-free water limpets could survive up to 36 hours, although Marshall & McQuaid (1989) found a lower tolerance for *Patella granularis*, which survived up to 11 hours in anoxic water. *Patella vulgata* and *Littorina littorea* are able to respire in the air, mitigating the effects of this pressure during the tidal cycle.

Sensitivity assessment. No direct evidence for the effects of hypoxia on *Corallina* turfs was found. As the biotope will only be exposed to this pressure when emersed and respiration will occur in air, biotope resistance was assessed as '**High**' and resilience as '**High**' (no effect to recover from), resulting in a sensitivity of '**Not sensitive**'.

Nutrient enrichment

High

Q: High A: Medium C: High

High

Q: High A: High C: High

Not sensitive

Q: High A: Medium C: High

The key characterizing *Corallina officinalis* and the associated green algae species have been identified worldwide as species that occur in areas subject to increased nutrient input within the vicinity of sewage outfalls and at intermediately polluted sites (Belgrove *et al.*, 2010; Littler & Murray, 1975; May, 1985; Brown *et al.*, 1990; Belgrove *et al.*, 1997). For example, Kindig & Littler (1980) demonstrated that *Corallina officinalis* var. *chilensis* in South California showed equivalent or enhanced health indices, highest productivity and lowest mortalities (amongst the species examined) when exposed to primary or secondary sewage effluent. Grazers in the biotope may benefit from increased availability of food resources, due to enhanced growth.

Atalah & Crowe (2010) added nutrients to rockpools occupied by a range of algae including encrusting corallines, turfs of *Mastocarpus stellatus*, *Chondrus crispus* and *Corallina officinalis* and green and red filamentous algae. The invertebrates present were mostly *Patella ulyssiponensis*, the winkle *Littorina littorea* and the flat top shell *Gibbula umbilicalis*. Nitrogen and phosphorous enhancement was via the addition of fertilisers, as either 40 g/litre or 20 g/litre. The treatments were applied for seven months and experimental conditions were maintained every two weeks. The experimental treatments do not directly relate to the pressure benchmark but indicate some general trends in sensitivity. Nutrients had no significant effect on the cover of *Corallina officinalis*. The cover of green filamentous algae was significantly increased both by reduced grazing and increased nutrients, although the effect size was synergistically magnified by the combined effect of grazer removal and nutrients. Nutrient enrichment caused an absolute increase in the average cover of green filamentous algae of 19% (± 3.9 S.E.) respect to the control treatments while the cover of red turfing algae was not affected by nutrient addition (Atalah & Crowe, 2010)

Sensitivity assessment. The pressure benchmark is relatively protective and may represent a reduced level of nutrient enrichment in previously polluted areas. Due to the tolerance of high levels of nutrient input demonstrated generally e.g. Belgrove *et al.* (2010) and Atalah & Crowe (2010), resistance to this pressure is assessed as '**High**' and resilience as '**High**' so that the biotope is assessed as '**Not sensitive**'. Where *Corallina* dominated biotopes have replaced canopy forming species in enriched areas it is not clear whether a change to the benchmark would lead to a shift in biotope type. Once established the presence of *Corallina* spp. and other turf forming species may limit recruitment by taller species (Belgrove *et al.*, 2010). No evidence was found to support an assessment of this indirect effect and it is not presented within the table.

Organic enrichment

High

Q: Low A: NR C: NR

High

Q: High A: High C: High

Not sensitive

Q: Low A: Low C: Low

No direct evidence was found to assess this pressure. Organic enrichment may lead to eutrophication with adverse environmental effects including deoxygenation, algal blooms and changes in community structure (see nutrient enrichment and de-oxygenation). Where the biotopes occur in tide-swept or wave exposed areas (Connor *et al.*, 2004) water movements will disperse organic matter reducing the level of exposure. The key characterizing

species *Corallina officinalis* has been noted to increase in abundance and may form extensive turfs within the vicinity of sewage outfalls and at intermediately polluted sites (Belgrove *et al.*, 2010; Littler & Murray, 1975; May, 1985; Brown *et al.*, 1990). As turf-forming algae *Corallina* spp. trap large amounts of sediment and are therefore not considered sensitive to sedimentation. The turfs host a variety of associated species and deposit feeders amongst these would be able to consume inputs of organic matter. Cabral-Oliveira *et al.* (2014), found higher abundances of juvenile *Patella* sp. and lower abundances of adults closer to sewage inputs, Cabral-Oliveira *et al.* (2014) suggested the structure of these populations was due to increased competition closer to the sewage outfalls.

Sensitivity assessment. Based on resistance to sedimentation, exposure to wave action, the presence of detrital consumers and the persistence of turfs in areas subject to sewage inputs resistance is assessed as '**High**' and resilience as '**High**' (by default). The biotope is therefore considered to be '**Not sensitive**' to this pressure at the benchmark.

A Physical Pressures

	Resistance	Resilience	Sensitivity
Physical loss (to land or freshwater habitat)	None	Very Low	High
	Q: High A: High C: High	Q: High A: High C: High	Q: High A: High C: High

All marine habitats and benthic species are considered to have a resistance of '**None**' to this pressure and to be unable to recover from a permanent loss of habitat (resilience is '**Very Low**'). Sensitivity within the direct spatial footprint of this pressure is, therefore '**High**'. Although no specific evidence is described confidence in this assessment is 'High', due to the incontrovertible nature of this pressure.

Physical change (to another seabed type)	None	Very Low	High
	Q: High A: High C: High	Q: High A: High C: High	Q: High A: High C: High

This biotope is characterized by the hard rock substratum to which the characterizing coralline turf and associated species such as red and green algae, barnacles limpets and anemones can firmly attach. A change to a sedimentary substratum would significantly alter the character of the biotope and would lead to the development of a biological assemblage more typical of the changed conditions. A change to an artificial substratum could also impact the development of this biotope as species may have settlement preferences for particular surface textures. Artificial hard substratum may also differ in other characteristics from natural hard substratum, so that replacement of natural surfaces with artificial may lead to changes in the biotope through changes in species composition, richness and diversity (Green *et al.*, 2012; Firth *et al.*, 2014) or the presence of non-native species (Bulleri & Airoidi, 2005). *Corallina officinalis* shows optimal settlement on finely rough artificial substrata (0.5 - 1 mm surface particle diameter). Although spores will settle and develop as crustose bases on smooth surfaces, fronds were only initiated on rough surfaces. *Corallina officinalis* settled on artificial substrata within one week in the field in summer months in New England (Harlin & Lindbergh 1977). However, in the laboratory fronds can grow from bases attached to smooth surfaces (Wiedeman pers comm) Similarly, tests with stone panels fixed to the sublittoral, mid-tide and high-tide levels of varying roughness found that *Ulva* species settle preferentially on smoother, fine-grained substratum (chalk, mottled sandstone) and *Porphyra purpurea* on rougher, granulated substratum (limestone, granite, basaltic larvae) (Luther, 1976).

Changes in substratum type can also lead to indirect effects. For example, Shanks & Wright (1986) observed that limpet mortalities were much higher at sites where the supply of loose cobbles and pebbles were greater, leading to increased abrasion through wave action 'throwing' rocks onto surfaces. Littorinids are found on a variety of shores, including sedimentary so a change in type may not significantly affect this species and some of the invertebrate species such as nematodes, amphipods and oligochaetes and polychaetes associated with sediments trapped in the *Corallina* turf are also found in sedimentary habitats

Sensitivity assessment. A change to a soft sedimentary habitat would remove the habitat for this biotope, resistance is assessed as '**None**' and resilience as '**Very low**' as the change is considered to be permanent. Sensitivity is therefore assessed as '**High**'.

Physical change (to another sediment type)

Not relevant (NR)

Q: NR A: NR C: NR

Not relevant (NR)

Q: NR A: NR C: NR

Not relevant (NR)

Q: NR A: NR C: NR

Not relevant to biotopes occurring on bedrock.

Habitat structure changes - removal of substratum (extraction)

Not relevant (NR)

Q: NR A: NR C: NR

Not relevant (NR)

Q: NR A: NR C: NR

Not relevant (NR)

Q: NR A: NR C: NR

The species characterizing this biotope are epifauna or epiflora occurring on rock and would be sensitive to the removal of the habitat. However, extraction of rock substratum is considered unlikely and this pressure is considered to be '**Not relevant**' to hard substratum habitats.

Abrasion/disturbance of the surface of the substratum or seabed

Medium

Q: High A: High C: High

High

Q: High A: Low C: Medium

Low

Q: High A: Low C: Medium

The species characterizing this biotope occur on the rock and therefore have no shelter from abrasion at the surface. Littler & Littler (1984) suggest that the basal crustose stage of *Corallina officinalis* is adapted to resist sand scour and wave shearing (as well as physiological stressors such as desiccation and heating). The base is much tougher than the fronds shown by experiments that demonstrate that the base has nearly twice the mechanical resistance (measured by penetration) of fronds (Littler & Kauker, 1984).

In general, studies show that *Corallina* and other turf-forming algae appear to be relatively resistant to single events and low levels of trampling. Brosnan & Crumrine (1994), for example, found that in experimentally trampled plots the cover of foliose and canopy forming species declined while turf-forming algae were relatively resistant. Similarly, a comparison of rocky intertidal ledges that received different amounts of visitors in Dorset, England, found that *Corallina officinalis* were present on both heavily visited and less visited ledges suggesting that this species has some resistance to trampling (Pinn & Rodgers, 2005). Povey & Keough (1991) in Mornington Peninsula, Australia investigated the effects of sustained trampling on intertidal coralline algal mats where upright branching *Corallina* spp. formed a turf with other red algae with sand and encrusting coralline algae between turfs. The experimental strips were 2 m long and 0.5 m wide. The percentage cover of upright *Corallina* spp. was significantly affected by 25 passages of a strip per day after 12 and 33 days. The algae appeared flattened and were shorter (1-2 cm high)

compared with the low intensity and control plots (3-4 cm high). However low intensity trampling within a strip (2 passages/ day) did not significantly affect the coralline turf.

Brown & Taylor (1999) also found that higher intensities of trampling damaged turfs. Moderate (50 steps per 0.09 sq. metres) or more trampling on intertidal articulated coralline algal turf in New Zealand reduced turf height by up to 50%, and weight of sand trapped within turf to about one third of controls. This resulted in declines in densities of the meiofaunal community within two days of trampling. Although the community returned to normal levels within three months of trampling events, it was suggested that the turf would take longer to recover its previous cover (Brown & Taylor 1999). Similarly, Schiel & Taylor (1999) noted that trampling had a direct detrimental effect on coralline turf species on the New Zealand rocky shore. At one site coralline bases were seen to peel from the rocks (Schiel & Taylor 1999), however, this was probably due to increased desiccation caused by loss of the algal canopy.

Species associated with the coralline turf may be more sensitive. Soft bodied species such as anemones are likely to be damaged or removed by abrasion, although anemones and sponges may repair and fragments may regrow. No evidence was found for the sensitivity of the small invertebrates associated with the coralline turf but abrasion could displace and damage these. The barnacles, limpets and littorinids that occur in low densities in this biotope, have some protection from hard shells or plates but abrasion may damage and kill individuals or detach these. All removed barnacles would be expected to die as there is no mechanism for these to reattach. Removal of limpets and barnacles may result in these being displaced to a less favourable habitat and injuries to foot muscles in limpets may prevent reattachment. Although limpets and littorinids may be able to repair shell damage, broken shells while healing will expose the individual to more risk of desiccation and predation. Evidence for the effects of abrasion is provided by a number of experimental studies on trampling (a source of abrasion) and on abrasion by wave thrown rocks and pebbles.

The effects of trampling on barnacles appear to be variable with some studies not detecting significant differences between trampled and controlled areas (Tyler-Walters & Arnold, 2008). However, this variability may be related to differences in trampling intensities and abundance of populations studied. The worst case incidence was reported by Brosnan & Crumrine (1994) who found that a trampling pressure of 250 steps in a 20x20 cm plot one day a month for a period of a year significantly reduced barnacle cover (*Semibalanus glandula* and *Chthamalus dalli*) at two study sites. Barnacle cover reduced from 66% to 7% cover in 4 months at one site and from 21% to 5% within six months at the second site. Overall barnacles were crushed and removed by trampling. Barnacle cover remained low until recruitment the following spring. Long *et al.* (2011) also found that heavy trampling (70 humans /km/hrs) led to reductions in barnacle cover. Single step experiments provide a clearer, quantitative indication of sensitivity to single events of direct abrasion. Povey & Keough (1991) in experiments on shores in Mornington Peninsula, Victoria, Australia, found that in single step experiments 10 out of 67 barnacles, (*Chthamalus antennatus* about 3 mm long), were crushed. However, on the same shore, the authors found that limpets may be relatively more resistant to abrasion from trampling. Following step and kicking experiments, few individuals of the limpet *Cellana trasomerica*, (similar size to *Patella vulgata*) suffered damage or relocated (Povey & Keough, 1991). One kicked limpet (out of 80) was broken and 2 (out of 80) limpets that were stepped on could not be relocated the following day (Povey & Keough, 1991). On the same shore, less than 5% of littorinids were crushed in single step experiments (Povey & Keough, 1991).

Shanks & Wright (1986), found that even small pebbles (<6 cm) that were thrown by wave action

in Southern California shores could create patches in aggregations of the barnacle, *Chthamalus fissus*, and could smash owl limpets (*Lottia gigantea*). Average, estimated survivorship of limpets at a wave exposed site, with many loose cobbles and pebbles allowing greater levels of abrasion was 40% lower than at a sheltered site. Severe storms were observed to lead to the almost total destruction of local populations of limpets through abrasion by large rocks and boulders. In sites with mobile cobbles and boulders increased scour results in lower densities of *Littorina* spp. compared with other, local sites with stable substratum (Carlson *et al.*, 2006).

Ulva spp. fronds are very thin and could be torn and damaged and individuals may be removed from the substratum, altering the biotope through changes in abundance and biomass. *Ulva* spp. cannot repair damage or reattach but torn fronds could still photosynthesise and produce gametes. Tearing and cutting of the fronds has been shown to stimulate gamete production and damaged plants would still be able to grow and reproduce. *Cladophora* spp. have a relatively tough thallus (Dodds & Gudder, 1992) but no direct evidence was found for resistance to abrasion. In Kimmeridge Bay in Southern England, Pinn & Rodgers (2005) found that the abundance of *Cladophora rupestris* was lower at a more heavily visited and trampled site.

Sensitivity assessment. The impact of surface abrasion will depend on the footprint, duration and magnitude of the pressure. Based on evidence from the step experiments and the relative robustness of the *Corallina officinalis* turf and associated species, resistance, to a single abrasion event is assessed as 'Medium' and recovery as 'High', so that sensitivity is assessed as 'Low'. Resistance and resilience will be lower (and hence sensitivity greater) to abrasion events that exert a greater crushing force and remove the bases than the trampling examples the assessment is based on).

Penetration or disturbance of the substratum subsurface

Not relevant (NR)

Q: NR A: NR C: NR

Not relevant (NR)

Q: NR A: NR C: NR

Not relevant (NR)

Q: NR A: NR C: NR

The species characterizing this biotope group are epifauna and epiflora occurring on rock which is resistant to subsurface penetration. The assessment for abrasion at the surface only is therefore considered to equally represent sensitivity to this pressure.

Changes in suspended solids (water clarity)

High

Q: High A: Medium C: High

High

Q: High A: High C: High

Not sensitive

Q: High A: Medium C: High

Intertidal biotopes will only be exposed to this pressure when submerged during the tidal cycle and thus have limited exposure. Siltation, which may be associated with increased suspended solids and the subsequent deposition of these is assessed separately (see siltation pressures). In general, increased suspended particles reduce light penetration and increase scour and deposition. They may enhance food supply to filter or deposit feeders (where the particles are organic in origin) or decrease feeding efficiency (where the particles are inorganic and require greater filtration efforts).

Increases in the cover of sediment trapping, turf-forming algae at the expense of canopy-forming species have been observed worldwide in temperate systems and have been linked to increased suspended solids linked to human activities worldwide (Airoldi, 2003). *Corallina* species accumulate more sediment than any other alga (Hicks, 1985). Hence an increase in suspended sediment is likely to accumulate in the coralline turf. A significant increase may result in

smothering (see above). An accumulation of sediment within the turf may attract more sediment dwelling interstitial invertebrates such as nematodes, harpacticoids and polychaetes, although in more wave exposed locations accumulation of sediment is likely to be minimal. Increased suspended sediment may also result in increased scour, which may adversely affect foliose red algae, and interfere with settling spores and recruitment if the factor is coincident with their major reproductive period. However, coralline algae, especially the crustose forms are thought to be resistant of sediment scour (Littler & Kauker, 1984), and will probably not be adversely affected at the benchmark level.

This community is unlikely to be dependent on suspended sediment. Although accumulated sediment within coralline turf habitats is likely to increase the species diversity of the epiphytic fauna, in very wave exposed locations, accumulated sediment in the habitat is likely to be minimal. A reduction in suspended sediment will probably reduce the risk of scour and reduce food availability for the few suspension feeding species in the biotope (e.g. barnacles and spirorbids present), although effects are not likely to be lethal.

The biotope occurs in shallow waters where light attenuation due to increases in turbidity is probably low. Red algae and coralline algae especially are known to be shade tolerant and are common components of the understory on seaweed dominated shores. Experiments have shown that *Ulva* is a shade tolerant genus and can compensate for reduced irradiance by increasing chlorophyll concentration and light absorption at low light levels. *Ulva* spp. were able to survive over two months in darkness and to begin photosynthesising immediately when returned to the light (Vermaat & Sand-Jensen, 1987). Limited shading from suspended sediments is therefore not considered to negatively affect this genus. Therefore, a decrease in light intensity is unlikely to adversely affect the biotope. An increase in light intensity is unlikely to adversely affect the biotope as plants can acclimate to different light levels.

Sensitivity assessment. The exposure of biotope to suspended sediments in the water column will be limited to immersion periods, and wave action will reduce accumulation. The biotope is considered to be 'Not sensitive' to a reduction in suspended solids, although this may reduce food supply to the barnacles and other filter and deposit feeders that occur in this biotope. An increase in suspended solids may lead to some sub-lethal abrasion of fronds however, evidence globally indicates that an increase in suspended solids favour the turf-forming algae that characterize this biotope (Airoldi, 2003). Resistance is therefore assessed as '**High**' and resilience as '**High**' (by default) so that the biotope is considered to be '**Not sensitive**'. An increase in suspended solids above the pressure benchmark may result in a change in species composition with an increase in species seen in very turbid, silty environments e.g. *Ahnfeltia plicata*, *Rhodothamniella floridula*, *Polyides rotunda* and *Furcellaria lumbricalis*.

Smothering and siltation rate changes (light)

High

Q: High A: High C: High

High

Q: High A: High C: High

Not sensitive

Q: High A: High C: High

Increased abundance of algal turfs worldwide has been linked to sediment perturbations although not all the pathways and mechanisms of these effects are clear (see review by Airoldi, 2003). However, even the most tolerant of organisms would eventually suffer from inhibition and mortality following smothering although the thresholds for these effects have not been identified (Airoldi, 2003).

Corallina officinalis and others within the genus (e.g. *Corallina pinnatifolia* and *Corallina vancouveriensis*) are found on shores subject to high rates of sedimentation that are periodically

disturbed by sand burial and scour (Stewart, 1989). Coralline turfs also trap sediments within the turf. The amount of sediment present and the associated fauna varies naturally depending on local conditions such as wave exposure (Dommasnes, 1969). On intertidal shores in southern California the amount of sediment trapped within turfs of *Corallina* spp. varied seasonally from < 5mm to >4.5 cm and was closely related to species composition and the structure of the turf. Airoidi (2003) identified a number of morphological, physiological and life history traits that conferred high levels of tolerance to sedimentation. Those shared by *Corallina* spp are the regeneration of upright fronds from a perennial basal crust resistant to burial and scour, calcified thalli, apical meristems, large reproductive outputs, lateral vegetative growth and slow growth rates (Airoidi, 2003). Experimental deposition of sand on coralline turfs and maintained at 3 cm or 6 cm for one month via daily top-ups did not remove the turfs but did lead to rapid (within 1 hours changes in the invertebrate species as highly mobile species moved away from the turf with later colonisation by sand adapted species (Huff & Jarett, 2007). The community had recovered one month after sand deposition ceased (Huff & Jarett, 2007).

Atalah & Crowe (2010) added sediment and nutrients to rockpools. The rock pools were occupied by a range of algae including encrusting corallines, turfs of *Mastocarpus stellatus*, *Chondrus crispus* and *Corallina officinalis* and green and red filamentous algae. The invertebrates present were mostly *Patella ulyssiponensis*, the wrinkle *Littorina littorea* and the flat top shell *Gibbula umbilicalis*. Sediment treatment involved the addition of a mixture of coarse and fine sand of either 300 mg/cm²/month or 600 mg/cm² every 15 days. The treatments were applied for seven months and experimental conditions were maintained every two weeks. The experimental treatments do not directly relate to the pressure benchmark but indicate some general trends in sensitivity. In the pools, the chronic addition of both levels of sediment led to a significant increase in the cover of *Corallina officinalis*. Sedimentation had no significant effect on the cover of green filamentous algae (*Ulva* sp.) but led to an increase in the mean cover of red turfing algae (*Mastocarpus stellatus* and *Chondrus crispus* and *Corallina officinalis*) from 11.7% (±1.0 S.E.) in controls to 26.1% (±4.7 S.E.), but there were no differences between the two levels of sedimentation. The abundance of the limpet *Patella ulyssiponensis* was significantly reduced by sedimentation. The average abundance of limpets in pools with high levels of sediment added was significantly lower ($P < 0.05$, mean 1.4 ind/144 cm² ± 0.2 S.E.) than in pools with ambient sediment loading (mean 2.7 ind/144 cm² ± 0.3 S.E.) (Atalah & Crowe, 2010).

Observations and experiments indicate that *Ulva* spp. have relatively high tolerances for the stresses induced by burial (darkness, hypoxia and exposure to sulphides) Vermaat & Sand-Jensen, 1987; Corradi et al., (2006; Kamermans et al., (1998). *Ulva lactuca* is a dominant species on sand-affected rocky shores in New Hampshire (Daly & Mathieson, 1977) although Littler et al., (1983) suggest that *Ulva* sp., are present in areas periodically subject to sand deposition not because they are able to withstand burial but because they are able to rapidly colonise sand-scoured areas. *Ulva* spp. have, however, been reported to form turfs that trap sediments (Airoidi, 2003, references therein) suggesting that resistance to chronic rather than acute siltation events may be higher.

The associated species, *Patella vulgata* and *Littorina* spp. are likely to be negatively affected by siltation (Airoidi & Hawkins, 2007; Chandrasekara & Frid, 1998; Albrecht & Reise, 1994). Experiments have shown that the addition of even thin layers of sediment (approximately 4 mm) inhibit grazing and result in loss of attachment and death after a few days Airoidi & Hawkins (2007). The laboratory experiments are supported by observations on exposed and sheltered shores with patches of sediment around Plymouth in the south west of England as *Patella vulgata* abundances were higher where deposits were absent (Airoidi & Hawkins (2007). Littler et al., (1983) found that another limpet species, *Lottia gigantea* on southern Californian shores was

restricted to refuges from sand burial on shores subject to periodic inundation by sands.

In general, propagules, early post-settlement stages and juveniles suffer severe stress and mortality from sediments (Vadas *et al.*, 1992; Airoidi, 2003). Moss *et al.*, (1973), for example, found that the growth of zygotes of *Himanthalia elongata* was inhibited by a layer of silt 1-2 mm thick and that attachment on silt was insecure.

Sensitivity assessment. The sensitivity assessment is based on the *Corallina officinalis* turf that characterizes this biotope. Resistance to siltation at the pressure benchmark is assessed as 'High', based on Airoidi (2003). Resilience is assessed as 'High' and the biotope is therefore considered to be 'Not sensitive'. The associated species within the biotope have higher sensitivities. The loss of grazing species could reduce species richness and may allow some growth of ephemeral red and green algae but this is not considered to significantly alter the biotope due to the space occupied by the coralline turf. However, the different siltation resistances of the red turf-forming algae and *Patella ulyssiponensis* (Atalah & Crowe, 2010) suggest that siltation events could alter the biotope classification between the two variants and that the variant characterized by *Patella ulyssiponensis* (LR.HLR.FR.Coff.Puly) is probably more sensitive to this pressure.

Smothering and siltation rate changes (heavy)

Low

Q: Low A: NR C: NR

Medium

Q: High A: Low C: Medium

Medium

Q: Low A: Low C: Low

No evidence was found to assess this pressure at the benchmark. A deposit at the pressure benchmark would cover all species with a thick layer of fine materials. Species associated with this biotope such as limpets and littorinids would not be able to escape and would likely suffer mortality. The tolerance of *Corallina officinalis* would be mediated by the length of time the deposit remained in place. The coralline turf and the red and green algae would be covered with sediment. Removal of the sediments by wave action and tidal currents would result in considerable scour.

Field observations and laboratory experiments have highlighted the sensitivity of limpets to sediment deposition (Airoidi & Hawkins, 2007) tested the effects of different grain sizes and deposit thickness in laboratory experiments using *Patella vulgata*. Sediments were added as a 'fine' rain to achieve deposit thicknesses of approximately 1mm, 2 mm, and 4 mm in controlled experiments and grazing and mortality observed over 8-12 days. Limpets were more sensitive to sediments with a higher fraction of fines (67% silt) than coarse (58% sand). Coarse sediments of thicknesses approximately 1, 2 and 4 mm decreased grazing activity by 35, 45 and 50 % respectively. At 1 and 2 mm thicknesses, fine sediments decreased grazing by 40 and 77 %. The addition of approximately 4 mm of fine sediment completely inhibited grazing. Limpets tried to escape the sediment but lost attachment and died after a few days (Airoidi & Hawkins, 2007). Observations on exposed and sheltered shores with patches of sediment around Plymouth in the south-west of England found that *Patella vulgata* abundances were higher where deposits were absent. The limpets were locally absent in plots with 50-65% sediment cover (Airoidi & Hawkins, 2007). Littler *et al.*, (1983) found that another limpet species, *Lottia gigantea* on southern Californian shores was restricted to refuges from sand burial on shores subject to periodic inundation by sands.

Sensitivity assessment. Sensitivity to this pressure will be mediated by site-specific hydrodynamic conditions and the footprint of the impact. Where a large area is covered sediments may be shifted by wave and tides rather than removed. However, mortality will depend on the duration of smothering, where wave action rapidly mobilises and removes fine sediments, survival may be much greater. Even small deposits of sediments are likely to result in local removal of limpets.

Resistance is assessed as '**Low**' as the impact is likely to be significant and would almost certainly result in the loss of grazers and a high proportion of the encrusting corallines and associated algae. Resilience is assessed as '**Medium**' and sensitivity is assessed as '**Medium**'.

Litter	Not Assessed (NA) Q: NR A: NR C: NR	Not assessed (NA) Q: NR A: NR C: NR	Not assessed (NA) Q: NR A: NR C: NR
---------------	--	--	--

Not assessed.

Electromagnetic changes	No evidence (NEv) Q: NR A: NR C: NR	Not relevant (NR) Q: NR A: NR C: NR	No evidence (NEv) Q: NR A: NR C: NR
--------------------------------	--	--	--

No evidence.

Underwater noise changes	Not relevant (NR) Q: NR A: NR C: NR	Not relevant (NR) Q: NR A: NR C: NR	Not relevant (NR) Q: NR A: NR C: NR
---------------------------------	--	--	--

Not relevant

Introduction of light or shading	High Q: High A: Medium C: High	High Q: High A: High C: High	Not sensitive Q: High A: Medium C: High
---	--	--	---

Corallina officinalis are shade tolerant algae, often occurring under a macroalgal canopy that reduces light penetration. In areas of higher light levels, the fronds may be lighter in colour due to bleaching (Colhart & Johansen, 1973). Other red algae in the biotope are flexible with regard to light levels. Canopy removal experiments in a rocky subtidal habitat in Nova Scotia, Canada by Schmidt & Scheibling (2007) did not find a shift in understorey macroalgal turfs (dominated by *Corallina officinalis*, *Chondrus crispus* and *Mastocarpus stellatus*) to more light-adapted species over 18 months. *Corallina officinalis* may be overgrown by epiphytes, especially during summer.

Sensitivity assessment. As the key structuring and characterizing species colonize a broad range of light environments from intertidal to deeper subtidal and shaded understorey habitats the biotope is considered to have '**High**' resistance and, by default, '**High**' resilience and, therefore, is '**Not sensitive**' to this pressure.

Barrier to species movement	High Q: Low A: NR C: NR	High Q: High A: High C: High	Not sensitive Q: Low A: Low C: Low
------------------------------------	-----------------------------------	--	--

Barriers that reduce the degree of tidal excursion may alter larval supply to suitable habitats from source populations. Conversely, the presence of barriers may enhance local population supply by preventing the loss of larvae from enclosed habitats. Barriers and changes in tidal excursion are not considered relevant to the characterizing *Corallina officinalis* as species dispersal is limited by the rapid rate of settlement and vegetative growth from bases rather than reliance on recruitment from outside of populations. Other species associated with the biotope are widely distributed and produce large numbers of larvae capable of long-distance transport and survival, resistance to this pressure is assessed as '**High**' and resilience as '**High**' by default. This biotope is therefore considered to be '**Not sensitive**'.

Death or injury by collision

Not relevant (NR)

Q: NR A: NR C: NR

Not relevant (NR)

Q: NR A: NR C: NR

Not relevant (NR)

Q: NR A: NR C: NR

Not relevant' to seabed habitats. NB. Collision by grounding vessels is addressed under 'surface abrasion.

Visual disturbance

Not relevant (NR)

Q: NR A: NR C: NR

Not relevant (NR)

Q: NR A: NR C: NR

Not relevant (NR)

Q: NR A: NR C: NR

Not relevant.

 Biological Pressures

Resistance

Resilience

Sensitivity

Genetic modification & translocation of indigenous species

No evidence (NEv)

Q: NR A: NR C: NR

Not relevant (NR)

Q: NR A: NR C: NR

No evidence (NEv)

Q: NR A: NR C: NR

The characterizing species and other common rocky shores species within the biotope, with the exception of *Mytilus edulis* which occurs in low densities, are not subject to translocation or cultivation. Commercial cultivation of *Mytilus edulis* involves the collection of juvenile mussel 'seed' or spat (newly settled juveniles ca 1-2 cm in length) from wild populations, with subsequent transportation around the UK for re-laying in suitable habitats. As the seed is harvested from wild populations from various locations the gene pool will not necessarily be decreased by translocations. Movement of mussel seed has the potential to transport pathogens and non-native species (see sensitivity assessments for *Mytilus edulis* bed biotopes). A review by Svåsand *et al.* (2007) concluded that there was a lack of evidence distinguishing between different *Mytilus edulis* populations to accurately assess the impacts of hybridisation with the congener *Mytilus galloprovincialis* and in particular how the gene flow may be affected by aquaculture. Therefore, it cannot be confirmed whether farming will have an impact on the genetics of wild individuals beyond a potential for increased hybridization.

Sensitivity assessment. No direct evidence was found regarding the potential for negative impacts of translocated mussel seed on wild *Mytilus edulis* populations. While it is possible that translocation of mussel seed could lead to gene flow between cultivated beds and local wild populations, there is currently no evidence to assess the impact (Svåsand *et al.*, 2007).

Introduction or spread of invasive non-indigenous species

High

Q: Low A: NR C: NR

High

Q: High A: High C: High

Not sensitive

Q: Low A: Low C: Low

The non-native wireweed *Sargassum muticum* may occur extensively in examples of this biotope. But the biotope persists, probably because of the small area of basal attachment of *Sargassum*.

The Australasian barnacle *Austrominius* (previously *Elminius*) *modestus* was introduced to British waters on ships during the second world war. However, its overall effect on the dynamics of rocky shores has been small as *Austrominius modestus* has simply replaced some individuals of a group of co-occurring barnacles (Raffaelli & Hawkins, 1999). Although present, monitoring indicates it

has not outnumbered native barnacles in the Isle of Cumbrae (Gallagher *et al.*, 2015) it may dominate in estuaries where it is more tolerant of lower salinities than *Semibalanus balanoides* (Gomes-Filho *et al.*, 2010). The degree of wave exposure experienced by this biotope will limit colonization by *Austrominius modestus* which tends to be present in more sheltered biotopes.

Beneath a canopy of the invasive *Codium fragile* ssp. *tomentosoides* on subtidal rocky shores in Nova Scotia, *Corallina officinalis* was the dominant species comprising 78-80% of the turf biomass, while *Chondrus crispus* and *Mastocarpus stellatus* comprised 18% (Schmidt & Scheibling, 2007). The biomass of *Corallina officinalis* was similar to those under a canopy of the native *Laminaria* species on the same shore (*Laminaria longicuris* and *Laminaria digitata*), suggesting little negative effect.

Sensitivity assessment. Overall, there is little evidence of this biotope being adversely affected by non-native species, resistance is therefore assessed as 'High', and resilience as 'High' (by default), and the biotope is considered to be 'Not sensitive'.

Introduction of microbial pathogens

High
Q: Low A: NR C: NR

High
Q: High A: High C: High

Not sensitive
Q: Low A: Low C: Low

Several coralline and non-coralline species are epiphytic on *Corallina officinalis*. Irvine & Chamberlain (1994) cite tissue destruction caused by *Titanoderma corallinae*. However, no information on pathogenic organisms in the UK was found. In Rhodophycota, viruses have been identified by means of electron microscopy (Lee, 1971) and they are probably widespread. However, nothing is known of their effects on growth or reproduction in red algae and experimental transfer from an infected to an uninfected specimen has not been achieved (Dixon & Irvine, 1977). *Corallina officinalis*, like many other algal species, has been demonstrated to produce antibacterial substances (Taskin *et al.*, 2007). Other species associated with this biotope such as littorinids, patellid limpets and other algae also experience low levels of infestation by pathogens but mass-mortalities have not been recorded. For example, parasitism by trematodes may cause sterility in *Littorina littorea*. *Littorina littorea* is also parasitized by the boring polychaete, *Polydora ciliata* and *Cliona* sp, which weakens the shell and increases crab predation. Outbreaks of the shellfish pathogen *Martelia* spp. may cause widespread mortality of *Mytilus edulis* (Mainwaring *et al.*, 2014) but populations within the UK have not been significantly impacted.

Sensitivity assessment. Based on the available evidence this biotope is considered to have 'High' resistance and hence 'High' resilience and is classed as 'Not sensitive' at the pressure benchmark.

Removal of target species

Low
Q: Low A: NR C: NR

Medium
Q: High A: Low C: Medium

Medium
Q: Low A: Low C: Low

Direct, physical impacts from harvesting are assessed through the abrasion and penetration of the seabed pressures. The sensitivity assessment for this pressure considers any biological/ecological effects resulting from the removal of target species on this biotope. The key characterizing and structuring species *Corallina officinalis* is collected for medical purposes; the fronds are dried and converted to hydroxyapatite and used as bone forming material (Ewers *et al.*, 1987). It is also sold as a powder for use in the cosmetic industry. Some species present in the biotope may also be targeted. The blue mussel *Mytilus edulis* is too small and patchy in this biotope to be targeted for commercial harvesting. However, some, unregulated recreational hand-gathering of this species and limpets, *Patella* spp., may occur. *Littorina littorea* may be targeted by commercial or recreational harvesters. Red and green algae may also be collected, *Mastocarpus stellatus*, for example, is

harvested to produce carrageen.

Littorinids are one of the most commonly harvested species of the rocky shore. Large-scale removal of *Littorina littorea* may allow a proliferation of opportunistic green algae, such as *Ulva*, on which it preferentially feeds. Experiments designed to test the effects of harvesting by removing individuals at Strangford Lough found that there was no effect of experimental treatments (either harvesting or simulated disturbance) on *Littorina littorea* abundance or body size over a 12 week period (Crossthwaite *et al.*, 2012). This suggests that these animals are generally abundant and highly mobile; thus, animals that were removed were quickly replaced by dispersal from the surrounding, un-harvested areas. However, long-term exploitation, as inferred by background levels of harvest intensity, did significantly influence population abundance and age structure (Crossthwaite *et al.*, 2012). A broadscale study of harvesting in Ireland using field studies and interviews with wholesalers and pickers did suggest that some areas were over-harvested but the lack of background data and quantitative records make this assertion difficult to test (Cummins *et al.*, 2002).

Sensitivity assessment. Collection of the key characterizing species would significantly alter the character and structure of the biotope and result in the loss of species inhabiting the turf. Collection of the associated limpet and littorinid grazers may allow red and green algae to increase in abundance and density. However, these algae may also be subject to harvesting limiting their dominance. The resistance of this biotope to targeted harvesting of characterizing and associated species is '**Low**' as the species are all relatively large, conspicuous and easily collected. Resilience is assessed '**Medium**' and sensitivity is assessed as '**Medium**'.

Removal of non-target species

Low

Q: Low A: NR C: NR

Medium

Q: High A: Low C: Medium

Medium

Q: Low A: Low C: Low

Incidental removal of the key characterizing species and associated species would alter the character of the biotope. The biotope is characterized by dense turfs of *Corallina officinalis*, these provide habitat and attachment surfaces for epiphytic species and where these trap sediments also provide a habitat for associated species. The loss of the turf due to incidental removal as by-catch would, therefore, alter the character of the habitat and result in the loss of species richness. The ecological services such as primary and secondary production provided by these species would also be lost.

Sensitivity assessment. Removal of a large percentage of the characterizing species resulting in bare rock would alter the character of the biotope, species richness and ecosystem function. Resistance is, therefore, assessed as '**Low**' and recovery as '**Medium**' so that sensitivity is assessed as '**Medium**'.

Bibliography

- Airoidi, L., 2003. The effects of sedimentation on rocky coast assemblages. *Oceanography and Marine Biology: An Annual Review*, **41**, 161-236
- Airoidi, L. & Hawkins, S.J., 2007. Negative effects of sediment deposition on grazing activity and survival of the limpet *Patella vulgata*. *Marine Ecology Progress Series*, **332**, 235-240.
- Albrecht, A. & Reise, K., 1994. Effects of *Fucus vesiculosus* covering intertidal mussel beds in the Wadden Sea. *Helgoländer Meeresuntersuchungen*, **48** (2-3), 243-256.
- Almada-Villela P.C., 1984. The effects of reduced salinity on the shell growth of small *Mytilus edulis* L. *Journal of the Marine Biological Association of the United Kingdom*, **64**, 171-182.
- Andrake, W. & Johansen, H.W., 1980. Alizarin red dye as a marker for measuring growth in *Corallina officinalis* L. (Corallinaceae, Rhodophyta). *Journal of Phycology*, **16** (4), 620-622.
- Atalah, J. & Crowe, T.P., 2010. Combined effects of nutrient enrichment, sedimentation and grazer loss on rock pool assemblages. *Journal of Experimental Marine Biology and Ecology*, **388** (1), 51-57.
- Bailey, J., Parsons, J. & Couturier, C., 1996. Salinity tolerance in the blue mussel, *Mytilus edulis*. Rep. Report no. 0840-5417, Aquaculture Association of Canada, New Brunswick, Canada
- Bamber, R.N. & Irving, P.W., 1993. The *Corallina* run-offs of Bridgewater Bay. *Porcupine Newsletter*, **5**, 190-197.
- Barnes, H., Finlayson, D.M. & Piatigorsky, J., 1963. The effect of desiccation and anaerobic conditions on the behaviour, survival and general metabolism of three common cirripedes. *Journal of Animal Ecology*, **32**, 233-252.
- Bellgrove, A., Clayton, M.N. & Quinn, G., 1997. Effects of secondarily treated sewage effluent on intertidal macroalgal recruitment processes. *Marine and Freshwater Research*, **48** (2), 137-146.
- Bellgrove, A., McKenzie, P.F., McKenzie, J.L. & Sfiligoj, B.J., 2010. Restoration of the habitat-forming fucoid alga *Hormosira banksii* at effluent-affected sites: competitive exclusion by coralline turfs. *Marine Ecology Progress Series*, **419**, 47-56.
- Blake, C. & Maggs, C.A., 2003. Comparative growth rates and internal banding periodicity of maerl species (Corallinales, Rhodophyta) from northern Europe. *Phycologia*, **42** (6), 606-612.
- Boney, A.D., 1971. Sub-lethal effects of mercury on marine algae. *Marine Pollution Bulletin*, **2**, 69-71.
- Bowman, R.S. & Lewis, J.R., 1977. Annual fluctuations in the recruitment of *Patella vulgata* L. *Journal of the Marine Biological Association of the United Kingdom*, **57**, 793-815.
- Bowman, R.S., 1981. The morphology of *Patella* spp. juveniles in Britain, and some phylogenetic inferences. *Journal of the Marine Biological Association of the United Kingdom*, **61**, 647-666.
- Brawley, S.H., 1992b. Mesoherbivores. In *Plant-animal interactions in the marine benthos* (ed. D.M John, S.J. Hawkins & J.H. Price), pp. 235-263. Oxford: Clarendon Press. [Systematics Association Special Volume, no. 46.]
- Brosnan, D.M. & Crumrine, L.L., 1994. Effects of human trampling on marine rocky shore communities. *Journal of Experimental Marine Biology and Ecology*, **177**, 79-97.
- Brown, P.J. & Taylor, R.B., 1999. Effects of trampling by humans on animals inhabiting coralline algal turf in the rocky intertidal. *Journal of Experimental Marine Biology and Ecology*, **235**, 45-53.
- Brown, V., Davies, S. & Synnot, R., 1990. Long-term monitoring of the effects of treated sewage effluent on the intertidal macroalgal community near Cape Schanck, Victoria, Australia. *Botanica Marina*, **33** (1), 85-98.
- Bryan, G.W. & Gibbs, P.E., 1983. *Heavy metals from the Fal estuary, Cornwall: a study of long-term contamination by mining waste and its effects on estuarine organisms*. Plymouth: Marine Biological Association of the United Kingdom. [Occasional Publication, no. 2.]
- Bryan, G.W., 1984. Pollution due to heavy metals and their compounds. In *Marine Ecology: A Comprehensive, Integrated Treatise on Life in the Oceans and Coastal Waters*, vol. 5. *Ocean Management*, part 3, (ed. O. Kinne), pp.1289-1431. New York: John Wiley & Sons.
- Bryan, G.W., Langston, W.J., Hummerstone, L.G., Burt, G.R. & Ho, Y.B., 1983. An assessment of the gastropod *Littorina littorea* (L.) as an indicator of heavy metal contamination in United Kingdom estuaries. *Journal of the Marine Biological Association of the United Kingdom*, **63**, 327-345.
- Bulleri, F. & Airoidi, L., 2005. Artificial marine structures facilitate the spread of a non-indigenous green alga, *Codium fragile* ssp. *tomentosoides*, in the north Adriatic Sea. *Journal of Applied Ecology*, **42** (6), 1063-1072.
- Burdin, K.S. & Bird, K.T., 1994. Heavy metal accumulation by carrageenan and agar producing algae. *Botanica Marina*, **37**, 467-470.
- Burdon, D., Dawes, O., Eades, R., Leighton, A., Musk, M. & Thompson, S. 2009. *BEEMS WP6 Intertidal Studies; Hinkley Survey-Report to Cefas*. Institute of Estuarine and Coastal Studies, University of Hull.
- Carlson, R.L., Shulman, M.J. & Ellis, J.C., 2006. Factors Contributing to Spatial Heterogeneity in the Abundance of the Common Periwinkle *Littorina Littorea* (L.). *Journal of Molluscan Studies*, **72** (2), 149-156.
- Carvalho, G.R., 1989. Microgeographic genetic differentiation and dispersal capacity in the intertidal isopod, *Jaera albifrons* Leach. In *Proceedings of the 23rd European Marine Biology Symposium, Swansea, 5-9 September 1988. Reproduction, Genetics and Distribution of Marine Organisms* (ed. J.S. Ryland & P.A. Tyler), pp. 265-271. Denmark: Olsen & Olsen.

- Chamberlain, Y.M., 1996. Lithophylloid Corallinaceae (Rhodophycota) of the genera *Lithophyllum* and *Titauserma* from southern Africa. *Phycologia*, **35**, 204-221.
- Chandrasekara, W.U. & Frid, C.L.J., 1998. A laboratory assessment of the survival and vertical movement of two epibenthic gastropod species, *Hydrobia ulvae*, (Pennant) and *Littorina littorea* (Linnaeus), after burial in sediment. *Journal of Experimental Marine Biology and Ecology*, **221**, 191-207.
- Choat, J.H. & Kingett, P.D., 1982. The influence of fish predation on the abundance cycles of an algal turf invertebrate fauna. *Oecologia*, **54**, 88-95.
- Cole, S., Codling, I.D., Parr, W. & Zabel, T., 1999. Guidelines for managing water quality impacts within UK European Marine sites. *Natura 2000 report prepared for the UK Marine SACs Project*. 441 pp., Swindon: Water Research Council on behalf of EN, SNH, CCW, JNCC, SAMS and EHS. [UK Marine SACs Project.], <http://www.ukmarinesac.org.uk/>
- Colhart, B.J., & Johanssen, H.W., 1973. Growth rates of *Corallina officinalis* (Rhodophyta) at different temperatures. *Marine Biology*, **18**, 46-49.
- Connor, D.W., Allen, J.H., Golding, N., Howell, K.L., Lieberknecht, L.M., Northen, K.O. & Reker, J.B., 2004. The Marine Habitat Classification for Britain and Ireland. Version 04.05. ISBN 1 861 07561 8. In JNCC (2015), *The Marine Habitat Classification for Britain and Ireland Version 15.03*. [2019-07-24]. Joint Nature Conservation Committee, Peterborough. Available from <https://mhc.jncc.gov.uk/>
- Connor, D.W., Brazier, D.P., Hill, T.O., & Northen, K.O., 1997b. Marine biotope classification for Britain and Ireland. Vol. 1. Littoral biotopes. *Joint Nature Conservation Committee, Peterborough, JNCC Report no. 229, Version 97.06., Joint Nature Conservation Committee, Peterborough, JNCC Report No. 230, Version 97.06.*
- Coull, B.C. & Wells, J.B.J., 1983. Refuges from fish predation: experiments with phytal meiofauna from the New Zealand rocky intertidal. *Ecology*, **64**, 1599-1609.
- Crisp, D.J. & Mwiseje, B., 1989. Diversity in intertidal communities with special reference to the *Corallina officinalis* community. *Scientia Marina*, **53**, 365-372.
- Crisp, D.J. (ed.), 1964. The effects of the severe winter of 1962-63 on marine life in Britain. *Journal of Animal Ecology*, **33**, 165-210.
- Crossthwaite, S.J., Reid, N. & Sigwart, J.D., 2012. Assessing the impact of shore-based shellfish collection on under-boulder communities in Strangford Lough. *Report prepared by the Natural Heritage Research Partnership (NHRP) between Queen's University Belfast and the Northern Ireland Environment Agency (NIEA) for the Research and Development Series No. 13/03.*
- Crump, R.G., Morley, H.S., & Williams, A.D., 1999. West Angle Bay, a case study. Littoral monitoring of permanent quadrats before and after the *Sea Empress* oil spill. *Field Studies*, **9**, 497-511.
- Cummins, V., Coughlan, S., McClean, O., Connolly, N., Mercer, J. & Burnell, G., 2002. An assessment of the potential for the sustainable development of the edible periwinkle, *Littorina littorea*, industry in Ireland. *Report by the Coastal and Marine Resources Centre, Environmental Research Institute, University College Cork.*
- Davenport, J. & Davenport, J.L., 2005. Effects of shore height, wave exposure and geographical distance on thermal niche width of intertidal fauna. *Marine Ecology Progress Series*, **292**, 41-50.
- Davies, C.E. & Moss, D., 1998. European Union Nature Information System (EUNIS) Habitat Classification. *Report to European Topic Centre on Nature Conservation from the Institute of Terrestrial Ecology, Monks Wood, Cambridgeshire*. [Final draft with further revisions to marine habitats.], Brussels: European Environment Agency.
- Davies, M.S., 1992. Heavy metals in seawater: effects on limpet pedal mucus production. *Water Research*, **26**, 1691-1693.
- Davies, S.P., 1970. Physiological ecology of *Patella* IV. Environmental and limpet body temperatures. *Journal of the Marine Biological Association of the United Kingdom*, **50** (04), 1069-1077.
- Delany, J., Myers, A.A. & McGrath, D., 1998. Recruitment, immigration and population structure of two coexisting limpet species in mid-shore tidepools, on the west coast of Ireland. *Journal of Experimental Marine Biology and Ecology*, **221**, 221-230.
- Dixon, P.S. & Irvine, L.M., 1977. *Seaweeds of the British Isles. Volume 1 Rhodophyta. Part 1 Introduction, Nemaliales, Gigartinales*. London: British Museum (Natural History) London.
- Dodds, W.K. & Gudder, D.A., 1992. The ecology of *Cladophora*. *Journal of Phycology*, **28**, 415-427.
- Dommasnes, A., 1968. Variation in the meiofauna of *Corallina officinalis* with wave exposure. *Sarsia*, **34**, 117-124.
- Dommasnes, A., 1969. On the fauna of *Corallina officinalis* L. in western Norway. *Sarsia*, **38**, 71-86.
- Dudgeon, S.R., Davison, I.R. & Vadas, R.L., 1990. Freezing tolerance in the intertidal red algae *Chondrus crispus* and *Mastocarpus stellatus*: relative importance of acclimation and adaptation. *Marine Biology*, **106**, 427-436.
- Ebere, A.G. & Akintonwa, A., 1992. Acute toxicity of pesticides to *Gobius* sp., *Palaemonetes africanus*, and *Desmocarip trispimos*. *Bulletin of Environmental Contamination and Toxicology*, **49**, 588-592.
- Ekaratne, S.U.K. & Crisp, D.J., 1984. Seasonal growth studies of intertidal gastropods from shell micro-growth band measurements, including a comparison with alternative methods. *Journal of the Marine Biological Association of the United Kingdom*, **64**, 183-210.
- Evans, R.G., 1948. The lethal temperatures of some common British littoral molluscs. *The Journal of Animal Ecology*, **17**, 165-173.
- Ewers, R., Kasperk, C. & Simmons, B., 1987. Biologisches Knochenimplantat aus Meeressalgen. *Zahnaerztliche Praxis*, **38**, 318-320.
- Firth, L., Thompson, R., Bohn, K., Abbiati, M., Airoidi, L., Bouma, T., Bozzeda, F., Ceccherelli, V., Colangelo, M. & Evans, A., 2014.

- Between a rock and a hard place: Environmental and engineering considerations when designing coastal defence structures. *Coastal Engineering*, **87**, 122-135.
- Fish, J.D. & Fish, S., 1996. *A student's guide to the seashore*. Cambridge: Cambridge University Press.
- Foster, B.A., 1970. Responses and acclimation to salinity in the adults of some balanomorph barnacles. *Philosophical Transactions of the Royal Society of London, Series B*, **256**, 377-400.
- Frazer, A.W.J., Brown, M.T. & Bannister, P., 1988. The frost resistance of some littoral and sub-littoral algae from southern New Zealand. *Botanica Marina*, **31**, 461-464.
- Fretter, V. & Graham, A., 1994. *British prosobranch molluscs: their functional anatomy and ecology*, revised and updated edition. London: The Ray Society.
- Gallagher, M.C., Davenport, J., Gregory, S., McAllen, R. & O'Riordan, R., 2015. The invasive barnacle species, *Austrominius modestus*: Its status and competition with indigenous barnacles on the Isle of Cumbrae, Scotland. *Estuarine, Coastal and Shelf Science*, **152**, 134-141.
- Gomes-Filho, J., Hawkins, S., Aquino-Souza, R. & Thompson, R., 2010. Distribution of barnacles and dominance of the introduced species *Elminius modestus* along two estuaries in South-West England. *Marine Biodiversity Records*, **3**, e58.
- Grahame, J., & Hanna, F.S., 1989. Factors affecting the distribution of the epiphytic fauna of *Corallina officinalis* (L.) on an exposed rocky shore. *Ophelia*, **30**, 113-129.
- Grandy, N., 1984. *The effects of oil and dispersants on subtidal red algae*. Ph.D. Thesis. University of Liverpool.
- Green, D., Chapman, M. & Blockley, D., 2012. Ecological consequences of the type of rock used in the construction of artificial boulder-fields. *Ecological Engineering*, **46**, 1-10.
- Grenon, J.F. & Walker, G., 1981. The tenacity of the limpet, *Patella vulgata* L.: an experimental approach. *Journal of Experimental Marine Biology and Ecology*, **54**, 277-308.
- Gressner, F. & Schramm, W., 1971. *Salinity - Plants*. In *Marine Ecology. A comprehensive, integrated treatise on life in oceans and coastal waters. Vol. 1 Environmental Factors. Part 2.* (ed. O. Kinne), pp. 705-820. London: John Wiley & Sons.
- Guiry, M.D. & Blunden, G., 1991. *Seaweed Resources in Europe: Uses and Potential*. Chichester: John Wiley & Sons.
- Guiry, M.D. & Guiry, G.M. 2015. AlgaeBase [Online], National University of Ireland, Galway [cited 30/6/2015]. Available from: <http://www.algaebase.org/>
- Hagerman, L., 1968. The ostracod fauna of *Corallina officinalis* L. in western Norway. *Sarsia*, **36**, 49-54.
- Harlin, M.M., & Lindbergh, J.M., 1977. Selection of substrata by seaweed: optimal surface relief. *Marine Biology*, **40**, 33-40.
- Hawkins, S.J. & Harkin, E., 1985. Preliminary canopy removal experiments in algal dominated communities low on the shore and in the shallow subtidal on the Isle of Man. *Botanica Marina*, **28**, 223-30.
- Hawkins, S.J. & Hartnoll, R.G., 1985. Factors determining the upper limits of intertidal canopy-forming algae. *Marine Ecology Progress Series*, **20**, 265-271.
- Hawkins, S.J., Proud, S.V., Spence, S.K. & Southward, A.J., 1994. From the individual to the community and beyond: water quality, stress indicators and key species in coastal systems. In *Water quality and stress indicators in marine and freshwater ecosystems: linking levels of organisation (individuals, populations, communities)* (ed. D.W. Sutcliffe), 35-62. Ambleside, UK: Freshwater Biological Association.
- Hicks, G.R.F. & Coull, B.C., 1983. The ecology of marine meiobenthic harpacticoid copepods. *Oceanography and Marine Biology: an Annual Review*, **21**, 67-175.
- Hicks, G.R.F., 1980. Structure of phytal harpacticoid copepod assemblages and the influence of habitat complexity and turbidity. *Journal of Experimental Marine Biology and Ecology*, **44**, 17-192.
- Hicks, G.R.F., 1985. Meiofauna associated with rocky shore algae. In *The Ecology of Rocky Coasts: essays presented to J.R. Lewis, D.Sc.*, (ed. P.G. Moore & R. Seed, ed.) pp. 36-56. London: Hodder & Stoughton Ltd.
- Hiscock, K., Southward, A., Tittley, I. & Hawkins, S., 2004. Effects of changing temperature on benthic marine life in Britain and Ireland. *Aquatic Conservation: Marine and Freshwater Ecosystems*, **14** (4), 333-362.
- Hoare, R. & Hiscock, K., 1974. An ecological survey of the rocky coast adjacent to the effluent of a bromine extraction plant. *Estuarine and Coastal Marine Science*, **2** (4), 329-348.
- Holt, T.J., Hartnoll, R.G. & Hawkins, S.J., 1997. The sensitivity and vulnerability to man-induced change of selected communities: intertidal brown algal shrubs, *Zostera* beds and *Sabellaria spinulosa* reefs. *English Nature, Peterborough, English Nature Research Report No. 234*.
- Holt, T.J., Jones, D.R., Hawkins, S.J. & Hartnoll, R.G., 1995. The sensitivity of marine communities to man induced change - a scoping report. *Countryside Council for Wales, Bangor, Contract Science Report*, no. 65.
- Holt, T.J., Rees, E.I., Hawkins, S.J. & Seed, R., 1998. Biogenic reefs (Volume IX). An overview of dynamic and sensitivity characteristics for conservation management of marine SACs. *Scottish Association for Marine Science (UK Marine SACs Project)*, 174 pp.
- Hong, J. & Reish, D.J., 1987. Acute toxicity of cadmium to eight species of marine amphipod and isopod crustaceans from southern California. *Bulletin of Environmental Contamination and Toxicology*, **39**, 884-888.
- Huff, T.M. & Jarett, J.K., 2007. Sand addition alters the invertebrate community of intertidal coralline turf. *Marine Ecology Progress*

Series, **345**, 75-82.

Irvine, L.M., 1983. *Seaweeds of the British Isles* vol. 1. *Rhodophyta Part 2A. Cryptonemiales (sensu stricto), Palmariales, Rhodymeniales*. London: British Museum (Natural History).

JNCC, 2015. The Marine Habitat Classification for Britain and Ireland Version 15.03. (20/05/2015). Available from <https://mhc.jncc.gov.uk/>

JNCC, 2015. The Marine Habitat Classification for Britain and Ireland Version 15.03. (20/05/2015). Available from <https://mhc.jncc.gov.uk/>

JNCC (Joint Nature Conservation Committee), 1999. *Marine Environment Resource Mapping And Information Database (MERMAID): Marine Nature Conservation Review Survey Database*. [on-line] <http://www.jncc.gov.uk/mermaid>

Johansen, W.H., 1974. Articulated coralline algae. *Oceanography and Marine Biology: an Annual Review*, **12**, 77-127.

Johansson, G., Eriksson, B.K., Pedersen, M. & Snoeijis, P., 1998. Long term changes of macroalgal vegetation in the Skagerrak area. *Hydrobiologia*, **385**, 121-138.

Kain, J.M., 1975a. Algal recolonization of some cleared subtidal areas. *Journal of Ecology*, **63**, 739-765.

Kennelly, S.J., 1987. Inhibition of kelp recruitment by turfing algae and consequences for an Australian kelp community. *Journal of Experimental Marine Biology and Ecology*, **112** (1), 49-60.

Kindig, A.C., & Littler, M.M., 1980. Growth and primary productivity of marine macrophytes exposed to domestic sewage effluents. *Marine Environmental Research*, **3**, 81-100.

Kinne, O. (ed.), 1971a. *Marine Ecology: A Comprehensive, Integrated Treatise on Life in Oceans and Coastal Waters. Vol. 1 Environmental Factors, Part 2*. Chichester: John Wiley & Sons.

Latham, H., 2008. Temperature stress-induced bleaching of the coralline alga *Corallina officinalis*: a role for the enzyme bromoperoxidase. *Bioscience Horizons*, 1-10

Lee, R.E., 1971. Systemic viral material in the cells of the freshwater alga *Sirodotia tenuissima* (Holden) Skuja. *Journal of Cell Science*, **8**, 623-631.

Lewis, J.R., 1964. *The Ecology of Rocky Shores*. London: English Universities Press.

Littler, M. & Murray, S., 1975. Impact of sewage on the distribution, abundance and community structure of rocky intertidal macro-organisms. *Marine Biology*, **30** (4), 277-291.

Littler, M. M., & Littler, D. S. 1984. Relationships between macroalgal functional form groups and substrata stability in a subtropical rocky-intertidal system. *Journal of Experimental Marine Biology and Ecology*, **74**(1), 13-34.

Littler, M.M., & Kauker, B.J., 1984. Heterotrichy and survival strategies in the red alga *Corallina officinalis* L. *Botanica Marina*, **27**, 37-44.

Littler, M.M., Murray, S.N. & Arnold, K.E., 1979. Seasonal variations in net photosynthetic performance and cover of intertidal macrophytes. *Aquatic Botany*, **7**, 35-46.

Long, J.D., Cochrane, E. & Dolecal, R., 2011. Previous disturbance enhances the negative effects of trampling on barnacles. *Marine Ecology Progress Series*, **437**, 165-173.

Lüning, K., 1990. *Seaweeds: their environment, biogeography, and ecophysiology*: John Wiley & Sons.

Luther, G., 1976. Bewuchsuntersuchungen auf Natursteinsubstraten im Gezeitenbereich des Nordsylter Wattenmeeres: Algen. *Helgoländer Wissenschaftliche Meeresuntersuchungen*, **28** (3-4), 318-351.

Maggs, C.A. & Hommersand, M.H., 1993. *Seaweeds of the British Isles: Volume 1 Rhodophycota Part 3A Ceramiales*. London: Natural History Museum, Her Majesty's Stationary Office.

Mainwaring, K., Tillin, H. & Tyler-Walters, H., 2014. Assessing the sensitivity of blue mussel beds to pressures associated with human activities. *Joint Nature Conservation Committee, JNCC Report No. 506.*, Peterborough, 96 pp.

Marchan, S., Davies, M.S., Fleming, S. & Jones, H.D., 1999. Effects of copper and zinc on the heart rate of the limpet *Patella vulgata* (L.) *Comparative Biochemistry and Physiology*, **123A**, 89-93.

Marshall, D.J. & McQuaid, C.D., 1989. The influence of respiratory responses on the tolerance to sand inundation of the limpets *Patella granularis* L.(Prosobranchia) and *Siphonaria capensis* Q. et G.(Pulmonata). *Journal of Experimental Marine Biology and Ecology*, **128** (3), 191-201.

Martel, A. & Chia, F.S., 1991b. Drifting and dispersal of small bivalves and gastropods with direct development. *Journal of Experimental Marine Biology and Ecology*, **150**, 131-147.

May, V., 1985. Observations on algal floras close to two sewerage outlets. *Cunninghamia*, **1**, 385-394.

McGrath, D., 1992. Recruitment and growth of the Blue-rayed limpet, *Helcion pellucidum* in south east Ireland. *Journal of Molluscan Studies*, **58**, 425-431.

Moss, B., Mercer, S., & Shearer, A., 1973. Factors Affecting the Distribution of *Himantalia elongata* (L.) S.F. Gray on the North-east Coast of England. *Estuarine and Coastal Marine Science*, **1**, 233-243.

Newell, R.C., 1979. *Biology of intertidal animals*. Faversham: Marine Ecological Surveys Ltd.

Norton, T.A., 1992. Dispersal by macroalgae. *British Phycological Journal*, **27**, 293-301.

O'Brien, P.J. & Dixon, P.S., 1976. Effects of oils and oil components on algae: a review. *British Phycological Journal*, **11**, 115-142.

- Perkol-Finkel, S. & Airoidi, L., 2010. Loss and recovery potential of marine habitats: an experimental study of factors maintaining resilience in subtidal algal forests at the Adriatic Sea. *PLoS One*, **5** (5), e10791.
- Pinn, E.H. & Rodgers, M., 2005. The influence of visitors on intertidal biodiversity. *Journal of the Marine Biological Association of the United Kingdom*, **85** (02), 263-268.
- Povey, A. & Keough, M.J., 1991. Effects of trampling on plant and animal populations on rocky shores. *Oikos*, **61**: 355-368.
- Raffaelli, D. & Hawkins, S., 1999. *Intertidal Ecology* 2nd edn.. London: Kluwer Academic Publishers.
- Reed, R.H. & Russell, G., 1979. Adaptation to salinity stress in populations of *Enteromorpha intestinalis* (L.) Link. *Estuarine and Coastal Marine Science*, **8**, 251-258.
- Riisgård, H.U., Bondo Christensen, P., Olesen, N.J., Petersen, J.K, Moller, M.M. & Anderson, P., 1993. Biological structure in a shallow cove (Kertinge Nor, Denmark) - control by benthic nutrient fluxes and suspension-feeding ascidians and jellyfish. *Ophelia*, **41**, 329-344.
- Schiel, D.R. & Foster, M.S., 1986. The structure of subtidal algal stands in temperate waters. *Oceanography and Marine Biology: an Annual Review*, **24**, 265-307.
- Schiel, D.R. & Taylor, D.I., 1999. Effects of trampling on a rocky intertidal algal assemblage in southern New Zealand. *Journal of Experimental Marine Biology and Ecology*, **235**, 213-235.
- Schmidt, A.L. & Scheibling, R.E., 2007. Effects of native and invasive macroalgal canopies on composition and abundance of mobile benthic macrofauna and turf-forming algae. *Journal of Experimental Marine Biology and Ecology*, **341** (1), 110-130.
- Seapy, R.R. & Littler, M.M., 1982. Population and Species Diversity Fluctuations in a Rocky Intertidal Community Relative to Severe Aerial Exposure and Sediment Burial. *Marine Biology*, **71**, 87-96.
- Seed, R. & Suchanek, T.H., 1992. Population and community ecology of *Mytilus*. In *The mussel Mytilus: ecology, physiology, genetics and culture*, (ed. E.M. Gosling), pp. 87-169. Amsterdam: Elsevier Science Publ. [Developments in Aquaculture and Fisheries Science, no. 25.]
- Shanks, A.L. & Wright, W.G., 1986. Adding teeth to wave action- the destructive effects of wave-bourne rocks on intertidal organisms. *Oecologia*, **69** (3), 420-428.
- Smith, J.E. (ed.), 1968. 'Torrey Canyon'. *Pollution and marine life*. Cambridge: Cambridge University Press.
- Sousa, W.P., 1979b. Experimental investigations of disturbance and ecological succession in a rocky intertidal algal community. *Ecological Monographs*, **49**, 227-254.
- Stewart, J.G., 1989. Establishment, persistence and dominance of *Corallina* (Rhodophyta) in algal turf. *Journal of Phycology*, **25** (3), 436-446.
- Storey, K.B., Lant, B., Anozie, O.O. & Storey, J.M., 2013. Metabolic mechanisms for anoxia tolerance and freezing survival in the intertidal gastropod, *Littorina littorea*. *Comparative Biochemistry and Physiology Part A: Molecular & Integrative Physiology*, **165** (4), 448-459.
- Suchanek, T.H., 1993. Oil impacts on marine invertebrate populations and communities. *American Zoologist*, **33**, 510-523.
- Svåsand, T., Crosetti, D., García-Vázquez, E. & Verspoor, E., 2007. Genetic impact of aquaculture activities on native populations. *Genimpact final scientific report (EU contract n. RICA-CT-2005-022802)*.
- Taskin, E., Ozturk, M. & Kurt, O., 2007. Antibacterial activities of some marine algae from the Aegean Sea (Turkey). *African Journal of Biotechnology*, **6** (24), 2746-2751.
- Tyler-Walters, H. & Arnold, C., 2008. Sensitivity of Intertidal Benthic Habitats to Impacts Caused by Access to Fishing Grounds. *Report to Cyngor Cefn Gwlad Cymru / Countryside Council for Wales from the Marine Life Information Network (MarLIN) [Contract no. FC 73-03-327]*, Marine Biological Association of the UK, Plymouth, 48 pp. Available from: www.marlin.ac.uk/publications
- Vadas, R.L., Johnson, S. & Norton, T.A., 1992. Recruitment and mortality of early post-settlement stages of benthic algae. *British Phycological Journal*, **27**, 331-351.
- Vadas, R.L., Keser, M. & Rusanowski, P.C., 1976. Influence of thermal loading on the ecology of intertidal algae. In *Thermal Ecology II*, (eds. G.W. Esch & R.W. McFarlane), ERDA Symposium Series (Conf-750425, NTIS), Augusta, GA, pp. 202-212.
- Vermaat J.E. & Sand-Jensen, K., 1987. Survival, metabolism and growth of *Ulva lactuca* under winter conditions: a laboratory study of bottlenecks in the life cycle. *Marine Biology*, **95** (1), 55-61.
- Watson, D.C. & Norton, T.A., 1985. Dietary preferences of the common periwinkle, *Littorina littorea* (L.). *Journal of Experimental Marine Biology and Ecology*, **88**, 193-211.
- Wiedemann, T., 1994. *Oekologische Untersuchungen in Gezeitentümpeln des Helgolaender Nord-Ost Felswatts*. , Diploma thesis, University of Kiel, Germany.
- Williams, G.A. & Seed, R., 1992. Interactions between macrofaunal epiphytes and their host algae. In *Plant-animal interactions in the marine benthos* (ed. D.M John, S.J. Hawkins & J.H. Price), pp. 189-211. Oxford: Clarendon Press. [Systematics Association Special Volume, no. 46.]