

Phylum **CTENOPHORA**Class **TENTACULATA****BOLINOPSIS INFUNDIBULUM** (O. F. Müller) [*Krumbach*, 1927, p. 15]

Abundant in May of particular years (W.G.): Sept. 1900 (A.J.S.): plentiful in spring and summer from 1916 to 1931, inside and outside Sound (M.V.L.): not seen in outside or inside tow-nettings 1946-56 (T.R.T., P.G.C.)

PLEUROBRACHIA PILEUS (O. F. Müller) [*Krumbach*, 1927, p. 6]

Always abundant towards the end of May (W.G.): adults not seen after June; minute specimens appeared in Sept. (E.J.B.): Aug. (T.V.H.): plentiful at times from spring to autumn and more rarely, in winter, from 1916 to 1931, from the coast to near Eddystone; young often in tow-nets (M.V.L.): occasionally numerous; scarce during the 4 or 5 years prior to 1956 (T.R.T., P.G.C.)

Class **NUDA****BEROË CUCUMIS** Fabricius [*Krumbach*, 1927, p. 22]

A few small specimens (E.T.B.): occasionally both large and small specimens in spring, summer and autumn from 1916 to 1931; young frequently in tow-nets; very abundant outside Breakwater to near Eddystone in summer, 1929 (M.V.L.): only very occasionally 1946-56 (T.R.T., P.G.C.)