

Characterization of Sea Surface Microlayer and Marine Aerosol Organic Composition Using STXM-NEXAFS Microscopy and FTIR Spectroscopy

Savannah L. Lewis, Lynn M. Russell,* Georges Saliba, Patricia K. Quinn, Timothy S. Bates, Craig A. Carlson, Nicholas Baetge, Lihini I. Aluwihare, Emmanuel Boss, Amanda A. Frossard, Thomas G. Bell, and Michael J. Behrenfeld



Cite This: <https://doi.org/10.1021/acsearthspacechem.2c00119>



Read Online

ACCESS |



Metrics & More




Article Recommendations



Supporting Information

ABSTRACT: Atmospheric submicron particles, generated primary marine aerosol (gPMA), seawater (depth 5 m), and sea surface microlayer (depth ~ 0.001 m) samples were collected during the North Atlantic Aerosol and Marine Ecosystem Study (NAAMES) on the R/V *Atlantis* in September 2017 and March 2018 and analyzed by scanning transmission X-ray microscopy with near-edge X-ray absorption fine structure (STXM-NEXAFS) and Fourier transform infrared (FTIR) spectroscopy. Three organic functional groups (hydroxyl, alkane, and amine) were present in all sample types, with the hydroxyl group typically being 50–90% of the quantified organic mass concentration. Microlayer and atmospheric particle samples both had a larger range of hydroxyl group to alkane group mass ratios than either seawater or gPMA. These measurements suggest that the sea surface microlayer organic composition contributes to the range of submicron atmospheric aerosol organic functional group composition. Atmospheric and microlayer sample alkane/hydroxyl group ratio variations were also associated with tracers of seawater biological activity, including chlorophyll and net primary production. Seawater and gPMA samples had relatively constant organic functional group composition for all of the sampled locations and varying phytoplankton activity conditions, suggesting that they are associated with the more consistent nature of dissolved organic carbon fraction that is ubiquitous in seawater. Eight *k*-means clusters of STXM-NEXAFS particles were identified from the spectra for all four sample types (atmospheric aerosol particles, gPMA, seawater, and microlayer), which showed that all four sample types had particles in four of the clusters and seven of the eight clusters included a mixture of sample types. These STXM-NEXAFS results support the FTIR measurements by showing consistent organic particle clusters across the four sample types as well as heterogeneity within each type.



KEYWORDS: microlayer, marine aerosol, FTIR, STXM-NEXAFS, North Atlantic Marine Aerosol Ecosystem Study, seawater

INTRODUCTION

The microlayer has been proposed to be a significant source of organic material to the marine atmosphere because it is the interface between the atmosphere and ocean,^{1,2} but there are few observations supporting this connection. The sea surface microlayer (SML) is operationally defined as the uppermost 1–1000 μm of the ocean.³ The SML is characterized by its enrichment of organic and inorganic components, including amino acids, proteins, and lipids, compared to subsurface seawater (SW). This enrichment is attributed to bubbles rising through the water column and scavenging organics that subsequently accumulate in the SML.^{1,4,5}

The SML plays a role in the enrichment of organic components of primary marine aerosols through the mechanism of bubbles bursting at the ocean surface. When the entrained bubbles reach the surface, seawater drains from the bubble film, leaving an organic-rich film that, when it bursts, ejects small particles into the atmosphere.⁶ The SML

has a heterogeneous composition, which can be composed of organic carbon, amino acids, carbohydrates, and transparent exopolymer particles (TEP).^{7,8} The chemical properties of the SML, including concentrations of TEP, free amino acids, and phosphate, have been shown to be variable, with the standard deviation of some variables being greater than the average of measured values.^{9,10}

For comparison, marine aerosol particles have been shown to be both homogeneous in terms of submicron organic functional group composition^{11,12} and heterogeneous in terms of a variety of micron-sized particle types.^{13,14} The consistency

Received: April 26, 2022

Revised: May 31, 2022

Accepted: June 1, 2022

Table 1. Concurrent Sampling Times and Station Information, Including Latitude, Longitude, Wind Speeds, Chlorophyll Concentrations, Net Primary Production (NPP) Rates, Seawater Temperatures, Number of above Detection Filters of the Four Sample Sources for FTIR, and the Number of above Detection Particles, Averaged Per Sample Source, for STXM-NEXAFS with the Number of Particles within the Average in Parenthesis

date	station	lat (deg)	long (deg)	wind speed (m s ⁻¹)	Chl-A (5 m, μg L ⁻¹)	NPP (μmol C L ⁻¹ day ⁻¹)	SW temp (°C)	FTIR ADL samples	STXM-NEXAFS ADL samples
9/10/2017	N3 S4	48	39	6.5	0.23	0.19	16.8	3	3 (13)
9/12/2017	N3 S5	51	39	5.8	0.31	0.42	14.3	4	4 (13)
9/15/2017	N3 S6	53	39	5.7	0.79	0.59	11.7	3	3 (22)
9/16/2017	N3 S6	53	39	7.3	0.83	0.72	11.7	2	2 (12)
3/27/2018	N4 S1	39	43	10.2	0.85	0.60	18.8	3	3 (18)
4/3/2018	N4 S2-RD	40	39	10.4	0.72	1.31	17.9	4	4 (17)

in organic composition and cloud condensation nuclei (CCN) activity of marine aerosols, defined as the aerosols present during clean marine conditions, has been shown across several open ocean regions of the Pacific and Atlantic Oceans.^{11,12,15,16} Aerosol particles collected from Sea Sweep, a device that generates primary aerosols by bubbling seawater from 1 m below the sea surface, have shown that generated primary marine aerosols (gPMA) results in relatively consistent CCN activity and size-resolved mass fractions.^{15,17} From these studies, it has been hypothesized that dissolved organic carbon (DOC) is the pool of carbon that has a greater influence on gPMA composition than particulate organic carbon (POC), since dissolved organic matter is the largest ocean reservoir of reduced carbon.^{18–21} This hypothesis is consistent with the fact that DOC has a relatively consistent fraction in comparison to particulate organic carbon (POC) and is present in both the SML and subsurface water.^{21–23} However, the few available measurements of organic composition for individual particles sampled from within the SML span a diverse range of types that might indicate a role for POC and the SML in influencing sea spray formation.

To understand what role the SML plays in the composition of submicron atmospheric aerosols, we collected and analyzed samples of the SML, atmospheric aerosol particles, subsurface seawater, and gPMA that were collected within 4 h of each other. There are very few studies in which SML, seawater, and atmospheric aerosol particles have been sampled simultaneously to allow assessment of how SML composition transfers to atmospheric aerosols.²² SML samples (depth ~0.001 m), subsurface seawater samples (depth 5 m), gPMA from Sea Sweep¹⁷ (depth of ~1 m), and atmospheric aerosol particles (height 18 m) collected in September 2017 and March 2018 during the North Atlantic Aerosol and Marine Ecosystem Study (NAAMES)²⁴ were analyzed using both single-particle microscopy from scanning transmission X-ray microscopy with near-edge X-ray absorption fine structure (STXM-NEXAFS) and Fourier transform infrared (FTIR). This study investigates this unique data set to assess the degree to which SML and seawater have similar organic compositions to gPMA and atmospheric aerosol particles.

METHODS

NAAMES. NAAMES was an interdisciplinary investigation conducted to improve the understanding of the earth's ocean–aerosol–cloud system.²⁴ The campaign consisted of four cruises on the R/V *Atlantis* in the North Atlantic that each included 26 days at sea in November 2015 (NAAMES 1, Winter), May–June 2016 (NAAMES 2, Late Spring),

September 2017 (NAAMES 3, Autumn), and March–April 2018 (NAAMES 4, Early Spring). The ship traveled to and from Woods Hole, MA, to ~55°N for the first three cruises. For NAAMES 4, the ship departed from San Juan, Puerto Rico, to ~44.5°N and returned to Woods Hole, MA. During each cruise, the ship occupied five to seven multiday sampling stations to conduct biological and oceanographic experiments. SML samples were only collected on NAAMES 3 and NAAMES 4, so those are the only data used in this study.

Sea Sweep. The Sea Sweep is a primary marine aerosol particle generator that has been used to provide a surrogate for particles formed from bubble bursting at the ocean surface to investigate the properties of the primary marine aerosols.¹⁷ Sea Sweep consists of a stainless steel frame attached to two inflatable pontoon floats and was deployed off the port bow of the R/V *Atlantis*. The sea surface inside the sampling hood is contained to capture aerosols produced from the microlayer, but splashing in high-wave conditions may modify the microlayer. The stainless steel hood on top of the frame maintains a laminar flow air curtain of particle-free air (charcoal and HEPA-filtered) at the bow and stern ends of the frame.¹⁷ Stainless steel frits, 200 μm in size, were positioned below the sea surface at approximately 0.75 m. The resulting particles were then transported to the instruments for analysis. Because the particles are transported directly to sampling instrumentation without time for atmospheric processes, they provide a convenient proxy for the “primary” component of sea spray aerosols.

Atmospheric Particle and gPMA Filter Collection. Atmospheric and gPMA particles were collected through a temperature- and humidity-controlled inlet during NAAMES. Atmospheric and generated primary marine aerosol (gPMA) particles were dried using diffusion driers filled with silica gel and collected on 37 mm Teflon filters (Pall Inc., 1 μm pore size) for FTIR spectroscopy analyses. <1 μm filters for atmospheric and gPMA were sampled at 8.3 L min⁻¹ after a <1 μm cyclone (SCC 2.229) at a flow rate of 16.7 L min⁻¹. Filters for <0.18 and <0.5 μm atmospheric and <1.1 μm gPMA particles were sampled at 10 L min⁻¹ after Berner impactors sampling at 30 L min⁻¹. Filters were collected for 12–23 h for atmospheric particle samples (<0.18, <0.5, and <1 μm) and 2 h for Sea Sweep samples (<0.18, <1, and <1.1 μm), resulting in filtered air volumes of 1–13.8 m³. After collection, the filters were immediately unloaded, stored in Petri dishes, and frozen at 0 °C for later analysis at Scripps Institution of Oceanography.

Seawater and Microlayer Collection. Seawater and sea surface microlayer (SML) samples were collected during the NAAMES 3 (September 2017) and NAAMES 4 (March–April

2018) cruises when on station, which is described in Behrenfeld et al.²⁴ Six SML samples were chosen for further investigation: four from NAAMES 3 (09/10/17, 09/12/17, 09/15/17, and 09/16/17) and two from NAAMES 4 (03/27/18 and 04/03/18) (Table 1). These six SML samples were chosen because they had start times within 4 h of gPMA and atmospheric particle filter start times. The SML samples were collected by a Garrett screen²⁵ (57.5 cm²), which was attached to a nylon rope and lowered over the starboard side of the ship until it touched the surface of the ocean while horizontal and then pulled back up to the deck. The screen was then held at an angle to allow the SML sample to drain off the screen for approximately 1 min, through an acid-cleaned glass funnel and into an acid-cleaned polycarbonate bottle, following the procedures used in prior studies.²⁶ This process was repeated until approximately 1.5 L was collected. Nitrile gloves were used whenever the Garrett screen was handled before and during sampling. The screen was rinsed with seawater between samples. The seawater samples were collected the same hour as the SML samples either using the ship in-line near-surface sampling system (NAAMES 3) or from 5-m CTD mounted Niskin bottles (NAAMES 4). The Imaging FlowCytobot (IFCB) cell biovolume was measured for the SML samples as well as for the in-line and Niskin seawater samples.²⁷ After SML collection, ~60 mL of seawater and SML samples were poured into amber storage bottles and immediately frozen at -20 °C to be later processed and analyzed for FTIR and STXM-NEXAFS analysis at Scripps Institution of Oceanography.

FTIR Spectroscopy and Algorithm. Samples on Teflon filters were analyzed using FTIR spectroscopy (Bruker Tensor 27 spectrometer with a deuterated triglycine sulfate (DTGS) detector) to measure the infrared transmission using 2 cm⁻¹ resolution.²⁸ The FTIR was located in a temperature- and humidity-controlled cleanroom, and the sample was purged with pure nitrogen at approximately 3 LPM for 3 min prior to the collection of each spectrum. FTIR spectra were checked for hydrate absorption, and no samples had an identifiable amount, so dehydration was not required.²⁹ The FTIR spectrum for each filter was analyzed using an automated fitting algorithm.^{28,30} When the baselining procedure resulted in degenerate spectra, defined as spectra that have no discernible peaks and low signal to noise, they were excluded from further analysis. In this data set, no filters had degenerate spectra. The FTIR spectrum for each filter was analyzed using an automated fitting algorithm.^{28,30} Five organic functional groups were quantified from these mixtures (alkane, hydroxyl, amine, carboxylic acid, and nonacidic carbonyl) and summed together to quantify the organic mass (OM) concentration, with alkene, aromatic, organosulfate, and organonitrate groups excluded because they were below detection.

OM group concentrations were considered above detection if they met three criteria: (1) the fitted peak area for the individual functional group exceeded the minimum observable peak area (defined below), (2) the fitted peak area for the individual functional group exceeded twice the standard deviation of the prescan background area, and (3) the alkane functional group was one of the groups that met the first two criteria.²⁰ OM from this technique has an uncertainty of ±20% due to functional groups that overlapped the Teflon absorption, unquantified functional groups, and semi-volatile properties.^{28,30,31} Cosine similarity (dot-product cosine on normalized spectra) was used to quantify spectral similarity

because it has been shown to be sensitive to small spectral differences in FTIR and similar chemical spectra.^{12,32}

Net Primary Production and Chlorophyll Measurements. Net primary production (NPP) ($\mu\text{mol C L}^{-1} \text{ day}^{-1}$) was estimated using the photoacclimation productivity model from Fox et al.,³³ which showed strong agreement with available measurements from 24 h ¹⁴C sodium bicarbonate uptake incubations. Shipboard chlorophyll concentrations were analyzed by a Wetlabs ACS, which measured hyperspectral particulate attenuation and absorption, which are optical measurements commonly used to estimate chlorophyll concentration, calibrated to the NAAMES campaign.³⁴ NPP was normalized to the integrated depth of the euphotic zone for each station to obtain a consistent metric for the maximum column productivity. The column maximum was used since the bubble entrainment depth is not known and the mixing layer depth was variable (6–231 m during the NAAMES campaigns).

Sample Preparation. Frozen seawater and SML samples were brought to room temperature, atomized with a constant output atomizer (TSI model 3076), dried using diffusion driers filled with silica gel, and collected on silicon nitride windows (Si₃N₄; Silson Ltd.) mounted on a rotating impactor (PIXE streaker; PIXE International, Inc.) at 1 LPM. The sampling times ranged from 3 to 5 min for STXM-NEXAFS. Teflon filters for FTIR were also atomized and dried using diffusion driers filled with silica gel at 1 LPM using the same sampling system after collection on the silicon nitride windows for STXM-NEXAFS. The sampling times ranged from 2 to 6 h for FTIR filters. The atomizer and impactor were cleaned with isopropanol between seawater and SML samples, and DI water was atomized for a minimum of 2 h between samples. Windows and filters were frozen directly after collection and stayed frozen until the time of analysis (samples were analyzed between September 2019 and June 2021). The size distribution was not measured during this process, but the particles collected after drying are expected to be smaller than 1 μm based on the design.

Atmospheric particles and gPMA were collected on Teflon filters that were used directly for nondestructive FTIR spectroscopy. After FTIR was completed, the particles on each filter were extracted and dried on silicon nitride windows to compare the composition by STXM-NEXAFS.³⁵ To extract the particles from the filters, the filters were placed in a fume hood on isopropanol-cleaned aluminum foil and 200 μL of high-performance liquid chromatography (HPLC)-grade DI water (Sigma-Aldrich) was aliquoted onto the center of the filter and agitated for 2 min. Then, 20 μL from the filter was transferred to the silicon nitride windows and left to dry overnight. The samples were frozen at 0 °C until analysis. The particles that formed after the extracts collected on these samples dried ranged from 0.2 to 7.5 μm in diameter.

Organic Functional Group Composition by STXM-NEXAFS. In all, 34 samples of extracted atmospheric particles, atomized SML, extracted gPMA, and atomized seawater from NAAMES 3 (September 2017) and NAAMES 4 (March 2018) were analyzed at the Advanced Light Source at Lawrence Berkeley National Laboratories Beamline 5.3.2.2 in a He-filled chamber using scanning transmission X-ray microscopy (STXM) with near-edge X-ray absorption fine structure (NEXAFS), following established protocols.³⁶ These 34 samples resulted in 219 stacks of spectra from particles that ranged in size from 0.2 to 7.5 μm , where these sizes reflect the




Figure 1. Normalized FTIR spectra from four sample types (atmospheric, gPMA, microlayer, and seawater) collected on six different days: 09/10/17 (red), 09/12/17 (orange), 09/15/17 (green), 09/16/17 (teal), 03/27/18 (blue), and 04/03/18 (pink). The dashed vertical lines indicate the hydroxyl group peak location at 3369 cm^{-1} (pink), alkane group peak locations at 2800 , 2852 , 2882 , and 2925 cm^{-1} (blue lines), carbonyl group peak location at 1717 cm^{-1} (teal), and amine group peak location at 1625 cm^{-1} (orange).

extracted or atomized sizes rather than their innate characteristics (and hence are not discussed here). These spectra were analyzed by an automated-shape recognition algorithm, which identified areas of high pre-edge density (non-carbon components) and high carbon density (post-edge minus pre-edge).³⁷ Of these 219 particle stacks, 122 (56%) particles passed the quality control checks, including having carbon absorbance sufficiently above the noise. The 122 particles were then clustered into eight categories using the MATLAB *k*-means clustering algorithm.

RESULTS

In this section, we report the differences in functional group composition for the different sample types (seawater, gPMA, SML, and atmospheric particle) and describe individual days on which multiple sample types were collected within four hours of each other. The morphology of the extracted (atmospheric particles and gPMA) or atomized (SML and seawater) STXM-NEXAFS particles are included (Table S1) as they contribute to chemical identification even though they are not representative of airborne samples.

Comparison of Different Sample Types. The four sample types (seawater, SML, gPMA, and atmospheric particle) show generally similar functional group composition by FTIR and STXM-NEXAFS, although indicators of small

differences in molecular structures are present (Figures 1 and 2, respectively).

FTIR (Figure 1) and STXM-NEXAFS (Figure 2) spectra for seawater were similar during the 3 days that were sampled and above the detection limit.

All samples had high cosine similarity values with low standard deviations of 0.98 ± 0.02 ($N = 3$) for FTIR and 0.97 ± 0.02 ($N = 7$) for STXM-NEXAFS (Table 2). Seawater typically had FTIR spectra with two hydroxyl group peaks, a broad alkane group peak at 2927 cm^{-1} , and a narrow amine group peak at 1615 cm^{-1} (Figure 1). The hydroxyl group fraction was 65–75% of the OM, followed by the alkane group (15–25%) and amine group (<10%). The STXM-NEXAFS spectra show an aromatic/alkene carbon peak, alkyl carbon peak, and a small carboxylic acid carbon peak, with about one in three also having potassium peaks. The carboxylic acid group was below the detection limit for FTIR, as is expected for seawater,¹² but STXM-NEXAFS is a more sensitive instrument, and carboxylic acids are a trace component of seawater composition.³⁸ STXM-NEXAFS shows four different *k*-means clusters of spectra for the seawater sample type.

Similar to previous studies,^{12,17} gPMA FTIR spectra showed a high cosine similarity value with a low standard deviation (0.94 ± 0.06 , $N = 4$) between different days and two seasons, with hydroxyl groups having the highest group mass concentration, followed by alkane and amine functional groups




Figure 2. Normalized STXM-NEXAFS centroids from four sample types (atmospheric, gPMA, microlayer, and seawater) collected on six different days: 09/10/17 (red), 09/12/17 (orange), 09/15/17 (green), 09/16/17 (teal), 03/27/18 (blue), and 04/03/18 (pink). Vertical gray lines in each panel indicate peak locations for the selected functional groups, which, from left to right, are aromatic (285 eV), ketone (286.7 eV), alkyl (287.7 eV), carboxylic acid (288.7 eV), carbonate (290.4 eV), and two potassium peaks (297.4 and 299.9 eV).

Table 2. Cosine Similarity Values with the Standard Deviation of the Four Different Sources (Atmospheric Particles (Atm), Sea Surface Microlayer (SML), Generated Primary Marine Aerosols (gPMA), and Seawater (SW)) between Both STXM and FTIR Spectra

	average STXM-NEXAFS cosine similarity				average FTIR cosine similarity			
	atm	SML	gPMA	SW	atm	SML	gPMA	SW
atm	0.95 ± 0.10				0.75 ± 0.15			
SML	0.97 ± 0.02	0.98 ± 0.02			0.73 ± 0.16	0.72 ± 0.18		
SS	0.97 ± 0.02	0.97 ± 0.02	0.97 ± 0.02		0.76 ± 0.17	0.78 ± 0.15	0.94 ± 0.06	
SW	0.97 ± 0.02	0.98 ± 0.02	0.97 ± 0.02	0.97 ± 0.02	0.83 ± 0.06	0.85 ± 0.08	0.88 ± 0.05	0.98 ± 0.02

(Figure 1). The STXM-NEXAFS spectra show carboxylic acid carbon, carbonate, and potassium peaks in four out of six of the samples and aromatic/alkene carbon peaks in half of the samples (Figure 2). The STXM-NEXAFS spectra were also similar to each other, with a cosine similarity value of 0.97 ± 0.02 ($N = 19$). There were six different k -means clusters of spectra for the gPMA sample type, showing a variety of individual particle compositions.

The SML FTIR spectra show the most variability, as demonstrated by the lowest cosine similarity value and highest standard deviation of the different sample types (0.72 ± 0.18 , $N = 6$). The different peak shapes and locations in the FTIR

spectra vary from sample to sample (Figure 1, top right panel). However, FTIR-based organic functional group composition showed that the OM in the SML samples predominately consisted of the hydroxyl group (50–80%), followed by smaller mass contributions of alkane and amine functional groups (5–35%) (Figure 3). STXM-NEXAFS had a cosine similarity of 0.98 ± 0.02 ($N = 38$). STXM-NEXAFS shows six different k -means clusters of spectra for the SML sample type, illustrating that the individual particle components atomized from the SML sample are variable.

Two of six atmospheric particle FTIR spectra and all six STXM-NEXAFS spectra had carboxylic acid groups that were




Figure 3. Organic functional group fraction from FTIR spectra with hydroxyl groups (pink), alkane groups (blue), amine groups (orange), and acid groups (green). The four sample types are atmospheric particles (atm), sea surface microlayer (SML), gPMA, and seawater (SW). The panels show the dates sampled: (A) 09/10/17, (B) 09/12/17, (C) 09/15/17, (D) 09/16/17, (E) 03/27/18, and (F) 04/03/18. The seasonal change occurs between panels (D) and (E) and is denoted with a thick black bar. The NPP values for each panel, as described in Table 1, are: (A) $0.19 \mu\text{mol C L}^{-1} \text{ day}^{-1}$, (B) $0.42 \mu\text{mol C L}^{-1} \text{ day}^{-1}$, (C) $0.59 \mu\text{mol C L}^{-1} \text{ day}^{-1}$, (D) $0.72 \mu\text{mol C L}^{-1} \text{ day}^{-1}$, (E) $0.6 \mu\text{mol C L}^{-1} \text{ day}^{-1}$, and (F) $1.31 \mu\text{mol C L}^{-1} \text{ day}^{-1}$.

likely from secondary photo-oxidative processing. The presence of carboxylic acid groups is consistent with previous observations of ambient marine aerosol samples in clean conditions.^{11,12,39} While the FTIR hydroxyl group fraction varied from 50 to 90% of the measured OM, there was no statistical difference ($p > 0.05$) when the fractions of hydroxyl, alkane, and amine groups from the different sources were compared. Atmospheric particle samples had the second-lowest FTIR cosine similarity value (0.75 ± 0.15 , $N = 6$) and the lowest STXM-NEXAFS cosine similarity value (0.95 ± 0.1 , $N = 58$). All six centroids of STXM-NEXAFS spectra had an aromatic/alkene carbon peak, a carboxylic acid carbon peak, and a carbonate peak. Four of the six STXM-NEXAFS spectra also had an alkyl carbon peak and potassium peaks. All eight of the k -means clusters were found in the atmospheric particle sample type, making it the only sample type that contains every cluster.

Comparison of Sampled Days. Six sets of samples that were each collected on the same day were used to investigate

connections between the four sample types (subsurface seawater, SML, gPMA, atmospheric aerosols) (Figure 4). This comparison was investigated to identify if there was a chemical signature that linked subsurface seawater and SML as sources for the gPMA or atmospheric particles.

The first set of samples analyzed from NAAMES 3 was on 10 September 2017 (Figure 4, row A), when there were three sample types: SML, gPMA, and atmospheric particles. The STXM-NEXAFS atmospheric particles and SML spectra both had an aromatic/alkene carbon peak, carboxylic acid carbon peak, and carbonate peaks. The gPMA spectrum also had a sharp alkyl carbon peak and carbonate peak.

Sampling on 12 September 2017 (Figure 4, row B) included all four sample types. STXM-NEXAFS showed that all four types have an aromatic/alkene carbon peak, and the gPMA, atmospheric particle, and seawater samples had a carbonate carbon peak. The FTIR showed the hydroxyl group peaks having nearly identical peak locations between 3338 and 3395




Figure 4. Centroids of normalized STXM-NEXAFS spectra from six separate days (left column) and normalized FTIR spectra (right column). Spectra were normalized to the highest absorbance peak before averaging. The rows show the dates sampled (with the number of particles within each STXM-NEXAFS spectra in parentheses): (A) 09/10/17 (atmospheric: 8, gPMA: 2, SML: 3), (B) 09/12/17 (atmospheric: 3, gPMA: 4, seawater: 3, SML: 3), (C) 09/15/17 (atmospheric: 8, gPMA: 5, SML: 9), (D) 09/16/17 (atmospheric: 7, SML: 5), (E) 03/27/18 (atmospheric: 7, seawater: 2, SML: 9), and (F) 04/03/18 (atmospheric: 8, gPMA: 3, seawater: 2, SML: 4). The colored bars on the top indicate shared functional group peak locations with aromatic/alkene absorption in black, hydroxyl in pink, alkane in blue, carbonyl in teal, and amine in orange. The STXM-NEXAFS functional groups identified by the gray lines, which, from left to right, are aromatic (285 eV), ketone (286.7 eV), alkyl (287.7 eV), carboxylic acid (288.7 eV), carbonate (290.4 eV), and two potassium peaks (297.4 and 299.9 eV). Table 1 contains the number of particles within each STXM-NEXAFS sample for each source for each day.

cm^{-1} . All four FTIR spectra had hydroxyl, alkane, and amine group peaks.

On 15 September 2017 (Figure 4, row C), three sample types were collected and analyzed (SML, gPMA, and atmospheric particle), with each corresponding STXM-NEXAFS spectrum showing a carboxylic acid carbon peak, carbonate carbon peak, and potassium peak. The atmospheric particle sample did not have an aromatic/alkene carbon peak, which was different than on 10 and 12 September 2017. The FTIR spectra from the three sample types were noticeably different from each other, with the atmospheric aerosol particle spectrum having an alkane group fraction that was twice as high as in the SML and seawater samples, as well as a large carboxylic acid group peak. The gPMA FTIR spectrum was

similar to those from 10 and 12 September 2017. The SML FTIR spectrum had a similar hydroxyl group peak shape but a larger amine group contribution than on 10 and 12 September 2017.

On 16 September 2017 (Figure 4, row D), the ship continued sampling at the same location as on 15 September 2017, and there were two types collected and analyzed: SML and atmospheric particle. The atmospheric particle sample had a STXM-NEXAFS spectrum with an alkyl carbon peak, carboxylic acid carbon peak, and carbonate peaks, as well as an aromatic/alkene carbon peak. The FTIR spectrum showed a broader hydroxyl group peak and larger alkane group peaks in the atmospheric particle sample than in the SML sample. Compared to 15 September 2017, the SML FTIR spectrum




Figure 5. Alkane-to-hydroxyl ratio of atmospheric particle (top left), SML (bottom left), gPMA (top right), and seawater (bottom right) samples compared to NPP. NPP was integrated and normalized to the depth of the euphotic zone and obtained from Baetge et al.⁴⁰ Linear fit lines are shown for $r > 0.3$. The points include measurements above detection for sample types collected on 09/10/17, 09/12/17, 09/15/17, 09/16/17, 03/27/18, and 04/03/18.

had a larger alkane group contribution and smaller hydroxyl group contribution.

The first set of samples from NAAMES 4 was analyzed on 27 March 2018 (Figure 4, row E), when there were three sample types: SML, seawater, and atmospheric particle. STXM-NEXAFS spectra showed that all three types have aromatic/alkene and carboxylic acid carbon peaks. The FTIR spectra showed that the three samples have similar hydroxyl group peak locations, with overlapping primary peaks between 3350 and 3370 cm^{-1} . The SML sample has a partial peak at 3510 cm^{-1} that is in three of the six SML samples in Figure 4. The atmospheric particle FTIR spectrum had sharp alkane group peaks, while the seawater had broad alkane group peaks, and the SML had very little alkane group absorption.

On 3 April 2018 (Figure 4, row F), all four sample types were collected and analyzed. STXM-NEXAFS spectra show a varied functional group composition, in which the seawater only had an aromatic/alkene carbon peak. The SML sample STXM-NEXAFS spectrum had an aromatic/alkene carbon peak, alkyl carbon peak, carboxylic acid carbon peak, carbonate peak, and potassium peaks. The gPMA sample has a carboxylic acid carbon peak, carbonate peak, and potassium peaks. The atmospheric aerosol particles STXM-NEXAFS spectrum had aromatic/alkene carbon, carboxylic acid carbon, and carbonate peaks. The FTIR spectra displayed two main composition types, where SML and atmospheric particle types had narrower hydroxyl group peaks that were smaller than the broad alkane group peak. gPMA and seawater spectra showed a broad hydroxyl group peak with smaller and narrower alkane group peaks.

Seawater and gPMA (<1 and $<1.1 \mu\text{m}$) samples (blue and orange in Figure 4) show FTIR spectra that do not show much variation in composition (cosine similarity of 0.94 ± 0.06 and 0.98 ± 0.02) throughout the 6 days of sampling, with two hydroxyl group peaks comprising most of the OM, followed by broad alkane group absorption and an amine group absorption. STXM-NEXAFS spectra consistently (five of six) had aromatic/alkene and alkyl carbon peaks and occasionally (three of six) had carbonate and potassium absorption peaks. Atmospheric and SML samples (red and brown in Figure 4) showed far more variability in their FTIR spectra throughout the 6 days, as demonstrated by the high standard deviation in cosine similarity (0.75 ± 0.15 and 0.72 ± 0.18). The FTIR spectra showed more variation than STXM-NEXAFS spectra, as indicated by cosine similarity values lower than 0.8 (Table 2). STXM-NEXAFS showed that the majority (11 of 12) of atmospheric particle and SML spectra had an aromatic/alkene carbon peak and carboxylic acid carbon peak. All six atmospheric particle filters also had carbonate peaks, whereas only two of six SML samples had carbonate peaks. The atmospheric particle filters were the most dissimilar of the four sample types using STXM-NEXAFS (0.95 ± 0.1).

FTIR spectra had lower cosine similarity values than NEXAFS spectra (Table 2), with hydroxyl, alkane, and amine groups being present in all samples and hydroxyl group accounting for 50–90% of the quantified OM. Seawater and gPMA samples had similar compositions with hydroxyl groups comprising 70–90% of the quantified OM, with the remainder from alkane groups (5–20%) and amine groups (2–20%), throughout the sampling days, which span two




Figure 6. Normalized *k*-means STXM-NEXAFS clusters from the NAAMES campaign (black) compared to reference STXM-NEXAFS spectra (red). The reference spectra in m, j, d, l, and g are from Takahama et al.,⁴¹ and the reference spectra from Macid and Mcarb are from Saliba et al.¹⁴ The vertical gray lines in each panel indicate peak locations for the selected functional groups, which, from left to right, are aromatic (285 eV), ketone (286.7 eV), alkyl (287.7 eV), carboxylic acid (288.7 eV), carbonate (290.4 eV), and two potassium peaks (297.4 and 299.9 eV). The pie chart indicates the number of particles from each sample type with seawater in blue, SML in brown, gPMA in orange, and atmospheric particle in red, with the number of particles within each source for each category described in Table 3.

different seasons. The SML and atmospheric particle samples had FTIR spectra that were more variable but still included the same three organic functional groups (hydroxyl, alkane, and amine), and the hydroxyl group accounted for 40–80% of quantified OM.

The variability in the SML and atmospheric particle spectra were compared using the alkane-to-hydroxyl functional group ratio and several biological proxies, including in-line chlorophyll and net primary production (NPP) (Figures 5 and S1). The alkane-to-hydroxyl group ratios had small ranges of alkane-to-hydroxyl group ratio, as evidenced by small standard deviations for both seawater (0.08 ± 0.04) and gPMA (0.26 ± 0.08). SML and atmospheric particle alkane-to-hydroxyl group ratios showed more variability, with 0.37 ± 0.42 for SML and 0.5 ± 0.37 for atmospheric particles. Here, we found weak and moderate correlations between the alkane-to-hydroxyl group ratio and both in-line chlorophyll (Figure S1) and NPP (Figure 5) for both atmospheric particle (with no significance, $p = 0.24$) and SML samples (with no significance, $p = 0.12$) over a range of alkane-to-hydroxyl ratios from 0 to 1 (0.5 ± 0.4 and 0.4 ± 0.4 , respectively). For seawater, there was no correlation, and the range of alkane-to-hydroxyl group ratios were 0.2 ± 0.1 . For gPMA, there was a strong negative correlation (with no significance, $p = 0.14$), but again low values and little variation in alkane-to-hydroxyl group ratio 0.1 ± 0.01 . While the sample size is too small to establish the statistical significance of the correlations, the difference in the

ranges of values is noteworthy, with atmospheric particle and SML ranges 2–10 times greater than seawater and gPMA ranges. The in-line chlorophyll had similar correlations to that of NPP for atmospheric particle and SML alkane-to-hydroxyl group ratios.

Clusters of STXM-NEXAFS Spectra for Sampled Particles. The average spectra of 122 particles were clustered into eight spectra categories (Figures 6 and 7). Five of the spectra categories are similar enough to previously reported categories presented by Takahama et al.⁴¹ that they are considered to be the same category (Figure 6 and Table 3). These five categories represented particles that were previously named d, g, j, l, and m and associated with combustion (g,m), freshly emitted black carbon (d), humic-like biogenic sources (g,j,m), and unidentified (l). Two other categories, Macid and Mcarb, were similar to particles identified as derived from ocean sea spray sources over the Southern Ocean, reported in Saliba et al.¹⁴

Particle spectra in cluster m had absorption in the aromatic/alkene, ketonic carbonyl, and alkyl carbon regions, similar to cluster m particles from Takahama et al.⁴¹ (cosine similarity = 0.99, two common peaks, Table 3). This cluster has particles from all four sample types (seawater, SML, gPMA, and atmospheric particle). The majority of SML particles (13 of 34) were in this cluster, which could mean that aromatic or alkene-containing substances such as phenols or humic materials are the source of the aromatic/alkene carbon




Figure 7. Normalized *k*-means cluster centroids of STXM-NEXAFS spectra from NAAMES campaign (black) and the individual particle-averaged STXM-NEXAFS spectra that comprise each cluster (gray). Reference spectra labeling is the same as Figure 6. Vertical gray lines in each panel indicate peak locations for the selected functional groups, which, from left to right, are aromatic (285 eV), ketone (286.7 eV), alkyl (287.7 eV), carboxylic acid (288.7 eV), carbonate (290.4 eV), and two potassium peaks (297.4 and 299.9 eV).

Table 3. Summary of Clustered Particle Types with the Cosine Similarity between the Literature Clusters^{14,41} and the Clusters Derived from *k*-Means Clustering in This Study^a

category	cosine similarity	common peaks	<i>N</i>	seawater	SML	gPMA	atmospheric
d	0.99	1	17	0	4	3	10
g	0.96	2	9	1	4	1	3
j	0.94	4	24	0	13	4	7
l	0.99	2	10	2	7	0	1
m	0.99	2	13	2	4	3	4
Macid	0.95	4	25	2	6	4	13
Mcarb	0.98	3	21	0	0	4	17
Csalt			3	0	0	0	3

^aThe number of functional group peaks shared between the literature clusters and the clusters of this study (common peaks), the number of particles within each cluster, and the number of particles within the cluster separated by sample type is displayed.

peak.⁴² It is also possible the aromatic/alkene carbon signal is from black carbon that had been deposited onto the sea surface or dissolved into the water column.⁴³ The particles had a high carbon content and were mostly spherical, although it is prudent to recall that the particle shape for atmospheric particle and gPMA filters represents the morphology of the dried extract rather than airborne particle morphology.

Particle spectra in cluster *j* had absorption in the aromatic/alkene carbon, carboxylic carbonyl carbon, carbonate, and potassium regions. This cluster was similar to type *j* particles identified by Takahama et al.,⁴¹ with a cosine similarity value of 0.94 and four common peaks (Table 3). This cluster had the second-highest number of spectra with 24 out of 122 particles, including 13 SML, four gPMA, and seven atmospheric

particles. Cluster *j* had the highest amount of SML particles. The morphology was a mixture of inorganic cores with organic coatings and clumps of organic-containing particles, indicating a likely marine biogenic source. Prior association of this type with humic and fulvic acids, soil substances, and biomass combustion by Takahama et al.⁴¹ suggests complex organic substances that may be surface-active⁴⁴ and have been measured previously in the SML.⁴⁵

Particle spectra in cluster *d* (cosine similarity 0.99, one common peak) have absorbance in the aromatic/alkene carbon region and small absorbance in the ketonic carbonyl carbon region. This cluster was present during both NAAMES 3 and NAAMES 4. This cluster had 17 out of 122 particles, including four SML, three gPMA, and 10 atmospheric particle samples.

The lack of peaks in this cluster other than the aromatic/alkene carbon absorption indicates it is likely fresh black carbon.⁴¹

Particle spectra in cluster Macid (cosine similarity 0.99, four common peaks, Table 3) were identified over the Eastern Pacific and over the Southern Ocean.^{14,46} This cluster is in NAAMES 3 and NAAMES 4 as well, along with carboxylic carbonyl carbon, carbonate carbon, and potassium peaks. This cluster had the highest number of spectra, with 25 particles from all four of the sample types. The morphology of atomized and extracted particles typically included low organic cuboidal shapes coated by organics, which likely represents a salt core with an organic coating.¹¹ The acidic groups in the atmospheric particle samples may indicate that secondary photochemical processes have occurred.

Particles in cluster I had a clear absorption in the aromatic/alkene carbon region and a broader absorption in the ketonic carbonyl carbon region. These particles were similar to type I identified by Takahama et al.⁴¹ (cosine similarity = 0.99, two common peaks, Table 3), which was classified as not having an identified source because of the limited number of particles found and the limited availability of tracers for some types of sources. This cluster may be indicative of biological remnants such as cell fragments and lipid-like materials (sometimes identified as TEP) that have been measured previously in both SML and nascent sea spray aerosols.^{4,7} This particle cluster had several different morphologies that included cuboidal, amorphous, and spherical shapes after atomization, making it different from previously reported BC morphology.⁴¹ This cluster was present during both NAAMES 3 and NAAMES 4. This cluster had 10 out of 122 particles, including seven SML, one atmospheric, and two seawater particles.

Particles in cluster Mcarb accounted for 21 out of 122 particles and mostly consisted of extracted atmospheric particles, which are 17 out of the 21 particles. This cluster was present during both NAAMES 3 and NAAMES 4. This cluster was first observed in Saliba et al.¹⁴ The cluster spectra include three peaks, namely alkyl carbon, carbonate, and potassium, and the spectra in the cluster have a cosine similarity of 0.98. This cluster is also similar to the calcareous phytoplankton cluster found in Hawkins and Russell.¹³ However, the morphology of the atmospheric particles from this work often had salt cores and diverse morphologies, whereas the calcareous phytoplankton type from Hawkins and Russell had structures that resembled cell or shell parts without salt cores. This particle type was likely sea spray particles, but it included only particles that were extracted from filters (atmospheric particle and gPMA) for which the extracted morphology may be different from their airborne state.

Particles in cluster g included all four sample types and contained nine out of 122 particles from both campaigns. This particle type was previously identified⁴¹ (cosine similarity = 0.96, two shared peaks, Table 3), and the particle cluster was identified as combustion-related due to the prominent aromatic/alkene carbon peak and amorphous carboxylic carbonyl carbon absorption. The high carbon content and the lack of potassium or alkyl groups in this cluster make it likely that the cluster g particles in NAAMES 3 and NAAMES 4 are from combustion sources.

Particle spectra in cluster crystallized salt (Csalt) only accounted for three out of 122 particles and were from a single extracted atmospheric particle sample from NAAMES 3. This cluster is distinguished by the peak splitting at the two potassium peaks (297.4 and 299.9 eV). This may be due to the

extracted particles crystallizing onto the silicon nitride windows, which has been shown to cause ligand field splitting.⁴⁷

DISCUSSION

Prior work has indicated the SML to be enriched in amino acids, lipids, and organics in comparison to subsurface seawater,^{1,48,49} suggesting that there would be a distinct difference in the organic composition between the SML and the other sample types (seawater, gPMA, and atmospheric particles). SML and atmospheric particle samples are different from the seawater and gPMA samples, as shown by their lower cosine similarity values for FTIR and higher standard deviations between cosine similarity values of individual FTIR spectra within the same sample type (Table 2 and Figures 1 and 2). However, we see many similarities in composition between the four types of samples with both FTIR and STXM-NEXAFS. All have three organic functional groups (hydroxyl, alkane, and amine), with hydroxyl consistently being the largest fraction (Figures 1 and 3). Most particles sampled have an aromatic/alkene carbon peak in STXM-NEXAFS spectra, regardless of sample source (Figure 2). Both FTIR and STXM-NEXAFS show hydroxyl groups and alkane groups; alkene groups are present in the STXM-NEXAFS spectra for the four sample types, although alkene groups are below the detection limit for FTIR quantification (Figure 4). Seawater and gPMA samples had very consistent FTIR spectra across the sampled days, with high cosine similarity values for spectra from the same sample type (Table 2). When the STXM-NEXAFS particles were separated into categories by *k*-means clustering, the only cluster that had particles from a single sample type was Csalt, which had only three atmospheric aerosol particles. Other than this new category (which may be associated with a salt artifact), there were no unique particle clusters that were from only one sample source, rather the particle clusters had particles from most of the sample sources.

While the FTIR organic functional group composition was similar on average for the four sample types, the FTIR spectra showed peaks at different locations and with different shapes, particularly in the alkane region, for SML and atmospheric particle samples. The variability of the FTIR composition of both atmospheric particle and SML samples is evident in the low cosine similarity values (<0.8) and high standard deviations among samples within the same type (Table 2). In contrast, seawater and gPMA samples had much more consistent FTIR spectra, shown by the high cosine similarity values (>0.93) and low standard deviations (Table 2). This range of different organic composition of SML and atmospheric aerosol samples are consistent with prior measurements showing the presence of varying amounts of polymers, proteinaceous matter, and TEP in both SML and marine aerosols.^{4,50} Longer chain saccharides have been observed to be enriched in atmospheric aerosol particles and SML that are present but not enriched in seawater.⁵¹ The SML has been shown to have a highly heterogeneous and variable composition,^{8,9,52} whereas seawater often has less variability.⁵² The IFCB cell biovolume was also more concentrated and more variable in the SML compared to the in-line seawater measurements ($p < 0.05$) (Figures S2 and S3). The individual particles measured by STXM-NEXAFS are also consistent with the heterogeneity of SML, with particles having a range of structures from salt cores with organic coatings to solid organic to mixtures of the two.

In Figure 3, panels C, D, and F show sampling times that have higher NPP than in panels A, B, E (Table 1). When NPP was higher, both SML and atmospheric aerosol particle samples had higher alkane/hydroxyl group ratios with high standard deviations (0.37 ± 0.42 and 0.49 ± 0.37 , respectively), whereas seawater and gPMA samples were nearly identical with low standard deviations (0.25 ± 0.08 and 0.08 ± 0.04 , respectively) despite the observed differences in NPP and Chl. The correlations of the four-sample alkane/hydroxyl group ratio to both chlorophyll (Figures S1 and S4) and NPP (Figure 5) indicate a possible role for the SML in contributing to the variability in atmospheric aerosol particle composition, perhaps driving the atmospheric particle correlation to biological tracers, as illustrated in Figure 8.




Figure 8. Diagram illustrating the influence of seawater (bottom left) on gPMA (top left with a trapezoid representing Sea Sweep) aerosol particle composition and the influence of SML (bottom right) on atmospheric particle composition (top right). The arrows are based on the similarity of the composition shown in the pie charts and the spectra and do not indicate that they are the only source as seawater and SML components can be mixed. The individual FTIR spectra shown are during a lower chlorophyll period (blue, 12 September 2017) and a higher chlorophyll period (green, 3 April 2018). The pie graphs, outlined in corresponding colors for low vs high chlorophyll, demonstrate the organic functional group FTIR composition with hydroxyl groups (pink), alkane groups (blue), and amine groups (orange).

The gPMA and seawater alkane-to-hydroxyl group ratios are consistently unrelated to NPP, showing almost no change for the 6 days that were sampled (0.1 ± 0.01 (gPMA) and 0.2 ± 0.1 (seawater) alkane/hydroxyl group ratio). This could be due to gPMA and seawater organic composition being largely influenced by DOC, which has a relatively consistent fraction in comparison to POC.^{12,16,18} This smaller seawater range may contribute to the similarly small range in gPMA, suggesting that for the NAAMES open ocean conditions, gPMA particles may have more contributions from seawater (and less from SML) than do the atmospheric particles. In contrast, atmospheric particles and SML samples show much wider variability, both spanning the range of 0.1–1 for alkane/

hydroxyl group ratio and showing weak positive correlations with both chlorophyll and NPP. STXM-NEXAFS showed multiple particle clusters for all sample types; however, the low number of particles sampled with STXM-NEXAFS means that we cannot rule out more similarity in seawater and gPMA than in SML and atmospheric aerosol particles.

Phytoplankton blooms are associated with an increase in surface-active compounds, particularly lipids, with increasing POC concentrations.⁵³ Recent contributions from recently produced DOC⁴⁰ may explain the changing composition of the SML during these times. Atmospheric aerosol particles could also be influenced by secondary atmospheric processes of organic components, but there is no evidence of such processes changing the alkane and hydroxyl group fractions.¹² The lack of expected changes to the alkane/hydroxyl group ratio from photochemical processes means that the ratio could serve as an approximately conserved tracer for the composition between these reservoirs. If this ratio is conserved on transfer between the ocean and the atmosphere, then the 0.1–1 range of alkane/hydroxyl group ratio in the atmospheric particles and the SML would not be explained by the seawater source range of 0–0.3.

Similar positive correlations of alkane and hydroxyl group contributions to biological activity have also been found in gPMA aerosol particles in more productive waters, where gPMA samples had higher alkane group fractions (and lower hydroxyl group fractions) than nonproductive waters during the Western Atlantic Climate Study (WACS) and California Nexus (CalNex) projects.¹² This difference from our results could be explained by the larger range of chlorophyll concentrations or the lower range of wind speeds (4 ± 2 m s⁻¹ for WACS and 5 ± 2 m s⁻¹ for CalNex) in the WACS/CalNEX data sets than in NAAMES 3 and NAAMES 4. In fact, the WACS/CalNEX measurements of hydroxyl and alkane groups also did not have a clear trend for chlorophyll <1 (which is the range of all NAAMES 3 and NAAMES 4 measurements); it was only at larger concentrations that the gPMA composition showed a moderate correlation to biological proxies. The contributions of SML and seawater to gPMA aerosol particles may vary with ocean conditions such as wind speed,^{49,54} since very little is known about the prevalence and variability of the SML or its effects on gPMA sampling.

CONCLUSIONS

Our results show the organic functional groups in atmospheric particles have chemical similarities to seawater, gPMA, and SML in the North Atlantic. The four sample types were analyzed using single-particle microscopy by STXM-NEXAFS and FTIR spectroscopy. Both measurements showed the presence of hydroxyl, alkyl, and alkene functional groups, and FTIR hydroxyl groups were consistently 50–90% of the quantified OM.

The larger range of alkane/hydroxyl group ratios, the greater FTIR spectral differences (low cosine similarity values and high standard deviations of cosine similarity values for individual spectra within the same sample type), and the weak correlations to tracers of biological activity for both atmospheric particle and SML samples suggest a biologically influenced source contributing to a wider range of composition for SML and atmospheric particles than was found for seawater or gPMA. Both atmospheric particle and SML samples showed an increased alkane/hydroxyl group ratio (associated with decreases in hydroxyl group fraction and increases in alkane group fraction) during periods of higher biological activity

tracers, which suggests that the SML was an important contributor to the organic composition of atmospheric aerosol particles. Seawater and gPMA samples had a more consistent composition, showing little variation for the range of biological activity that was observed and supporting previous findings that seawater contributes a relatively consistent organic functional group composition to atmospheric aerosol particles.

Of the eight STXM-NEXAFS clusters that were identified, there were no clusters that contained only SML particle types and only one small cluster, Csalt, that was only found in a single atmospheric particle sample. This variety of particle compositions is consistent with other observations in open ocean conditions around the world,^{11,13,14} illustrating the heterogeneity of components within the SML.

The SML showed more variability in FTIR spectra among the six case-study days than either seawater or gPMA, with an FTIR cosine similarity value of 0.72 ± 0.18 for SML compared to FTIR cosine similarity values of 0.98 ± 0.02 for seawater and 0.94 ± 0.06 for gPMA. This heterogeneity of composition could explain why some reports show a robust link between marine aerosol composition and SML and others do not,^{1,2,10,22,23} particularly, given the limited sample number of the few available studies of microlayer and aerosol composition.

The variability of the SML during sampled days displays the need for a longer time series in which day-to-day variability can be teased apart from various environmental factors, including wind speed and biological activity. Sampling the atmospheric particle and gPMA samples directly onto substrates for STXM-NEXAFS analysis would have allowed for the characterization of airborne-relevant particle morphology and diameter, and additional sampling would have provided a more robust data set for understanding seasonal and regional differences.

■ ASSOCIATED CONTENT

SI Supporting Information

The Supporting Information is available free of charge at <https://pubs.acs.org/doi/10.1021/acsearthspacechem.2c00119>.

Additional details; detailed information about every particle that was above the detection limit for STXM-NEXAFS (Table S1); alkane-to-hydroxyl ratio of concurrently sampled ambient, SML, Sea Sweep, and seawater filters compared to 5 m in-line (Figure S1); a box-and-whisker plot of IFCB cell biovolumes (Figure S2); IFCB cell biovolume on six of the sample days during NAAMES 3 comparing in-line data with SML data (Figure S3); alkane-to-hydroxyl ratios from three campaigns (Figure S4) (PDF)

■ AUTHOR INFORMATION

Corresponding Author

Lynn M. Russell – Scripps Institution of Oceanography, University of California, San Diego, La Jolla, California 92037, United States; orcid.org/0000-0002-6108-2375; Email: lmrussell@ucsd.edu

Authors

Savannah L. Lewis – Scripps Institution of Oceanography, University of California, San Diego, La Jolla, California 92037, United States; orcid.org/0000-0002-0194-8050

Georges Saliba – California Air Resource Board, Sacramento, California 95814, United States; orcid.org/0000-0002-4305-1305

Patricia K. Quinn – Pacific Marine Environmental Laboratory, NOAA, Seattle, Washington 98115, United States

Timothy S. Bates – Pacific Marine Environmental Laboratory, NOAA, Seattle, Washington 98115, United States; Cooperative Institute for Climate, Oceans, and Ecosystem Studies (CICOES), University of Washington, Seattle, Washington 98105, United States

Craig A. Carlson – Department of Ecology, Evolution and Marine Biology, University of California, Santa Barbara, Santa Barbara, California 93117, United States

Nicholas Baetge – Department of Botany and Plant Pathology, Oregon State University, Corvallis, Oregon 97331, United States

Lihini I. Aluwihare – Scripps Institution of Oceanography, University of California, San Diego, La Jolla, California 92037, United States

Emmanuel Boss – School of Marine Sciences, University of Maine, Orono, Maine 04469, United States

Amanda A. Frossard – Department of Chemistry, University of Georgia, Athens, Georgia 30602, United States; orcid.org/0000-0002-5728-0854

Thomas G. Bell – Plymouth Marine Laboratory, Plymouth PL1 3DH, U.K.; Department of Earth System Science, University of California, Irvine, Irvine, California 92697, United States; orcid.org/0000-0002-4108-7048

Michael J. Behrenfeld – Department of Botany and Plant Pathology, Oregon State University, Corvallis, Oregon 97331, United States

Complete contact information is available at:

<https://pubs.acs.org/10.1021/acsearthspacechem.2c00119>

Author Contributions

S.L.L. led the sampling and analysis of both the STXM-NEXAFS and FTIR spectra and the writing of this manuscript; L.M.R. provided a review of results and their interpretation, as well as manuscript editing; G.S., P.K.Q., T.S.B., C.A.C., N.B., L.I.A., E.B., A.A.F., T.G.B., and M.J.B. provided supporting measurements and editorial comments.

Notes

The authors declare no competing financial interest.

All NAAMES data is available at <https://naames.larc.nasa.gov>. Scripps measurements are available at <https://library.ucsd.edu/dc/collection/bb34508432>. Shipboard measurements are available at <https://seabass.gsfc.nasa.gov/>.

■ ACKNOWLEDGMENTS

The authors would like to thank the dedicated officers and crew of the R/V *Atlantis*. The authors would also like to acknowledge Laura Ravelin, Chia-Li Chen, Raghu Betha, Derek Price, Maryam Askari Lamjiri, Derek Coffman, and Lucia Upchurch for their contributions to collecting and reducing data. The authors would also like to thank Nils Haëntjens for his help retrieving and processing the IFCB data. This work was funded by NASA Grant NNX15AE66G, and Savannah Lewis was supported by the National Science Foundation Graduate Research Fellowship. This is PMEL contribution number 5369.

REFERENCES

- (1) Engel, A.; Bange, H. W.; Cunliffe, M.; Burrows, S. M.; Friedrichs, G.; Galgani, L.; Herrmann, H.; Hertkorn, N.; Johnson, M.; Liss, P. S.; et al. The Ocean's Vital Skin: Toward an Integrated Understanding of the Sea Surface Microlayer. *Front. Mar. Sci.* **2017**, *4*, No. 165.
- (2) Cunliffe, M.; Engel, A.; Frka, S.; Gašparović, B.; Guitart, C.; Murrell, J. C.; Salter, M.; Stolle, C.; Upstill-Goddard, R.; Wurl, O. Sea surface microlayers: A unified physicochemical and biological perspective of the air–ocean interface. *Prog. Oceanogr.* **2013**, *109*, 104–116.
- (3) Liss, P.; Duce, R. *The Sea Surface and Global Change*; Cambridge University Press, 1997. Hunter, K. A.; Liss, P. S. The input of organic material to the oceans: air–sea interactions and the organic chemical composition of the sea surface. *Mar. Chem.* **1977**, *5*, 361–379.
- (4) Aller, J. Y.; Radway, J. C.; Kiltath, W. P.; Bothe, D. W.; Wilson, T. W.; Vaillancourt, R. D.; Quinn, P. K.; Coffman, D. J.; Murray, B. J.; Knopf, D. A. Size-resolved characterization of the polysaccharidic and proteinaceous components of sea spray aerosol. *Atmos. Environ.* **2017**, *154*, 331–347.
- (5) Tseng, R.-S.; Viechnicki, J. T.; Skop, R. A.; Brown, J. W. Sea-to-air transfer of surface-active organic compounds by bursting bubbles. *J. Geophys. Res.: Oceans* **1992**, *97*, S201–S206.
- (6) Blanchard, D. C. Sea-to-Air Transport of Surface Active Material. *Science* **1964**, *146*, 396. Facchini, M. C.; Rinaldi, M.; Decesari, S.; Carbone, C.; Finessi, E.; Mircea, M.; Fuzzi, S.; Ceburnis, D.; Flanagan, R.; Nilsson, E. D.; et al. Primary submicron marine aerosol dominated by insoluble organic colloids and aggregates. *Geophys. Res. Lett.* **2008**, *35*, No. L17814.
- (7) Wurl, O.; Holmes, M. The gelatinous nature of the sea-surface microlayer. *Mar. Chem.* **2008**, *110*, 89–97.
- (8) Engel, A.; Galgani, L. The organic sea-surface microlayer in the upwelling region off the coast of Peru and potential implications for air–sea exchange processes. *Biogeosciences* **2016**, *13*, 989–1007.
- (9) Zäncker, B.; Bracher, A.; Röttgers, R.; Engel, A. Variations of the Organic Matter Composition in the Sea Surface Microlayer: A Comparison between Open Ocean, Coastal, and Upwelling Sites Off the Peruvian Coast. *Front. Microbiol.* **2017**, *8*, No. 2369.
- (10) Reinthaler, T.; Sintes, E.; Herndl, G. J. Dissolved organic matter and bacterial production and respiration in the sea-surface microlayer of the open Atlantic and the western Mediterranean Sea. *Limnol. Oceanogr.* **2008**, *53*, 122–136.
- (11) Russell, L. M.; Hawkins, L. N.; Frossard, A. A.; Quinn, P. K.; Bates, T. S. Carbohydrate-like composition of submicron atmospheric particles and their production from ocean bubble bursting. *Proc. Natl. Acad. Sci. U.S.A.* **2010**, *107*, 6652–6657.
- (12) Frossard, A. A.; Russell, L. M.; Burrows, S. M.; Elliott, S. M.; Bates, T. S.; Quinn, P. K. Sources and composition of submicron organic mass in marine aerosol particles. *J. Geophys. Res.: Atmos.* **2014**, *119*, 12977–913003.
- (13) Hawkins, L. N.; Russell, L. M. Polysaccharides, Proteins, and Phytoplankton Fragments: Four Chemically Distinct Types of Marine Primary Organic Aerosol Classified by Single Particle Spectromicroscopy. *Adv. Meteorol.* **2010**, *2010*, No. 612132.
- (14) Saliba, G.; Sanchez, K. J.; Russell, L. M.; Twohy, C. H.; Roberts, G. C.; Lewis, S.; Dedrick, J.; McCluskey, C. S.; Moore, K.; DeMott, P. J.; Toohey, D. W. Organic composition of three different size ranges of aerosol particles over the Southern Ocean. *Aerosol Sci. Technol.* **2021**, *55*, 268–288.
- (15) Bates, T. S.; Quinn, P. K.; Coffman, D. J.; Johnson, J. E.; Upchurch, L.; Saliba, G.; Lewis, S.; Graff, J.; Russell, L. M.; Behrenfeld, M. J. Variability in Marine Plankton Ecosystems Are Not Observed in Freshly Emitted Sea Spray Aerosol over the North Atlantic Ocean. *Geophys. Res. Lett.* **2020**, *47*, No. e2019GL085938.
- (16) Quinn, P. K.; Bates, T. S.; Schulz, K. S.; Coffman, D. J.; Frossard, A. A.; Russell, L. M.; Keene, W. C.; Kieber, D. J. Contribution of sea surface carbon pool to organic matter enrichment in sea spray aerosol. *Nat. Geosci.* **2014**, *7*, 228–232.
- (17) Bates, T. S.; Quinn, P. K.; Frossard, A. A.; Russell, L. M.; Hakala, J.; Petaja, T.; Kulmala, M.; Covert, D. S.; Cappa, C. D.; Li, S. M.; et al. Measurements of ocean derived aerosol off the coast of California. *J. Geophys. Res.: Atmos.* **2012**, *117*, No. D00V15.
- (18) Beaupré, S. R.; Kieber, D. J.; Keene, W. C.; Long, M. S.; Maben, J. R.; Lu, X.; Zhu, Y.; Frossard, A. A.; Kinsey, J. D.; Duplessis, P.; et al. Oceanic efflux of ancient marine dissolved organic carbon in primary marine aerosol. *Sci. Adv.* **2019**, *5*, No. eaax6535.
- (19) Hansell, D. A.; Carlson, C. A.; Repeta, D. J.; Schlitzer, R. Dissolved Organic Matter in the Ocean: A Controversy Stimulates New Insights. *Oceanography* **2009**, *22*, 202–211.
- (20) Lewis, S. L.; Saliba, G.; Russell, L. M.; Quinn, P. K.; Bates, T. S.; Behrenfeld, M. J. Seasonal Differences in Submicron Marine Aerosol Particle Organic Composition in the North Atlantic. *Front. Mar. Sci.* **2021**, *8*, No. 720208.
- (21) Carlson, C. A.; Hansell, D. A. DOM Sources, Sinks, Reactivity, and Budgets. In *Biogeochemistry of Marine Dissolved Organic Matter*, 2nd ed.; Hansell, D. A., Carlson, C. A., Eds.; Academic Press, 2015; Chapter 3, pp 65–126.
- (22) van Pinxteren, M.; Barthel, S.; Fomba, K. W.; Müller, K.; von Tümpling, W.; Herrmann, H. The influence of environmental drivers on the enrichment of organic carbon in the sea surface microlayer and in submicron aerosol particles – measurements from the Atlantic Ocean. *Elementa* **2017**, *5*, No. 35.
- (23) Gašparović, B.; Plavšić, M.; Čosović, B.; Reigstad, M. Organic matter characterization and fate in the sub-arctic Norwegian fjords during the late spring/summer period. *Estuarine, Coastal Shelf Sci.* **2005**, *62*, 95–107.
- (24) Behrenfeld, M. J.; Moore, R. H.; Hostetler, C. A.; Graff, J.; Gaube, P.; Russell, L. M.; Chen, G.; Doney, S. C.; Giovannoni, S.; Liu, H.; et al. The North Atlantic Aerosol and Marine Ecosystem Study (NAAMES): Science Motive and Mission Overview. *Front. Mar. Sci.* **2019**, *6*, No. 122.
- (25) Garrett, W. D. The organic chemical composition of the ocean surface. *Deep-Sea Res. Oceanogr. Abstr.* **1967**, *14*, 221–227.
- (26) Cunliffe, M.; Wurl, O. *Guide to Best Practices to Study the Ocean's Surface*; Marine Biological Association of the United Kingdom for SCOR, 2014; p 118.
- (27) Chase, A. P.; Kramer, S. J.; Haëntjens, N.; Boss, E. S.; Karp-Boss, L.; Edmondson, M.; Graff, J. R. Evaluation of diagnostic pigments to estimate phytoplankton size classes. *Limnol. Oceanogr.: Methods* **2020**, *18*, 570–584. Moberg, E. A.; Sosik, H. M. Distance maps to estimate cell volume from two-dimensional plankton images. *Limnol. Oceanogr.: Methods* **2012**, *10*, 278–288.
- (28) Takahama, S.; Johnson, A.; Russell, L. M. Quantification of Carboxylic and Carbonyl Functional Groups in Organic Aerosol Infrared Absorbance Spectra. *Aerosol Sci. Technol.* **2013**, *47*, 310–325.
- (29) Frossard, A. A.; Russell, L. M. Removal of Sea Salt Hydrate Water from Seawater-Derived Samples by Dehydration. *Environ. Sci. Technol.* **2012**, *46*, 13326–13333.
- (30) Maria, S. F.; Russell, L. M.; Turpin, B. J.; Porcja, R. J. FTIR measurements of functional groups and organic mass in aerosol samples over the Caribbean. *Atmos. Environ.* **2002**, *36*, 5185–5196. Russell, L. M.; Takahama, S.; Liu, S.; Hawkins, L. N.; Covert, D. S.; Quinn, P. K.; Bates, T. S. Oxygenated fraction and mass of organic aerosol from direct emission and atmospheric processing measured on the R/V Ronald Brown during TEXAQS/GoMACCS 2006. *J. Geophys. Res.: Atmos.* **2009**, *114*, No. D00F05.
- (31) Russell, L. M.; Bahadur, R.; Hawkins, L. N.; Allan, J.; Baumgardner, D.; Quinn, P. K.; Bates, T. S. Organic aerosol characterization by complementary measurements of chemical bonds and molecular fragments. *Atmos. Environ.* **2009**, *43*, 6100–6105. Russell, L. M. Aerosol Organic-Mass-to-Organic-Carbon Ratio Measurements. *Environ. Sci. Technol.* **2003**, *37*, 2982–2987.
- (32) Wan, K. X.; Vidavsky, I.; Gross, M. L. Comparing similar spectra: From similarity index to spectral contrast angle. *J. Am. Soc. Mass Spectrom.* **2002**, *13*, 85–88. Stein, S. E.; Scott, D. R. Optimization and testing of mass spectral library search algorithms for compound identification. *J. Am. Soc. Mass Spectrom.* **1994**, *5*, 859–866. Liu, J.; Russell, L. M.; Lee, A. K. Y.; McKinney, K. A.; Surratt, J. D.; Ziemann, P. J. Observational evidence for pollution-induced

selective uptake contributing to biogenic secondary organic aerosols in the southeastern U.S. *Geophys. Res. Lett.* **2017**, *44*, 8056–8064.

(33) Fox, J.; Behrenfeld, M. J.; Haëntjens, N.; Chase, A.; Kramer, S. J.; Boss, E.; Karp-Boss, L.; Fisher, N. L.; Penta, W. B.; Westberry, T. K.; Halsey, K. H. Phytoplankton Growth and Productivity in the Western North Atlantic: Observations of Regional Variability From the NAAMES Field Campaigns. *Front. Mar. Sci.* **2020**, *7*, No. 24.

(34) Boss, E.; Twardowski, M.; Herring, S. Shape of the particulate beam attenuation spectrum and its inversion to obtain the shape of the particulate size distribution. *Appl. Opt.* **2001**, *40*, 4885–4893. Cetinić, I.; Perry, M. J.; Briggs, N. T.; Kallin, E.; D'Asaro, E. A.; Lee, C. M. Particulate organic carbon and inherent optical properties during 2008 North Atlantic Bloom Experiment. *J. Geophys. Res.: Oceans* **2012**, *117*, No. C06028.

(35) Crilley, L. R.; Ayoko, G. A.; Morawska, L. Analysis of organic aerosols collected on filters by Aerosol Mass Spectrometry for source identification. *Anal. Chim. Acta* **2013**, *803*, 91–96.

(36) Maria, S. F.; Russell Lynn, M.; Gilles Mary, K.; Myneni Satish, C. B. Organic Aerosol Growth Mechanisms and Their Climate-Forcing Implications. *Science* **2004**, *306*, 1921–1924. Russell, L. M.; Maria, S. F.; Myneni, S. C. B. Mapping organic coatings on atmospheric particles. *Geophys. Res. Lett.* **2002**, *29*, 26-21–26-24.

(37) Takahama, S.; Liu, S.; Russell, L. M. Coatings and clusters of carboxylic acids in carbon-containing atmospheric particles from spectromicroscopy and their implications for cloud-nucleating and optical properties. *J. Geophys. Res.: Atmos.* **2010**, *115*, No. D01202.

(38) Gagosian, R. B.; Stuermer, D. H. The cycling of biogenic compounds and their diagenetically transformed products in seawater. *Mar. Chem.* **1977**, *5*, 605–632. Hertkorn, N.; Benner, R.; Frommberger, M.; Schmitt-Kopplin, P.; Witt, M.; Kaiser, K.; Kettrup, A.; Hedges, J. I. Characterization of a major refractory component of marine dissolved organic matter. *Geochim. Cosmochim. Acta* **2006**, *70*, 2990–3010.

(39) Saliba, G.; Chen, C.-L.; Lewis, S.; Russell, L. M.; Quinn, P. K.; Bates, T. S.; Bell, T. G.; Lawler, M. J.; Saltzman, E. S.; Sanchez, K. J.; et al. Seasonal Differences and Variability of Concentrations, Chemical Composition, and Cloud Condensation Nuclei of Marine Aerosol Over the North Atlantic. *J. Geophys. Res.: Atmos.* **2020**, *125*, No. e2020JD033145.

(40) Baetge, N.; Behrenfeld, M. J.; Fox, J.; Halsey, K. H.; Mojica, K. D. A.; Novoa, A.; Stephens, B. M.; Carlson, C. A. The Seasonal Flux and Fate of Dissolved Organic Carbon Through Bacterioplankton in the Western North Atlantic. *Front. Microbiol.* **2021**, *12*, No. 669883.

(41) Takahama, S.; Gilardoni, S.; Russell, L. M.; Kilcoyne, A. L. D. Classification of multiple types of organic carbon composition in atmospheric particles by scanning transmission X-ray microscopy analysis. *Atmos. Environ.* **2007**, *41*, 9435–9451.

(42) Carlson, D. J.; Mayer, L. M. Enrichment of dissolved phenolic material in the surface microlayer of coastal waters. *Nature* **1980**, *286*, 482–483. Knulst, J. C.; Boerschke, R. C.; Loemo, S. Differences in Organic Surface Microlayers from an Artificially Acidified and Control Lake, Elucidated by XAD-8/XAD-4 Tandem Separation and Solid State ¹³C NMR Spectroscopy. *Environ. Sci. Technol.* **1998**, *32*, 8–12.

(43) Bao, H.; Niggemann, J.; Luo, L.; Dittmar, T.; Kao, S.-J. Aerosols as a source of dissolved black carbon to the ocean. *Nat. Commun.* **2017**, *8*, No. 510.

(44) Hayase, K.; Tsubota, H. Sedimentary humic acid and fulvic acid as surface active substances. *Geochim. Cosmochim. Acta* **1983**, *47*, 947–952.

(45) Drozdowska, V.; Wrobel, I.; Markuszewski, P.; Makuch, P.; Raczkowska, A.; Kowalczyk, P. Study on organic matter fractions in the surface microlayer in the Baltic Sea by spectrophotometric and spectrofluorometric methods. *Ocean Sci.* **2017**, *13*, 633–647.

(46) Laskin, A.; Moffet, R. C.; Gilles, M. K.; Fast, J. D.; Zaveri, R. A.; Wang, B.; Nigge, P.; Shutthanandan, J. Tropospheric chemistry of internally mixed sea salt and organic particles: Surprising reactivity of NaCl with weak organic acids. *J. Geophys. Res.: Atmos.* **2012**, *117*, No. D15302.

(47) Vedrinskii, R. V.; Bugaev, L. A.; Gegusin, I. I.; Kraizman, V. L.; Novakovich, A. A.; Prosandeev, S. A.; Ruus, R. E.; Maiste, A. A.; Elango, M. A. X-ray absorption near edge structure (XANES) for KCl. *Solid State Commun.* **1982**, *44*, 1401–1407.

(48) Aller, J. Y.; Kuznetsova, M. R.; Jahns, C. J.; Kemp, P. F. The sea surface microlayer as a source of viral and bacterial enrichment in marine aerosols. *J. Aerosol Sci.* **2005**, *36*, 801–812.

(49) Obernosterer, I.; Catala, P.; Lami, R.; Caparros, J.; Ras, J.; Bricaud, A.; Dupuy, C.; van Wambeke, F.; Lebaron, P. Biochemical characteristics and bacterial community structure of the sea surface microlayer in the South Pacific Ocean. *Biogeosciences* **2008**, *5*, 693–705.

(50) Hendrickson, B. N.; Brooks, S. D.; Thornton, D. C. O.; Moore, R. H.; Crosbie, E.; Ziemba, L. D.; Carlson, C. A.; Baetge, N.; Mirrielees, J. A.; Alsante, A. N. Role of Sea Surface Microlayer Properties in Cloud Formation. *Front. Mar. Sci.* **2021**, *7*, No. 596225. Kuznetsova, M.; Lee, C.; Aller, J. Characterization of the proteinaceous matter in marine aerosols. *Mar. Chem.* **2005**, *96*, 359–377.

(51) Hasenecz, E. S.; Kaluarachchi, C. P.; Lee, H. D.; Tivanski, A. V.; Stone, E. A. Saccharide Transfer to Sea Spray Aerosol Enhanced by Surface Activity, Calcium, and Protein Interactions. *ACS Earth Space Chem.* **2019**, *3*, 2539–2548.

(52) Chance, R. J.; Hamilton, J. F.; Carpenter, L. J.; Hackenberg, S. C.; Andrews, S. J.; Wilson, T. W. Water-Soluble Organic Composition of the Arctic Sea Surface Microlayer and Association with Ice Nucleation Ability. *Environ. Sci. Technol.* **2018**, *52*, 1817–1826.

(53) Kharbush, J. J.; Close, H. G.; Van Mooy, B. A. S.; Arnosti, C.; Smittenberg, R. H.; Le Moigne, F. A. C.; Mollenhauer, G.; Scholz-Böttcher, B.; Obrecht, I.; Koch, B. P.; et al. Particulate Organic Carbon Deconstructed: Molecular and Chemical Composition of Particulate Organic Carbon in the Ocean. *Front. Mar. Sci.* **2020**, *7*, No. 518.

(54) Rahlff, J.; Stolle, C.; Giebel, H.-A.; Brinkhoff, T.; Ribas-Ribas, M.; Hodapp, D.; Wurl, O. High wind speeds prevent formation of a distinct bacterioneuston community in the sea-surface microlayer. *FEMS Microbiol. Ecol.* **2017**, *93*, No. fix041.